CS70 Discrete Probability

Kelvin Lee

kelvinlee@berkeley.edu

October 26, 2020

Overview

Probabilistic Models

Probability Space

Oiscrete Uniform Probability Space

A **probabilistic model** is a mathematical description of an uncertain situation. The elements of a probabilistic model includes

A **probabilistic model** is a mathematical description of an uncertain situation. The elements of a probabilistic model includes

• sample space Ω : set of all all possible outcomes of an experiment.

A **probabilistic model** is a mathematical description of an uncertain situation. The elements of a probabilistic model includes

- sample space Ω : set of all all possible outcomes of an experiment.
- probability law: assigns to a set A of possible outcomes (event) a nonnegative value P(A) (probability of A) that encodes the knowledge about the likelihood of the elements of A.

Some basic terminologies:

Some basic terminologies:

Experiment:

Some basic terminologies:

Experiment:

An **experiment** is a procedure that yields one of a given set of possible outcomes.

Some basic terminologies:

Experiment:

An **experiment** is a procedure that yields one of a given set of possible outcomes.

Sample space Ω :

Some basic terminologies:

Experiment:

An **experiment** is a procedure that yields one of a given set of possible outcomes.

Sample space Ω :

The **sample space** of the experiment is the set of possible outcomes.

Some basic terminologies:

Experiment:

An **experiment** is a procedure that yields one of a given set of possible outcomes.

Sample space Ω :

The **sample space** of the experiment is the set of possible outcomes.

Sample point ω :

4/8

Some basic terminologies:

Experiment:

An **experiment** is a procedure that yields one of a given set of possible outcomes.

Sample space Ω :

The **sample space** of the experiment is the set of possible outcomes.

Sample point ω :

A **sample point** is an element of the sample space.

Some basic terminologies:

Experiment:

An **experiment** is a procedure that yields one of a given set of possible outcomes.

Sample space Ω :

The **sample space** of the experiment is the set of possible outcomes.

Sample point ω :

A **sample point** is an element of the sample space.

Event:

4/8

Some basic terminologies:

Experiment:

An **experiment** is a procedure that yields one of a given set of possible outcomes.

Sample space Ω :

The **sample space** of the experiment is the set of possible outcomes.

Sample point ω :

A **sample point** is an element of the sample space.

Event:

An **event** is a subset of the sample space.

The **probability space** is defined by the triple (Ω, \mathscr{A}, P) where Ω is the sample space, $\mathscr{A} \subseteq \Omega$ is the event space and P is the probability function, satisfying the following axioms:

The **probability space** is defined by the triple (Ω, \mathcal{A}, P) where Ω is the sample space, $\mathcal{A} \subseteq \Omega$ is the event space and P is the probability function, satisfying the following axioms:

Nonnegativity:

The **probability space** is defined by the triple (Ω, \mathcal{A}, P) where Ω is the sample space, $\mathcal{A} \subseteq \Omega$ is the event space and P is the probability function, satisfying the following axioms:

• Nonnegativity: $P(\omega) \ge 0$ for all sample points $\omega \in \Omega$.

The **probability space** is defined by the triple (Ω, \mathcal{A}, P) where Ω is the sample space, $\mathcal{A} \subseteq \Omega$ is the event space and P is the probability function, satisfying the following axioms:

- Nonnegativity: $P(\omega) \ge 0$ for all sample points $\omega \in \Omega$.
- Additivity:

The **probability space** is defined by the triple (Ω, \mathscr{A}, P) where Ω is the sample space, $\mathscr{A} \subseteq \Omega$ is the event space and P is the probability function, satisfying the following axioms:

- Nonnegativity: $P(\omega) \ge 0$ for all sample points $\omega \in \Omega$.
- Additivity: any countable sequence of disjoint sets (mutually exclusive events) E_1, E_2, \ldots satisfies

$$P\left(\bigcup_{i=1}^{\infty}E_{i}\right)=\sum_{i=1}^{\infty}P\left(E_{i}\right).$$

The **probability space** is defined by the triple (Ω, \mathscr{A}, P) where Ω is the sample space, $\mathscr{A} \subseteq \Omega$ is the event space and P is the probability function, satisfying the following axioms:

- Nonnegativity: $P(\omega) \ge 0$ for all sample points $\omega \in \Omega$.
- Additivity: any countable sequence of disjoint sets (mutually exclusive events) E_1, E_2, \ldots satisfies

$$P\left(\bigcup_{i=1}^{\infty}E_{i}\right)=\sum_{i=1}^{\infty}P\left(E_{i}\right).$$

Normalization:

The **probability space** is defined by the triple (Ω, \mathscr{A}, P) where Ω is the sample space, $\mathscr{A} \subseteq \Omega$ is the event space and P is the probability function, satisfying the following axioms:

- Nonnegativity: $P(\omega) \ge 0$ for all sample points $\omega \in \Omega$.
- Additivity: any countable sequence of disjoint sets (mutually exclusive events) E_1, E_2, \ldots satisfies

$$P\left(\bigcup_{i=1}^{\infty}E_{i}\right)=\sum_{i=1}^{\infty}P\left(E_{i}\right).$$

Normalization:

$$\sum_{\omega \in \Omega} P(\omega) = P(\Omega) = 1.$$



Let's define probability.

Let's define probability.

For any event $A \subseteq \Omega$, we define the **probability** of A to be

Let's define probability.

For any event $A \subseteq \Omega$, we define the **probability** of A to be

$$P(A) = \sum_{\omega \in A} P(\omega).$$

Let's define probability.

For any event $A \subseteq \Omega$, we define the **probability** of A to be

$$P(A) = \sum_{\omega \in A} P(\omega).$$

Let's define probability.

For any event $A \subseteq \Omega$, we define the **probability** of A to be

$$P(A) = \sum_{\omega \in A} P(\omega).$$

•
$$P(\emptyset) = 0$$
.

Let's define probability.

For any event $A \subseteq \Omega$, we define the **probability** of A to be

$$P(A) = \sum_{\omega \in A} P(\omega).$$

- $P(\emptyset) = 0$.
- $P(\overline{A}) = 1 P(A)$, where \overline{A} (or A^c) is the **complement** of A.

Let's define probability.

For any event $A \subseteq \Omega$, we define the **probability** of A to be

$$P(A) = \sum_{\omega \in A} P(\omega).$$

- $P(\emptyset) = 0$.
- $P(\overline{A}) = 1 P(A)$, where \overline{A} (or A^c) is the **complement** of A.
- $P(A \cup B) = P(A) + P(A) P(A \cap B)$.

Let's define probability.

For any event $A \subseteq \Omega$, we define the **probability** of A to be

$$P(A) = \sum_{\omega \in A} P(\omega).$$

- $P(\emptyset) = 0$.
- $P(\overline{A}) = 1 P(A)$, where \overline{A} (or A^c) is the **complement** of A.
- $P(A \cup B) = P(A) + P(A) P(A \cap B)$.
- If $A \subseteq B$, then $P(A) \le P(B)$.



Discrete Uniform Probability Law:

Discrete Uniform Probability Law:

In a **uniform** probability space, all sample points have the same probability $\frac{1}{|\Omega|}$ and the probability of an event A is

Discrete Uniform Probability Law:

In a **uniform** probability space, all sample points have the same probability $\frac{1}{|\Omega|}$ and the probability of an event A is

$$P(A) = \frac{|A|}{|\Omega|}.$$

Discrete Uniform Probability Law:

In a **uniform** probability space, all sample points have the same probability $\frac{1}{|\Omega|}$ and the probability of an event A is

$$P(A) = \frac{|A|}{|\Omega|}.$$

For uniform spaces, computing probabilities is simply counting sample points.

Problem Time!