

# CS70 Counting

Kelvin Lee

*kelvinlee@berkeley.edu*

October 7, 2020

# Overview

- 1 Rules of Counting
- 2 Stars and Bars
- 3 Binomial Theorem
- 4 Combinatorial Proofs
- 5 Principle of Inclusion-Exclusion
- 6 Problems
- 7 Summary/Tips

# First Rule of Counting

# First Rule of Counting

## First Rule of Counting(Product Rule):

# First Rule of Counting

## First Rule of Counting(Product Rule):

If there are  $n$  ways of doing something, and  $m$  ways of doing another thing after that, then there are  $n \times m$  ways to perform both of these actions.

# First Rule of Counting

## First Rule of Counting(Product Rule):

If there are  $n$  ways of doing something, and  $m$  ways of doing another thing after that, then there are  $n \times m$  ways to perform both of these actions.

- **Order matters** (permutations).

# First Rule of Counting

## First Rule of Counting(Product Rule):

If there are  $n$  ways of doing something, and  $m$  ways of doing another thing after that, then there are  $n \times m$  ways to perform both of these actions.

- **Order matters** (permutations).
- Sampling  $k$  elements from  $n$  items:

# First Rule of Counting

## First Rule of Counting(Product Rule):

If there are  $n$  ways of doing something, and  $m$  ways of doing another thing after that, then there are  $n \times m$  ways to perform both of these actions.

- **Order matters** (permutations).
- Sampling  $k$  elements from  $n$  items:
  - **With replacement:**



# First Rule of Counting

## First Rule of Counting(Product Rule):

If there are  $n$  ways of doing something, and  $m$  ways of doing another thing after that, then there are  $n \times m$  ways to perform both of these actions.

- **Order matters** (permutations).
- Sampling  $k$  elements from  $n$  items:
  - **With replacement:**  $n^k$ .

# First Rule of Counting

## First Rule of Counting(Product Rule):

If there are  $n$  ways of doing something, and  $m$  ways of doing another thing after that, then there are  $n \times m$  ways to perform both of these actions.

- **Order matters** (permutations).
- Sampling  $k$  elements from  $n$  items:
  - **With replacement:**  $n^k$ .
  - **Without replacement:**

# First Rule of Counting

## First Rule of Counting(Product Rule):

If there are  $n$  ways of doing something, and  $m$  ways of doing another thing after that, then there are  $n \times m$  ways to perform both of these actions.

- **Order matters** (permutations).
- Sampling  $k$  elements from  $n$  items:
  - **With replacement:**  $n^k$ .
  - **Without replacement:**  $\frac{n!}{(n-k)!}$

# Second Rule of Counting

# Second Rule of Counting

## Second Rule of Counting:

# Second Rule of Counting

## Second Rule of Counting:

If order doesn't matter count ordered objects and then divide by number of orderings.

# Second Rule of Counting

## Second Rule of Counting:

If order doesn't matter count ordered objects and then divide by number of orderings.

- **Without replacement** and **ordering doesn't matter** (combinations).

# Second Rule of Counting

## Second Rule of Counting:

If order doesn't matter count ordered objects and then divide by number of orderings.

- **Without replacement** and **ordering doesn't matter** (combinations).
- Number of ways of choosing  $k$ -element subsets out of a set of size  $n$ :



# Second Rule of Counting

## Second Rule of Counting:

If order doesn't matter count ordered objects and then divide by number of orderings.

- **Without replacement** and **ordering doesn't matter** (combinations).
- Number of ways of choosing  $k$ -element subsets out of a set of size  $n$ :

$$\binom{n}{k} = \frac{n!}{(n-k)!k!}.$$

# Starts and Bars

# Starts and Bars

**Problem:** Consider the equation  $a + b + c + d = 12$  where  $a, b, c, d$  are non-negative integers. How many solutions are there to this equation?

# Starts and Bars

**Problem:** Consider the equation  $a + b + c + d = 12$  where  $a, b, c, d$  are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 stars and 3 bars.

# Starts and Bars

**Problem:** Consider the equation  $a + b + c + d = 12$  where  $a, b, c, d$  are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 stars and 3 bars.

★★ | ★★ | ★★★ | ★★★★★

# Starts and Bars

**Problem:** Consider the equation  $a + b + c + d = 12$  where  $a, b, c, d$  are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 stars and 3 bars.

★★ | ★★ | ★★★ | ★★★★★

- How many ways can we arrange them?

# Starts and Bars

**Problem:** Consider the equation  $a + b + c + d = 12$  where  $a, b, c, d$  are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 stars and 3 bars.

★ ★ | ★ ★ | ★ ★ ★ | ★ ★ ★ ★

- How many ways can we arrange them?  $\binom{12+3}{3}$

# Starts and Bars

**Problem:** Consider the equation  $a + b + c + d = 12$  where  $a, b, c, d$  are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 stars and 3 bars.

★★ | ★★ | ★★★ | ★★★★★

- How many ways can we arrange them?  $\binom{12+3}{3} = \binom{15}{3}$



# Starts and Bars

**Problem:** Consider the equation  $a + b + c + d = 12$  where  $a, b, c, d$  are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 stars and 3 bars.

★ ★ | ★ ★ | ★ ★ ★ | ★ ★ ★ ★

- How many ways can we arrange them?  $\binom{12+3}{3} = \binom{15}{3}$
- This is the answer to our original problem! Do you see the bijection between the two problems?

# Starts and Bars

**Problem:** Consider the equation  $a + b + c + d = 12$  where  $a, b, c, d$  are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 stars and 3 bars.

★★ | ★★ | ★★★ | ★★★★★

- How many ways can we arrange them?  $\binom{12+3}{3} = \binom{15}{3}$
- This is the answer to our original problem! Do you see the bijection between the two problems?

**Zeroth Rule of Counting:**

# Starts and Bars

**Problem:** Consider the equation  $a + b + c + d = 12$  where  $a, b, c, d$  are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 stars and 3 bars.

★★ | ★★ | ★★★ | ★★★★★

- How many ways can we arrange them?  $\binom{12+3}{3} = \binom{15}{3}$
- This is the answer to our original problem! Do you see the bijection between the two problems?

## Zeroth Rule of Counting:

If a set  $A$  has a bijection relationship with a set  $B$ , then  $|A| = |B|$ .

# Starts and Bars

# Stars and Bars

## Stars and Bars:

# Starts and Bars

## Stars and Bars:

How many ways can one distribute  $n$  indistinguishable objects into  $k$  distinguishable bins?

# Stars and Bars

## Stars and Bars:

How many ways can one distribute  $n$  indistinguishable objects into  $k$  distinguishable bins?

$$\binom{n + k - 1}{k - 1}.$$

# Stars and Bars

## Stars and Bars:

How many ways can one distribute  $n$  indistinguishable objects into  $k$  distinguishable bins?

$$\binom{n + k - 1}{k - 1}.$$

- Useful for **with replacement** but **order doesn't matter** type of problems.



# Binomial Theorem

# Binomial Theorem

## Binomial Theorem:

# Binomial Theorem

## Binomial Theorem:

For all  $n \in \mathbb{N}$ ,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

# Binomial Theorem

## Binomial Theorem:

For all  $n \in \mathbb{N}$ ,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

***Proof:***

# Binomial Theorem

## Binomial Theorem:

For all  $n \in \mathbb{N}$ ,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

***Proof:***

See notes. □

# Binomial Theorem

## Binomial Theorem:

For all  $n \in \mathbb{N}$ ,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

***Proof:***

See notes. □

**Corollary:**

# Binomial Theorem

## Binomial Theorem:

For all  $n \in \mathbb{N}$ ,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

***Proof:***

See notes. □

**Corollary:**

$$\sum_{k=0}^n (-1)^k \binom{n}{k} = 0.$$

# Binomial Theorem

## Binomial Theorem:

For all  $n \in \mathbb{N}$ ,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

***Proof:***

See notes. □

## Corollary:

$$\sum_{k=0}^n (-1)^k \binom{n}{k} = 0.$$

***Proof:***



# Binomial Theorem

## Binomial Theorem:

For all  $n \in \mathbb{N}$ ,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

**Proof:**

See notes. □

## Corollary:

$$\sum_{k=0}^n (-1)^k \binom{n}{k} = 0.$$

**Proof:**

Plug in  $a = -1$  and  $b = 1$  for the binomial theorem.

# Binomial Theorem

## Binomial Theorem:

For all  $n \in \mathbb{N}$ ,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

**Proof:**

See notes. □

## Corollary:

$$\sum_{k=0}^n (-1)^k \binom{n}{k} = 0.$$

**Proof:**

Plug in  $a = -1$  and  $b = 1$  for the binomial theorem. □

# Combinatorial Proofs

# Combinatorial Proofs

- Intuitive counting arguments. No tedious algebraic manipulation.

# Combinatorial Proofs

- Intuitive counting arguments. No tedious algebraic manipulation.
- Proofs by stories: same story from multiple perspectives.

# Combinatorial Proofs

- Intuitive counting arguments. No tedious algebraic manipulation.
- Proofs by stories: same story from multiple perspectives.
- Proving an identity by counting the same thing in two different ways.

# Combinatorial Proofs

- Intuitive counting arguments. No tedious algebraic manipulation.
- Proofs by stories: same story from multiple perspectives.
- Proving an identity by counting the same thing in two different ways.
- Useful identity:

# Combinatorial Proofs

- Intuitive counting arguments. No tedious algebraic manipulation.
- Proofs by stories: same story from multiple perspectives.
- Proving an identity by counting the same thing in two different ways.
- Useful identity:

$$\binom{n}{k} = \binom{n}{n-k}.$$



# Combinatorial Proofs

- Intuitive counting arguments. No tedious algebraic manipulation.
- Proofs by stories: same story from multiple perspectives.
- Proving an identity by counting the same thing in two different ways.
- Useful identity:

$$\binom{n}{k} = \binom{n}{n-k}.$$

- Choosing  $k$  objects to include is equivalent to choosing  $n - k$  objects to exclude.

# Example

## Combinatorial Identity:

## Combinatorial Identity:

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

# Example

## Combinatorial Identity:

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

***Proof:***

# Example

## Combinatorial Identity:

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

### ***Proof:***

Although we can use binomial theorem by letting  $a = b = 1$ , we use combinatorial argument to prove this.

# Example

## Combinatorial Identity:

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

### ***Proof:***

Although we can use binomial theorem by letting  $a = b = 1$ , we use combinatorial argument to prove this.

### **RHS:**

# Example

## Combinatorial Identity:

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

### ***Proof:***

Although we can use binomial theorem by letting  $a = b = 1$ , we use combinatorial argument to prove this.

**RHS:** Total number of subsets of a set of size  $n$ .



# Example

## Combinatorial Identity:

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

### ***Proof:***

Although we can use binomial theorem by letting  $a = b = 1$ , we use combinatorial argument to prove this.

**RHS:** Total number of subsets of a set of size  $n$ .

**LHS:**

# Example

## Combinatorial Identity:

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

### ***Proof:***

Although we can use binomial theorem by letting  $a = b = 1$ , we use combinatorial argument to prove this.

**RHS:** Total number of subsets of a set of size  $n$ .

**LHS:** The number of ways to choose a subset of size  $i$  is  $\binom{n}{i}$ . To find the total number of subsets, we simply add all the cases when  $i = 0, 1, 2, \dots, n$ .

□

# Principle of Inclusion-Exclusion

# Principle of Inclusion-Exclusion

## Principle of Inclusion-Exclusion(General):

# Principle of Inclusion-Exclusion

## Principle of Inclusion-Exclusion(General):

Let  $A_1, \dots, A_n$  be arbitrary subsets of the same finite set  $A$ . Then,

$$|A_1 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k-1} \sum_{S \subseteq \{1, \dots, n\}: |S|=k} |\cap_{i \in S} A_i|.$$

# Principle of Inclusion-Exclusion

## Principle of Inclusion-Exclusion(General):

Let  $A_1, \dots, A_n$  be arbitrary subsets of the same finite set  $A$ . Then,

$$|A_1 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k-1} \sum_{S \subseteq \{1, \dots, n\}: |S|=k} |\cap_{i \in S} A_i|.$$

***Proof:***

# Principle of Inclusion-Exclusion

## Principle of Inclusion-Exclusion(General):

Let  $A_1, \dots, A_n$  be arbitrary subsets of the same finite set  $A$ . Then,

$$|A_1 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k-1} \sum_{S \subseteq \{1, \dots, n\}: |S|=k} |\cap_{i \in S} A_i|.$$

***Proof:***

See notes.



# Principle of Inclusion-Exclusion

## Principle of Inclusion-Exclusion(General):

Let  $A_1, \dots, A_n$  be arbitrary subsets of the same finite set  $A$ . Then,

$$|A_1 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k-1} \sum_{S \subseteq \{1, \dots, n\}: |S|=k} |\cap_{i \in S} A_i|.$$

### ***Proof:***

See notes. □

## Principle of Inclusion-Exclusion(Simplified):



# Principle of Inclusion-Exclusion

## Principle of Inclusion-Exclusion(General):

Let  $A_1, \dots, A_n$  be arbitrary subsets of the same finite set  $A$ . Then,

$$|A_1 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k-1} \sum_{S \subseteq \{1, \dots, n\}: |S|=k} |\cap_{i \in S} A_i|.$$

### ***Proof:***

See notes. □

## Principle of Inclusion-Exclusion(Simplified):

$$|A \cup B| = |A| + |B| - |A \cap B|.$$

# Problems

## SP19 MT2 6.3

## SP19 MT2 6.3

How many permutations of the numbers 1 through  $n$  are there such that 1 comes before 2 and after 3? Assume  $n > 3$ .

## SP19 MT2 6.3

How many permutations of the numbers 1 through  $n$  are there such that 1 comes before 2 and after 3? Assume  $n > 3$ .

**Hint:**

## SP19 MT2 6.3

How many permutations of the numbers 1 through  $n$  are there such that 1 comes before 2 and after 3? Assume  $n > 3$ .

**Hint:** how many ways to choose the positions for the three numbers? What do we do with the remaining numbers?

## SP19 MT2 6.3

How many permutations of the numbers 1 through  $n$  are there such that 1 comes before 2 and after 3? Assume  $n > 3$ .

**Hint:** how many ways to choose the positions for the three numbers? What do we do with the remaining numbers?

**Solution:**

## SP19 MT2 6.3

How many permutations of the numbers 1 through  $n$  are there such that 1 comes before 2 and after 3? Assume  $n > 3$ .

**Hint:** how many ways to choose the positions for the three numbers? What do we do with the remaining numbers?

### Solution:

- There are  $\binom{n}{3}$  ways to pick positions for 1, 2, 3. For the positions picked, we place the three numbers in a way such that the conditions are met, i.e, we place them in the order of 3, 1, 2.



## SP19 MT2 6.3

How many permutations of the numbers 1 through  $n$  are there such that 1 comes before 2 and after 3? Assume  $n > 3$ .

**Hint:** how many ways to choose the positions for the three numbers? What do we do with the remaining numbers?

### Solution:

- There are  $\binom{n}{3}$  ways to pick positions for 1, 2, 3. For the positions picked, we place the three numbers in a way such that the conditions are met, i.e, we place them in the order of 3, 1, 2.
- Now for the remaining numbers, there are  $(n - 3)!$  to arrange them.

## SP19 MT2 6.3

How many permutations of the numbers 1 through  $n$  are there such that 1 comes before 2 and after 3? Assume  $n > 3$ .

**Hint:** how many ways to choose the positions for the three numbers? What do we do with the remaining numbers?

### Solution:

- There are  $\binom{n}{3}$  ways to pick positions for 1, 2, 3. For the positions picked, we place the three numbers in a way such that the conditions are met, i.e, we place them in the order of 3, 1, 2.
- Now for the remaining numbers, there are  $(n - 3)!$  to arrange them.
- Finally, by the **first rule of counting**, we have  $\boxed{\frac{n!}{6}}$  permutations.

# Problems

## SP18 MT2 5

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

**Hint:**

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

**Hint:** how many edges are there? How do we choose them?



## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

**Hint:** how many edges are there? How do we choose them?

**Solution:**

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

**Hint:** how many edges are there? How do we choose them?

**Solution:**

- If every vertex has degree 1, then we can only have 3 edges.

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

**Hint:** how many edges are there? How do we choose them?

**Solution:**

- If every vertex has degree 1, then we can only have 3 edges.
- Each edge requires 2 vertices, so  $\binom{6}{2} = 15$  ways to choose an edge.

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

**Hint:** how many edges are there? How do we choose them?

### Solution:

- If every vertex has degree 1, then we can only have 3 edges.
- Each edge requires 2 vertices, so  $\binom{6}{2} = 15$  ways to choose an edge.
- After choosing the first edge, we have 4 vertices remaining, so there are  $\binom{4}{2} = 6$  ways to choose the second edge and similarly  $\binom{2}{2} = 1$  way to choose the final edge.

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

**Hint:** how many edges are there? How do we choose them?

**Solution:**

- If every vertex has degree 1, then we can only have 3 edges.
- Each edge requires 2 vertices, so  $\binom{6}{2} = 15$  ways to choose an edge.
- After choosing the first edge, we have 4 vertices remaining, so there are  $\binom{4}{2} = 6$  ways to choose the second edge and similarly  $\binom{2}{2} = 1$  way to choose the final edge.
- However, since order doesn't matter, by the **second rule of counting**, we divide by  $3! = 6$ . So our final answer is 15.

# Problems

## SP18 MT2 5

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.



## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(b) How many ways can we form two disjoint cycles of length 3 with 6 vertices?

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(b) How many ways can we form two disjoint cycles of length 3 with 6 vertices?

**Hint:**

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(b) How many ways can we form two disjoint cycles of length 3 with 6 vertices?

**Hint:** how many ways to pick two groups?

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(b) How many ways can we form two disjoint cycles of length 3 with 6 vertices?

**Hint:** how many ways to pick two groups?

**Solution:**

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(b) How many ways can we form two disjoint cycles of length 3 with 6 vertices?

**Hint:** how many ways to pick two groups?

**Solution:**

- We are choosing two sets of 3 vertices. There are  $\binom{6}{3} \binom{3}{3} = 20$  ways.

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(b) How many ways can we form two disjoint cycles of length 3 with 6 vertices?

**Hint:** how many ways to pick two groups?

**Solution:**

- We are choosing two sets of 3 vertices. There are  $\binom{6}{3} \binom{3}{3} = 20$  ways.
- But order doesn't matter here again. So we divide by  $2!$ . Thus, the answer is 10.

# Problems

## SP18 MT2 5



## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

**Hint:**

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

**Hint:** how many ways can we permute the vertices?

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

**Hint:** how many ways can we permute the vertices?

**Solution:**

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

**Hint:** how many ways can we permute the vertices?

### Solution:

- We think of the cycle as a permutation of the vertices, which has  $6!$  possibilities.

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

**Hint:** how many ways can we permute the vertices?

### Solution:

- We think of the cycle as a permutation of the vertices, which has  $6!$  possibilities.
- However, it doesn't matter where we start, so divide by 6.

## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

**Hint:** how many ways can we permute the vertices?

### Solution:

- We think of the cycle as a permutation of the vertices, which has  $6!$  possibilities.
- However, it doesn't matter where we start, so divide by 6.
- The direction in which we travel along the cycle also doesn't matter, so divide by 2.



## SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

**Hint:** how many ways can we permute the vertices?

**Solution:**

- We think of the cycle as a permutation of the vertices, which has  $6!$  possibilities.
- However, it doesn't matter where we start, so divide by 6.
- The direction in which we travel along the cycle also doesn't matter, so divide by 2.
- Thus, our answer is  $\frac{6!}{2 \cdot 6} = \boxed{60}$ .

# Problems

## SP17 MT2 4

## SP17 MT2 4

What are the number of ways to divide  $m$  dollar bills among  $z$  people?

## SP17 MT2 4

What are the number of ways to divide  $m$  dollar bills among  $z$  people?

**Hint:**

## SP17 MT2 4

What are the number of ways to divide  $m$  dollar bills among  $z$  people?

**Hint:** Stars and bars.

## SP17 MT2 4

What are the number of ways to divide  $m$  dollar bills among  $z$  people?

**Hint:** Stars and bars.

**Solution:**

## SP17 MT2 4

What are the number of ways to divide  $m$  dollar bills among  $z$  people?

**Hint:** Stars and bars.

### Solution:

- This is a stars and bars problem where we have  $z - 1$  bars and  $m$  stars.



## SP17 MT2 4

What are the number of ways to divide  $m$  dollar bills among  $z$  people?

**Hint:** Stars and bars.

### Solution:

- This is a stars and bars problem where we have  $z - 1$  bars and  $m$  stars.
- So  $n = m$  and  $k = z$  in this case.

## SP17 MT2 4

What are the number of ways to divide  $m$  dollar bills among  $z$  people?

**Hint:** Stars and bars.

### Solution:

- This is a stars and bars problem where we have  $z - 1$  bars and  $m$  stars.
- So  $n = m$  and  $k = z$  in this case.

- Thus, the answer is  $\binom{n+k-1}{k-1} = \boxed{\binom{m+z-1}{z-1}}.$

# Problems

## SP16 MT2 6

## SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

## SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

**Hint:**

## SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

**Hint:** Ternary strings.

## SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

**Hint:** Ternary strings.

**Solution:**



## SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

**Hint:** Ternary strings.

**Solution:**

- **LHS:**

## SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

**Hint:** Ternary strings.

**Solution:**

- **LHS:** the number of ternary strings of length  $n$ .

## SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

**Hint:** Ternary strings.

**Solution:**

- **LHS:** the number of ternary strings of length  $n$ .
- **RHS:**

## SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

**Hint:** Ternary strings.

**Solution:**

- **LHS:** the number of ternary strings of length  $n$ .
- **RHS:** There are  $\binom{n}{i}$  positions of the 2's, and there are  $2^{n-i}$  possible patterns of 0 and 1's in the remaining positions. The sum gives you all the ternary strings.

# Summary/Tips

## Summary:

## Summary:

# Summary/Tips

## Summary:

	with replacement	w/o replacement
order matters	$n^k$	$\frac{n!}{(n-k)!}$
order doesn't matter	$\binom{n+k-1}{k-1}$	$\binom{n}{k}$

## Tips:



# Summary/Tips

## Summary:

	with replacement	w/o replacement
order matters	$n^k$	$\frac{n!}{(n-k)!}$
order doesn't matter	$\binom{n+k-1}{k-1}$	$\binom{n}{k}$

## Tips:

- Don't memorize formulas. Understand them by counting.

# Summary/Tips

## Summary:

	with replacement	w/o replacement
order matters	$n^k$	$\frac{n!}{(n-k)!}$
order doesn't matter	$\binom{n+k-1}{k-1}$	$\binom{n}{k}$

## Tips:

- Don't memorize formulas. Understand them by counting.
- Identify which categories does the problem fall under.

# Summary/Tips

## Summary:

	with replacement	w/o replacement
order matters	$n^k$	$\frac{n!}{(n-k)!}$
order doesn't matter	$\binom{n+k-1}{k-1}$	$\binom{n}{k}$

## Tips:

- Don't memorize formulas. Understand them by counting.
- Identify which categories does the problem fall under.
- Double check answers by using two different counting approaches.

# Summary/Tips

## Summary:

	with replacement	w/o replacement
order matters	$n^k$	$\frac{n!}{(n-k)!}$
order doesn't matter	$\binom{n+k-1}{k-1}$	$\binom{n}{k}$

## Tips:

- Don't memorize formulas. Understand them by counting.
- Identify which categories does the problem fall under.
- Double check answers by using two different counting approaches.
- Check for overcounting.

# Summary/Tips

## Summary:

	with replacement	w/o replacement
order matters	$n^k$	$\frac{n!}{(n-k)!}$
order doesn't matter	$\binom{n+k-1}{k-1}$	$\binom{n}{k}$

## Tips:

- Don't memorize formulas. Understand them by counting.
- Identify which categories does the problem fall under.
- Double check answers by using two different counting approaches.
- Check for overcounting.
- Relax and have fun!