

CS70 Modular Arithmetic

Kelvin Lee

kelvinlee@berkeley.edu

September 21, 2020

Overview

- 1 Basic Definitions
- 2 Multiplicative Inverse
- 3 Euclid's Algorithm
- 4 Extended Euclid's algorithm

Basic Definitions

Definition (Congruence)

Basic Definitions

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m
- x and y have the same remainder w.r.t. m

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m
- x and y have the same remainder w.r.t. m
- $x = y + km$ for some integer k

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m
 - x and y have the same remainder w.r.t. m
 - $x = y + km$ for some integer k
-
- In modulo m , only the numbers $\{0, 1, 2, \dots, m - 1\}$ exist.

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m
 - x and y have the same remainder w.r.t. m
 - $x = y + km$ for some integer k
-
- In modulo m , only the numbers $\{0, 1, 2, \dots, m - 1\}$ exist.
 - Division is not well-defined.

Basic Definitions

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m
 - x and y have the same remainder w.r.t. m
 - $x = y + km$ for some integer k
-
- In modulo m , only the numbers $\{0, 1, 2, \dots, m - 1\}$ exist.
 - Division is not well-defined.

Definition (Multiplicative Inverse)

Basic Definitions

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m
 - x and y have the same remainder w.r.t. m
 - $x = y + km$ for some integer k
-
- In modulo m , only the numbers $\{0, 1, 2, \dots, m - 1\}$ exist.
 - Division is not well-defined.

Definition (Multiplicative Inverse)

Normally we say that the **multiplicative inverse** of x is y if $xy = 1$.

Basic Definitions

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y(\text{mod } m)$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m
 - x and y have the same remainder w.r.t. m
 - $x = y + km$ for some integer k
-
- In modulo m , only the numbers $\{0, 1, 2, \dots, m - 1\}$ exist.
 - Division is not well-defined.

Definition (Multiplicative Inverse)

Normally we say that the **multiplicative inverse** of x is y if $xy = 1$. In the modular space, the **multiplicative inverse** of $x \text{ mod } m$ is y if $xy \equiv 1(\text{mod } m)$.

Theorems

Theorem (Modular operations)

Theorem (Modular operations)

$a \equiv c \pmod{m}$ and $b \equiv d \pmod{m} \implies a + b \equiv c + d \pmod{m}$ and $a \cdot b \equiv c \cdot d \pmod{m}$.

Theorem (Modular operations)

$a \equiv c \pmod{m}$ and $b \equiv d \pmod{m} \implies a + b \equiv c + d \pmod{m}$ and $a \cdot b \equiv c \cdot d \pmod{m}$.

Theorem (Existence of multiplicative inverse)

Theorem (Modular operations)

$a \equiv c \pmod{m}$ and $b \equiv d \pmod{m} \implies a + b \equiv c + d \pmod{m}$ and $a \cdot b \equiv c \cdot d \pmod{m}$.

Theorem (Existence of multiplicative inverse)

$\gcd(x, m) = 1 \implies x$ has a multiplicative inverse modulo m and it is **unique**.

Euclid's Algorithm

Euclid's Algorithm

How do we compute \gcd of two numbers x and y ?

Euclid's Algorithm

How do we compute \gcd of two numbers x and y ?

Theorem (Euclid's Algorithm)

Euclid's Algorithm

How do we compute gcd of two numbers x and y ?

Theorem (Euclid's Algorithm)

Let $x \geq y > 0$. Then

$$\gcd(x, y) = \gcd(y, x \bmod y)$$

Euclid's Algorithm

How do we compute gcd of two numbers x and y ?

Theorem (Euclid's Algorithm)

Let $x \geq y > 0$. Then

$$\gcd(x, y) = \gcd(y, x \bmod y)$$

Example

Euclid's Algorithm

How do we compute gcd of two numbers x and y ?

Theorem (Euclid's Algorithm)

Let $x \geq y > 0$. Then

$$\gcd(x, y) = \gcd(y, x \bmod y)$$

Example

Compute $\gcd(16, 10)$:

Euclid's Algorithm

How do we compute gcd of two numbers x and y ?

Theorem (Euclid's Algorithm)

Let $x \geq y > 0$. Then

$$\gcd(x, y) = \gcd(y, x \bmod y)$$

Example

Compute $\gcd(16, 10)$:

$$\begin{aligned}\gcd(16, 10) &= \gcd(10, 6) \\ &= \gcd(6, 4) \\ &= \gcd(4, 2) \\ &= \gcd(2, 0) \\ &= 2.\end{aligned}$$

Extended Euclid's algorithm

Extended Euclid's algorithm

How to compute the multiplicative inverse?

Extended Euclid's algorithm

How to compute the multiplicative inverse?

- Need an algorithm that returns integers a and b such that:

$$\gcd(x, y) = ax + by.$$

Extended Euclid's algorithm

How to compute the multiplicative inverse?

- Need an algorithm that returns integers a and b such that:

$$\gcd(x, y) = ax + by.$$

- When $\gcd(x, y) = 1$, we can deduce that b is an inverse of y mod x .

Extended Euclid's algorithm

How to compute the multiplicative inverse?

- Need an algorithm that returns integers a and b such that:

$$\gcd(x, y) = ax + by.$$

- When $\gcd(x, y) = 1$, we can deduce that b is an inverse of y mod x .
- So basically repetitively substitute values so that the final expression is in terms of x and y .

Problem Time!