

# CS70 Random Variables

Kelvin Lee

*kelvinlee@berkeley.edu*

November 9, 2020

# Overview

- 1 Discrete Random Variables
- 2 Expectation
- 3 Variance
- 4 Bernoulli Distribution
- 5 Binomial Distribution
- 6 Indicator Random Variable
- 7 Geometric Distribution
- 8 Poisson Distribution

# Definitions

# Definitions

**(Random Variable).**

# Definitions

## (Random Variable).

A **random variable**  $X$  on a sample space  $\Omega$  is a function  $X : \Omega \rightarrow \mathbb{R}$  that assigns to each sample point  $\omega \in \Omega$  a real number  $X(\omega)$ .

# Definitions

## (Random Variable).

A **random variable**  $X$  on a sample space  $\Omega$  is a function  $X : \Omega \rightarrow \mathbb{R}$  that assigns to each sample point  $\omega \in \Omega$  a real number  $X(\omega)$ .

## (Distribution).

# Definitions

## (Random Variable).

A **random variable**  $X$  on a sample space  $\Omega$  is a function  $X : \Omega \rightarrow \mathbb{R}$  that assigns to each sample point  $\omega \in \Omega$  a real number  $X(\omega)$ .

## (Distribution).

The **distribution** of a discrete random variable  $X$  is the collection of values  $\{(x, P(X = x)) : x \in \mathcal{X}\}$ , where  $\mathcal{X}$  is the set of all possible values taken by  $X$ .

# Definitions

## (Random Variable).

A **random variable**  $X$  on a sample space  $\Omega$  is a function  $X : \Omega \rightarrow \mathbb{R}$  that assigns to each sample point  $\omega \in \Omega$  a real number  $X(\omega)$ .

## (Distribution).

The **distribution** of a discrete random variable  $X$  is the collection of values  $\{(x, P(X = x)) : x \in \mathcal{X}\}$ , where  $\mathcal{X}$  is the set of all possible values taken by  $X$ .

## (Probability Mass Function).



# Definitions

## (Random Variable).

A **random variable**  $X$  on a sample space  $\Omega$  is a function  $X : \Omega \rightarrow \mathbb{R}$  that assigns to each sample point  $\omega \in \Omega$  a real number  $X(\omega)$ .

## (Distribution).

The **distribution** of a discrete random variable  $X$  is the collection of values  $\{(x, P(X = x)) : x \in \mathcal{X}\}$ , where  $\mathcal{X}$  is the set of all possible values taken by  $X$ .

## (Probability Mass Function).

The **probability mass function**, or **PMF**, of a discrete random variable  $X$  is a function mapping  $X$ 's values to their associated probabilities. It is the function  $p : \mathcal{X} \rightarrow [0, 1]$  defined by

$$p_X(x) := P(X = x).$$

# Definitions

**(Joint Distribution).**

## (Joint Distribution).

The **joint distribution** for two discrete random variables  $X$  and  $Y$  is the collection of values  $\{((x, y), P(X = x, Y = y)) : x \in \mathcal{X}, y \in \mathcal{Y}\}$ , where  $\mathcal{X}$  is the set of all possible values taken by  $X$  and  $\mathcal{Y}$  is the set of all possible values taken by  $Y$ .

# Definitions

## (Joint Distribution).

The **joint distribution** for two discrete random variables  $X$  and  $Y$  is the collection of values  $\{((x, y), P(X = x, Y = y)) : x \in \mathcal{X}, y \in \mathcal{Y}\}$ , where  $\mathcal{X}$  is the set of all possible values taken by  $X$  and  $\mathcal{Y}$  is the set of all possible values taken by  $Y$ .

## (Marginal Distribution).

# Definitions

## (Joint Distribution).

The **joint distribution** for two discrete random variables  $X$  and  $Y$  is the collection of values  $\{((x, y), P(X = x, Y = y)) : x \in \mathcal{X}, y \in \mathcal{Y}\}$ , where  $\mathcal{X}$  is the set of all possible values taken by  $X$  and  $\mathcal{Y}$  is the set of all possible values taken by  $Y$ .

## (Marginal Distribution).

Given the joint distribution for  $X$  and  $Y$ , the **marginal distribution** for  $X$  is as follows:

# Definitions

## (Joint Distribution).

The **joint distribution** for two discrete random variables  $X$  and  $Y$  is the collection of values  $\{((x, y), P(X = x, Y = y)) : x \in \mathcal{X}, y \in \mathcal{Y}\}$ , where  $\mathcal{X}$  is the set of all possible values taken by  $X$  and  $\mathcal{Y}$  is the set of all possible values taken by  $Y$ .

## (Marginal Distribution).

Given the joint distribution for  $X$  and  $Y$ , the **marginal distribution** for  $X$  is as follows:

$$P(X = x) = \sum_{y \in \mathcal{Y}} P(X = x, Y = y).$$

# Independence



# Independence

(Independence).

# Independence

## (Independence).

Random variables  $X$  and  $Y$  are said to be **independent** if the events  $X = x$  and  $Y = y$  are independent for all values  $x, y$ . Equivalently, the joint distribution of independent R.V's decomposes as

# Independence

## (Independence).

Random variables  $X$  and  $Y$  are said to be **independent** if the events  $X = x$  and  $Y = y$  are independent for all values  $x, y$ . Equivalently, the joint distribution of independent R.V's decomposes as

$$P(X = x, Y = y) = P(X = x)P(Y = y), \quad \forall x, y.$$

# Expectation

# Expectation

(Expectation).

# Expectation

(Expectation).

The **expectation** of a discrete random variable  $X$  is defined as

# Expectation

## (Expectation).

The **expectation** of a discrete random variable  $X$  is defined as

$$\mathbb{E}[X] = \sum_{x \in \mathcal{X}} x \cdot P(X = x).$$

# Expectation

## (Expectation).

The **expectation** of a discrete random variable  $X$  is defined as

$$\mathbb{E}[X] = \sum_{x \in \mathcal{X}} x \cdot P(X = x).$$

Alternatively, we also have



# Expectation

## (Expectation).

The **expectation** of a discrete random variable  $X$  is defined as

$$\mathbb{E}[X] = \sum_{x \in \mathcal{X}} x \cdot P(X = x).$$

Alternatively, we also have

$$\mathbb{E}[X] = \sum_{\omega \in \Omega} X(\omega) \cdot P(\omega).$$

# Variance

# Variance

(Variance).

# Variance

(Variance).

The **variance** of a random variable  $X$  is

# Variance

**(Variance).**

The **variance** of a random variable  $X$  is

$$\text{Var}(X) = \mathbb{E}[(X - \mathbb{E}[X])^2].$$

# Variance

## (Variance).

The **variance** of a random variable  $X$  is

$$\text{Var}(X) = \mathbb{E}[(X - \mathbb{E}[X])^2].$$

## (Standard Deviation).

# Variance

## (Variance).

The **variance** of a random variable  $X$  is

$$\text{Var}(X) = \mathbb{E}[(X - \mathbb{E}[X])^2].$$

## (Standard Deviation).

The **standard deviation** of a random variable  $X$

# Variance

## (Variance).

The **variance** of a random variable  $X$  is

$$\text{Var}(X) = \mathbb{E}[(X - \mathbb{E}[X])^2].$$

## (Standard Deviation).

The **standard deviation** of a random variable  $X$

$$\sigma := \sqrt{\text{Var}(X)}.$$



# Variance

# Variance

## Theorem

# Variance

## Theorem

For a random variable  $X$ ,

# Variance

## Theorem

For a random variable  $X$ ,

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2.$$

# Variance

## Theorem

For a random variable  $X$ ,

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2.$$

***Proof:***

# Variance

## Theorem

For a random variable  $X$ ,

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2.$$

***Proof:***

$$\text{Var}(X) = \mathbb{E} \left[ (X - \mathbb{E}[X])^2 \right]$$

# Variance

## Theorem

For a random variable  $X$ ,

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2.$$

## *Proof:*

$$\begin{aligned}\text{Var}(X) &= \mathbb{E} \left[ (X - \mathbb{E}[X])^2 \right] \\ &= \mathbb{E} \left[ X^2 - 2X\mathbb{E}[X] + \mathbb{E}[X]^2 \right]\end{aligned}$$

# Variance

## Theorem

For a random variable  $X$ ,

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2.$$

## *Proof:*

$$\begin{aligned}\text{Var}(X) &= \mathbb{E} \left[ (X - \mathbb{E}[X])^2 \right] \\ &= \mathbb{E} \left[ X^2 - 2X\mathbb{E}[X] + \mathbb{E}[X]^2 \right] \\ &= \mathbb{E} \left[ X^2 \right] - \mathbb{E}[2X\mathbb{E}[X]] + \mathbb{E} \left[ \mathbb{E}[X]^2 \right]\end{aligned}$$



# Variance

## Theorem

For a random variable  $X$ ,

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2.$$

## *Proof:*

$$\begin{aligned}\text{Var}(X) &= \mathbb{E} \left[ (X - \mathbb{E}[X])^2 \right] \\ &= \mathbb{E} \left[ X^2 - 2X\mathbb{E}[X] + \mathbb{E}[X]^2 \right] \\ &= \mathbb{E} \left[ X^2 \right] - \mathbb{E}[2X\mathbb{E}[X]] + \mathbb{E} \left[ \mathbb{E}[X]^2 \right] \\ &= \mathbb{E} \left[ X^2 \right] - 2\mathbb{E}[X]^2 + \mathbb{E}[X]^2\end{aligned}$$

# Variance

## Theorem

For a random variable  $X$ ,

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2.$$

## *Proof:*

$$\begin{aligned}\text{Var}(X) &= \mathbb{E} \left[ (X - \mathbb{E}[X])^2 \right] \\ &= \mathbb{E} \left[ X^2 - 2X\mathbb{E}[X] + \mathbb{E}[X]^2 \right] \\ &= \mathbb{E} \left[ X^2 \right] - \mathbb{E}[2X\mathbb{E}[X]] + \mathbb{E} \left[ \mathbb{E}[X]^2 \right] \\ &= \mathbb{E} \left[ X^2 \right] - 2\mathbb{E}[X]^2 + \mathbb{E}[X]^2 \\ &= \mathbb{E} \left[ X^2 \right] - \mathbb{E}[X]^2.\end{aligned}$$

# Variance

## Theorem

## Theorem

For independent random variables  $X, Y$ ,  $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$ .

## Theorem

For independent random variables  $X, Y$ ,  $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$ .

***Proof:***

## Theorem

For independent random variables  $X, Y$ ,  $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$ .

**Proof:**

$$\mathbb{E}[XY] = \sum_x \sum_y xy \cdot P(X = x, Y = y)$$

## Theorem

For independent random variables  $X, Y$ ,  $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$ .

**Proof:**

$$\begin{aligned}\mathbb{E}[XY] &= \sum_x \sum_y xy \cdot P(X = x, Y = y) \\ &= \sum_x \sum_y xy \cdot P(X = x) \cdot P(Y = y)\end{aligned}$$



## Theorem

For independent random variables  $X, Y$ ,  $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$ .

**Proof:**

$$\begin{aligned}\mathbb{E}[XY] &= \sum_x \sum_y xy \cdot P(X = x, Y = y) \\ &= \sum_x \sum_y xy \cdot P(X = x) \cdot P(Y = y) \\ &= \left( \sum_x x \cdot P(X = x) \right) \cdot \left( \sum_y y \cdot P(Y = y) \right)\end{aligned}$$

## Theorem

For independent random variables  $X, Y$ ,  $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$ .

**Proof:**

$$\begin{aligned}\mathbb{E}[XY] &= \sum_x \sum_y xy \cdot P(X = x, Y = y) \\ &= \sum_x \sum_y xy \cdot P(X = x) \cdot P(Y = y) \\ &= \left( \sum_x x \cdot P(X = x) \right) \cdot \left( \sum_y y \cdot P(Y = y) \right) \\ &= \mathbb{E}[X] \cdot \mathbb{E}[Y]\end{aligned}$$

## Theorem

For independent random variables  $X, Y$ ,  $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$ .

**Proof:**

$$\begin{aligned}\mathbb{E}[XY] &= \sum_x \sum_y xy \cdot P(X = x, Y = y) \\ &= \sum_x \sum_y xy \cdot P(X = x) \cdot P(Y = y) \\ &= \left( \sum_x x \cdot P(X = x) \right) \cdot \left( \sum_y y \cdot P(Y = y) \right) \\ &= \mathbb{E}[X] \cdot \mathbb{E}[Y]\end{aligned}$$

where the second line follows from independence. □

# Variance

## Theorem

## Theorem

For **independent** random variables  $X, Y$ ,

## Theorem

For **independent** random variables  $X, Y$ ,

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

## Theorem

For **independent** random variables  $X, Y$ ,

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

***Proof:***



## Theorem

For **independent** random variables  $X, Y$ ,

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

## ***Proof:***

Try it on your own.

## Theorem

For **independent** random variables  $X, Y$ ,

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y).$$

## *Proof:*

Try it on your own. □

# Bernoulli Distribution

# Bernoulli Distribution

- $X \sim \text{Bernoulli}(p)$ .

# Bernoulli Distribution

- $X \sim \text{Bernoulli}(p)$ .
- One coin flip with probability of  $p$  getting a head.

# Bernoulli Distribution

- $X \sim \text{Bernoulli}(p)$ .
- One coin flip with probability of  $p$  getting a head.
- **PMF:**

# Bernoulli Distribution

- $X \sim \text{Bernoulli}(p)$ .
- One coin flip with probability of  $p$  getting a head.
- **PMF:**

$$P(X = i) = \begin{cases} p, & \text{if } i = 1 \\ 1 - p, & \text{if } i = 0 \end{cases}$$

# Bernoulli Distribution

- $X \sim \text{Bernoulli}(p)$ .
- One coin flip with probability of  $p$  getting a head.
- **PMF:**

$$P(X = i) = \begin{cases} p, & \text{if } i = 1 \\ 1 - p, & \text{if } i = 0 \end{cases}$$

where  $0 \leq p \leq 1$ .



# Bernoulli Distribution

- $X \sim \text{Bernoulli}(p)$ .
- One coin flip with probability of  $p$  getting a head.

- **PMF:**

$$P(X = i) = \begin{cases} p, & \text{if } i = 1 \\ 1 - p, & \text{if } i = 0 \end{cases}$$

where  $0 \leq p \leq 1$ .

- **Expectation:**

# Bernoulli Distribution

- $X \sim \text{Bernoulli}(p)$ .
- One coin flip with probability of  $p$  getting a head.

- **PMF:**

$$P(X = i) = \begin{cases} p, & \text{if } i = 1 \\ 1 - p, & \text{if } i = 0 \end{cases}$$

where  $0 \leq p \leq 1$ .

- **Expectation:**

$$\mathbb{E}[X] = p.$$

# Bernoulli Distribution

- $X \sim \text{Bernoulli}(p)$ .
- One coin flip with probability of  $p$  getting a head.

- **PMF:**

$$P(X = i) = \begin{cases} p, & \text{if } i = 1 \\ 1 - p, & \text{if } i = 0 \end{cases}$$

where  $0 \leq p \leq 1$ .

- **Expectation:**

$$\mathbb{E}[X] = p.$$

- **Variance:**

# Bernoulli Distribution

- $X \sim \text{Bernoulli}(p)$ .
- One coin flip with probability of  $p$  getting a head.

- **PMF:**

$$P(X = i) = \begin{cases} p, & \text{if } i = 1 \\ 1 - p, & \text{if } i = 0 \end{cases}$$

where  $0 \leq p \leq 1$ .

- **Expectation:**

$$\mathbb{E}[X] = p.$$

- **Variance:**

$$\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = p - p^2 = p(1 - p).$$

# Binomial Distribution

# Binomial Distribution

- $X \sim \text{Bin}(n, p)$ .

# Binomial Distribution

- $X \sim \text{Bin}(n, p)$ .
- $n$  coin flips with probability of  $p$  getting a head.

# Binomial Distribution

- $X \sim \text{Bin}(n, p)$ .
- $n$  coin flips with probability of  $p$  getting a head.
- **PMF:**



# Binomial Distribution

- $X \sim \text{Bin}(n, p)$ .
- $n$  coin flips with probability of  $p$  getting a head.
- **PMF:**

$$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}, \quad \text{for } k = 0, 1, \dots, n.$$

# Binomial Distribution

- $X \sim \text{Bin}(n, p)$ .
- $n$  coin flips with probability of  $p$  getting a head.

- **PMF:**

$$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}, \quad \text{for } k = 0, 1, \dots, n.$$

- **Expectation:**

# Binomial Distribution

- $X \sim \text{Bin}(n, p)$ .
- $n$  coin flips with probability of  $p$  getting a head.

- **PMF:**

$$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}, \quad \text{for } k = 0, 1, \dots, n.$$

- **Expectation:**

$$\mathbb{E}[X] = \mathbb{E}\left[\sum_{i=1}^n Y_i\right] = \sum_{i=1}^n \mathbb{E}[Y_i] = \sum_{i=1}^n p = np, \quad Y_i \sim \text{Bernoulli}(p).$$

# Binomial Distribution

- $X \sim \text{Bin}(n, p)$ .
- $n$  coin flips with probability of  $p$  getting a head.
- **PMF:**

$$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}, \quad \text{for } k = 0, 1, \dots, n.$$

- **Expectation:**

$$\mathbb{E}[X] = \mathbb{E}\left[\sum_{i=1}^n Y_i\right] = \sum_{i=1}^n \mathbb{E}[Y_i] = \sum_{i=1}^n p = np, \quad Y_i \sim \text{Bernoulli}(p).$$

- **Variance:**

# Binomial Distribution

- $X \sim \text{Bin}(n, p)$ .
- $n$  coin flips with probability of  $p$  getting a head.

- **PMF:**

$$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}, \quad \text{for } k = 0, 1, \dots, n.$$

- **Expectation:**

$$\mathbb{E}[X] = \mathbb{E}\left[\sum_{i=1}^n Y_i\right] = \sum_{i=1}^n \mathbb{E}[Y_i] = \sum_{i=1}^n p = np, \quad Y_i \sim \text{Bernoulli}(p).$$

- **Variance:**

$$\text{Var}(X) = \text{Var}\left(\sum_{i=1}^n Y_i\right) = np(1 - p).$$

# Indicator Random Variable

# Indicator Random Variable

(Indicator Random variable).

# Indicator Random Variable

**(Indicator Random variable).**

We use  $\mathbb{I}_i$ , or  $X_i$  to denotes the **indicator random variable** that takes on values  $\{0, 1\}$  according to whether a specified event occurs or not.



# Indicator Random Variable

## (Indicator Random variable).

We use  $\mathbb{I}_i$ , or  $X_i$  to denotes the **indicator random variable** that takes on values  $\{0, 1\}$  according to whether a specified event occurs or not.

- Usually  $\{\mathbb{I}_i\}_{i=1}^n$  are mutually independent and they are said to be *independent and identically distributed (i.i.d)*.

# Indicator Random Variable

## (Indicator Random variable).

We use  $\mathbb{I}_i$ , or  $X_i$  to denotes the **indicator random variable** that takes on values  $\{0, 1\}$  according to whether a specified event occurs or not.

- Usually  $\{\mathbb{I}_i\}_{i=1}^n$  are mutually independent and they are said to be ***independent and identically distributed (i.i.d)***.
- Will be very useful soon for computing expectations.

# Geometric Distribution

# Geometric Distribution

- $X \sim \text{Geo}(p)$ .

# Geometric Distribution

- $X \sim \text{Geo}(p)$ .
- Number of coin flips until first head appears.

# Geometric Distribution

- $X \sim \text{Geo}(p)$ .
- Number of coin flips until first head appears.
- **PMF:**

# Geometric Distribution

- $X \sim \text{Geo}(p)$ .
- Number of coin flips until first head appears.
- **PMF:**

$$P(X = k) = (1 - p)^{k-1}p, \quad \text{for } k = 1, 2, 3, \dots$$

# Geometric Distribution

- $X \sim \text{Geo}(p)$ .
- Number of coin flips until first head appears.

- **PMF:**

$$P(X = k) = (1 - p)^{k-1}p, \quad \text{for } k = 1, 2, 3, \dots$$

- **Expectation:**



# Geometric Distribution

- $X \sim \text{Geo}(p)$ .
- Number of coin flips until first head appears.
- **PMF:**

$$P(X = k) = (1 - p)^{k-1}p, \quad \text{for } i = 1, 2, 3, \dots$$

- **Expectation:**

$$\mathbb{E}[X] = \sum_{i=1}^{\infty} P(X \geq i) = \sum_{x=1}^{\infty} (1 - p)^{i-1} = \frac{1}{1 - (1 - p)} = \frac{1}{p},$$

# Geometric Distribution

- $X \sim \text{Geo}(p)$ .
- Number of coin flips until first head appears.
- **PMF:**

$$P(X = k) = (1 - p)^{k-1}p, \quad \text{for } i = 1, 2, 3, \dots$$

- **Expectation:**

$$\mathbb{E}[X] = \sum_{i=1}^{\infty} P(X \geq i) = \sum_{x=1}^{\infty} (1 - p)^{i-1} = \frac{1}{1 - (1 - p)} = \frac{1}{p},$$

where the first equality uses the **tail sum formula** (discussed in next slide).

# Geometric Distribution

- $X \sim \text{Geo}(p)$ .
- Number of coin flips until first head appears.
- **PMF:**

$$P(X = k) = (1 - p)^{k-1}p, \quad \text{for } i = 1, 2, 3, \dots$$

- **Expectation:**

$$\mathbb{E}[X] = \sum_{i=1}^{\infty} P(X \geq i) = \sum_{x=1}^{\infty} (1 - p)^{i-1} = \frac{1}{1 - (1 - p)} = \frac{1}{p},$$

where the first equality uses the **tail sum formula** (discussed in next slide).

- **Variance:**

# Geometric Distribution

- $X \sim \text{Geo}(p)$ .
- Number of coin flips until first head appears.
- **PMF:**

$$P(X = k) = (1 - p)^{k-1}p, \quad \text{for } i = 1, 2, 3, \dots$$

- **Expectation:**

$$\mathbb{E}[X] = \sum_{i=1}^{\infty} P(X \geq i) = \sum_{x=1}^{\infty} (1 - p)^{i-1} = \frac{1}{1 - (1 - p)} = \frac{1}{p},$$

where the first equality uses the **tail sum formula** (discussed in next slide).

- **Variance:**

$$\text{Var}(X) = \frac{1 - p}{p^2}.$$

# Tail Sum Formula

# Tail Sum Formula

## Tail Sum Formula.

# Tail Sum Formula

## Tail Sum Formula.

Let  $X$  be a random variable that takes values in  $\{0, 1, 2, \dots\}$ . Then

# Tail Sum Formula

## Tail Sum Formula.

Let  $X$  be a random variable that takes values in  $\{0, 1, 2, \dots\}$ . Then

$$\mathbb{E}[X] = \sum_{i=1}^{\infty} P(X \geq i).$$



# Tail Sum Formula

## Tail Sum Formula.

Let  $X$  be a random variable that takes values in  $\{0, 1, 2, \dots\}$ . Then

$$\mathbb{E}[X] = \sum_{i=1}^{\infty} P(X \geq i).$$

***Proof:***

# Tail Sum Formula

## Tail Sum Formula.

Let  $X$  be a random variable that takes values in  $\{0, 1, 2, \dots\}$ . Then

$$\mathbb{E}[X] = \sum_{i=1}^{\infty} P(X \geq i).$$

**Proof:**

$$\mathbb{E}[X] = \sum_{x=1}^{\infty} xP(X = x)$$

# Tail Sum Formula

## Tail Sum Formula.

Let  $X$  be a random variable that takes values in  $\{0, 1, 2, \dots\}$ . Then

$$\mathbb{E}[X] = \sum_{i=1}^{\infty} P(X \geq i).$$

**Proof:**

$$\begin{aligned}\mathbb{E}[X] &= \sum_{x=1}^{\infty} xP(X = x) \\ &= \sum_{x=1}^{\infty} \sum_{i=1}^x P(X = x)\end{aligned}$$

# Tail Sum Formula

## Tail Sum Formula.

Let  $X$  be a random variable that takes values in  $\{0, 1, 2, \dots\}$ . Then

$$\mathbb{E}[X] = \sum_{i=1}^{\infty} P(X \geq i).$$

**Proof:**

$$\begin{aligned}\mathbb{E}[X] &= \sum_{x=1}^{\infty} xP(X = x) \\ &= \sum_{x=1}^{\infty} \sum_{i=1}^x P(X = x) \\ &= \sum_{i=1}^{\infty} \sum_{x=i}^{\infty} P(X = x)\end{aligned}$$

# Tail Sum Formula

## Tail Sum Formula.

Let  $X$  be a random variable that takes values in  $\{0, 1, 2, \dots\}$ . Then

$$\mathbb{E}[X] = \sum_{i=1}^{\infty} P(X \geq i).$$

**Proof:**

$$\begin{aligned}\mathbb{E}[X] &= \sum_{x=1}^{\infty} xP(X = x) \\ &= \sum_{x=1}^{\infty} \sum_{i=1}^x P(X = x) \\ &= \sum_{i=1}^{\infty} \sum_{x=i}^{\infty} P(X = x) \\ &= \sum_{i=1}^{\infty} P(X \geq i).\end{aligned}$$

# A Clever Way of Computing the Geometric Expectation

# A Clever Way of Computing the Geometric Expectation

The **Tail Sum Formula** certainly makes the computation a lot easier, but there's an even more simpler way.

# A Clever Way of Computing the Geometric Expectation

The **Tail Sum Formula** certainly makes the computation a lot easier, but there's an even more simpler way.

- Suppose we toss our first coin. There are two possibilities: (1) we get a head with probability  $p$  and call it a day, (2) we get a tail with probability  $1 - p$  and we are right back where we just started.



# A Clever Way of Computing the Geometric Expectation

The **Tail Sum Formula** certainly makes the computation a lot easier, but there's an even more simpler way.

- Suppose we toss our first coin. There are two possibilities: (1) we get a head with probability  $p$  and call it a day, (2) we get a tail with probability  $1 - p$  and we are right back where we just started.
- In the latter case, we expect  $1 + \mathbb{E}[X]$  trials until our first success because we already wasted one trial. Hence,

# A Clever Way of Computing the Geometric Expectation

The **Tail Sum Formula** certainly makes the computation a lot easier, but there's an even more simpler way.

- Suppose we toss our first coin. There are two possibilities: (1) we get a head with probability  $p$  and call it a day, (2) we get a tail with probability  $1 - p$  and we are right back where we just started.
- In the latter case, we expect  $1 + \mathbb{E}[X]$  trials until our first success because we already wasted one trial. Hence,

$$\mathbb{E}[X] = p \cdot 1 + (1 - p)(1 + \mathbb{E}[X]) \implies \mathbb{E}[X] = \frac{1}{p}.$$

# A Clever Way of Computing the Geometric Expectation

The **Tail Sum Formula** certainly makes the computation a lot easier, but there's an even more simpler way.

- Suppose we toss our first coin. There are two possibilities: (1) we get a head with probability  $p$  and call it a day, (2) we get a tail with probability  $1 - p$  and we are right back where we just started.
- In the latter case, we expect  $1 + \mathbb{E}[X]$  trials until our first success because we already wasted one trial. Hence,

$$\mathbb{E}[X] = p \cdot 1 + (1 - p)(1 + \mathbb{E}[X]) \implies \mathbb{E}[X] = \frac{1}{p}.$$

This makes use of an important property called the **memoryless property**, which will be covered later in the class.

# Poisson Distribution

# Poisson Distribution

- $X \sim \text{Poisson}(\lambda)$ .

# Poisson Distribution

- $X \sim \text{Poisson}(\lambda)$ .
- Models rare events, such as number of arrivals of a bus in an hour.

# Poisson Distribution

- $X \sim \text{Poisson}(\lambda)$ .
- Models rare events, such as number of arrivals of a bus in an hour.
- Defined in terms of a rate  $\lambda$ , which specifies the average number of times an event occurs in a time interval.

# Poisson Distribution

- $X \sim \text{Poisson}(\lambda)$ .
- Models rare events, such as number of arrivals of a bus in an hour.
- Defined in terms of a rate  $\lambda$ , which specifies the average number of times an event occurs in a time interval.
- **PMF:**



# Poisson Distribution

- $X \sim \text{Poisson}(\lambda)$ .
- Models rare events, such as number of arrivals of a bus in an hour.
- Defined in terms of a rate  $\lambda$ , which specifies the average number of times an event occurs in a time interval.
- **PMF:**

$$P(X = k) = \frac{\lambda^k e^{-\lambda}}{k!}, \quad \text{for } i = 0, 1, 2, \dots$$

# Poisson Distribution

- $X \sim \text{Poisson}(\lambda)$ .
- Models rare events, such as number of arrivals of a bus in an hour.
- Defined in terms of a rate  $\lambda$ , which specifies the average number of times an event occurs in a time interval.

- **PMF:**

$$P(X = k) = \frac{\lambda^k e^{-\lambda}}{k!}, \quad \text{for } i = 0, 1, 2, \dots$$



$$\sum_{i=0}^{\infty} P(X = i) = \sum_{i=0}^{\infty} \frac{\lambda^i e^{-\lambda}}{i!} = e^{-\lambda} \sum_{i=0}^{\infty} \frac{\lambda^i}{i!} = e^{-\lambda} \cdot e^{\lambda} = 1.$$

# Poisson Distribution

- $X \sim \text{Poisson}(\lambda)$ .
- Models rare events, such as number of arrivals of a bus in an hour.
- Defined in terms of a rate  $\lambda$ , which specifies the average number of times an event occurs in a time interval.

- **PMF:**

$$P(X = k) = \frac{\lambda^k e^{-\lambda}}{k!}, \quad \text{for } k = 0, 1, 2, \dots$$



$$\sum_{i=0}^{\infty} P(X = i) = \sum_{i=0}^{\infty} \frac{\lambda^i e^{-\lambda}}{i!} = e^{-\lambda} \sum_{i=0}^{\infty} \frac{\lambda^i}{i!} = e^{-\lambda} \cdot e^{\lambda} = 1.$$

Recall the **Taylor series expansion** from calculus:

# Poisson Distribution

- $X \sim \text{Poisson}(\lambda)$ .
- Models rare events, such as number of arrivals of a bus in an hour.
- Defined in terms of a rate  $\lambda$ , which specifies the average number of times an event occurs in a time interval.

- **PMF:**

$$P(X = k) = \frac{\lambda^k e^{-\lambda}}{k!}, \quad \text{for } i = 0, 1, 2, \dots$$



$$\sum_{i=0}^{\infty} P(X = i) = \sum_{i=0}^{\infty} \frac{\lambda^i e^{-\lambda}}{i!} = e^{-\lambda} \sum_{i=0}^{\infty} \frac{\lambda^i}{i!} = e^{-\lambda} \cdot e^{\lambda} = 1.$$

Recall the **Taylor series expansion** from calculus:

$$e^x = \sum_{i=0}^{\infty} \frac{x^i}{i!}.$$

# Poisson Distribution

# Poisson Distribution

- **Expectation:**

# Poisson Distribution

- **Expectation:**

$$\mathbb{E}[X] = \sum_{i=0}^{\infty} i \cdot P(X = i)$$

# Poisson Distribution

- **Expectation:**

$$\begin{aligned}\mathbb{E}[X] &= \sum_{i=0}^{\infty} i \cdot P(X = i) \\ &= \sum_{i=1}^{\infty} \frac{\lambda^i e^{-\lambda}}{i!}\end{aligned}$$



# Poisson Distribution

- **Expectation:**

$$\begin{aligned}\mathbb{E}[X] &= \sum_{i=0}^{\infty} i \cdot P(X = i) \\ &= \sum_{i=1}^{\infty} \frac{\lambda^i e^{-\lambda}}{i!} \\ &= \lambda e^{-\lambda} \sum_{i=1}^{\infty} \frac{\lambda^{i-1}}{(i-1)!}\end{aligned}$$

# Poisson Distribution

- **Expectation:**

$$\begin{aligned}\mathbb{E}[X] &= \sum_{i=0}^{\infty} i \cdot P(X = i) \\ &= \sum_{i=1}^{\infty} \frac{\lambda^i e^{-\lambda}}{i!} \\ &= \lambda e^{-\lambda} \sum_{i=1}^{\infty} \frac{\lambda^{i-1}}{(i-1)!} \\ &= \lambda e^{-\lambda} e^{\lambda}\end{aligned}$$

$$(e^{\lambda} = \sum_{j=1}^{\infty} \frac{\lambda^j}{j!} \text{ with } j = i - 1)$$

# Poisson Distribution

- **Expectation:**

$$\begin{aligned}\mathbb{E}[X] &= \sum_{i=0}^{\infty} i \cdot P(X = i) \\ &= \sum_{i=1}^{\infty} \frac{\lambda^i e^{-\lambda}}{i!} \\ &= \lambda e^{-\lambda} \sum_{i=1}^{\infty} \frac{\lambda^{i-1}}{(i-1)!} \\ &= \lambda e^{-\lambda} e^{\lambda} \\ &= \lambda.\end{aligned}$$

$$(e^{\lambda} = \sum_{j=1}^{\infty} \frac{\lambda^j}{j!} \text{ with } j = i - 1)$$

# Poisson Distribution

- **Expectation:**

$$\begin{aligned}\mathbb{E}[X] &= \sum_{i=0}^{\infty} i \cdot P(X = i) \\ &= \sum_{i=1}^{\infty} \frac{\lambda^i e^{-\lambda}}{i!} \\ &= \lambda e^{-\lambda} \sum_{i=1}^{\infty} \frac{\lambda^{i-1}}{(i-1)!} \\ &= \lambda e^{-\lambda} e^{\lambda} \\ &= \lambda.\end{aligned}$$

$$(e^{\lambda} = \sum_{j=1}^{\infty} \frac{\lambda^j}{j!} \text{ with } j = i - 1)$$

- **Variance:**

# Poisson Distribution

- **Expectation:**

$$\begin{aligned}\mathbb{E}[X] &= \sum_{i=0}^{\infty} i \cdot P(X = i) \\ &= \sum_{i=1}^{\infty} \frac{\lambda^i e^{-\lambda}}{i!} \\ &= \lambda e^{-\lambda} \sum_{i=1}^{\infty} \frac{\lambda^{i-1}}{(i-1)!} \\ &= \lambda e^{-\lambda} e^{\lambda} \quad \left( e^{\lambda} = \sum_{j=1}^{\infty} \frac{\lambda^j}{j!} \text{ with } j = i - 1 \right) \\ &= \lambda.\end{aligned}$$

- **Variance:**

$$\text{Var}(X) = \lambda.$$

# Problem Time!