

CS70 Modular Arithmetic

Kelvin Lee

kelvinlee@berkeley.edu

September 28, 2020

Overview

- 1 Basic Definitions
- 2 Multiplicative Inverse
- 3 Euclid's Algorithm
- 4 Extended Euclid's algorithm
- 5 Functions
- 6 Bijection
- 7 Fermat's Little Theorem
- 8 Chinese Remainder Theorem

Basic Definitions

Definition (Congruence)

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m
- x and y have the same remainder w.r.t. m

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m
- x and y have the same remainder w.r.t. m
- $x = y + km$ for some integer k

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m
 - x and y have the same remainder w.r.t. m
 - $x = y + km$ for some integer k
-
- In modulo m , only the numbers $\{0, 1, 2, \dots, m - 1\}$ exist.

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m
 - x and y have the same remainder w.r.t. m
 - $x = y + km$ for some integer k
-
- In modulo m , only the numbers $\{0, 1, 2, \dots, m - 1\}$ exist.
 - Division is not well-defined.

Basic Definitions

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m
- x and y have the same remainder w.r.t. m
- $x = y + km$ for some integer k
- In modulo m , only the numbers $\{0, 1, 2, \dots, m - 1\}$ exist.
- Division is not well-defined.

Definition (Multiplicative Inverse)

Basic Definitions

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y \pmod{m}$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m
 - x and y have the same remainder w.r.t. m
 - $x = y + km$ for some integer k
-
- In modulo m , only the numbers $\{0, 1, 2, \dots, m - 1\}$ exist.
 - Division is not well-defined.

Definition (Multiplicative Inverse)

Normally we say that the **multiplicative inverse** of x is y if $xy = 1$.

Basic Definitions

Definition (Congruence)

x is **congruent** to y modulo m or $x \equiv y(\text{mod } m)$ if and only if any one of the following is true:

- $(x - y)$ is divisible by m
 - x and y have the same remainder w.r.t. m
 - $x = y + km$ for some integer k
-
- In modulo m , only the numbers $\{0, 1, 2, \dots, m - 1\}$ exist.
 - Division is not well-defined.

Definition (Multiplicative Inverse)

Normally we say that the **multiplicative inverse** of x is y if $xy = 1$. In the modular space, the **multiplicative inverse** of $x \text{ mod } m$ is y if $xy \equiv 1(\text{mod } m)$.

Theorems

Theorem (Modular operations)

Theorem (Modular operations)

$a \equiv c \pmod{m}$ and $b \equiv d \pmod{m} \implies a + b \equiv c + d \pmod{m}$ and $a \cdot b \equiv c \cdot d \pmod{m}$.

Theorems

Theorem (Modular operations)

$a \equiv c \pmod{m}$ and $b \equiv d \pmod{m} \implies a + b \equiv c + d \pmod{m}$ and $a \cdot b \equiv c \cdot d \pmod{m}$.

Theorem (Existence of multiplicative inverse)

Theorems

Theorem (Modular operations)

$a \equiv c \pmod{m}$ and $b \equiv d \pmod{m} \implies a + b \equiv c + d \pmod{m}$ and $a \cdot b \equiv c \cdot d \pmod{m}$.

Theorem (Existence of multiplicative inverse)

$\gcd(x, m) = 1 \implies x$ has a multiplicative inverse modulo m and it is **unique**.

Euclid's Algorithm

Euclid's Algorithm

How do we compute \gcd of two numbers x and y ?

Euclid's Algorithm

How do we compute \gcd of two numbers x and y ?

Theorem (Euclid's Algorithm)

Euclid's Algorithm

How do we compute gcd of two numbers x and y ?

Theorem (Euclid's Algorithm)

Let $x \geq y > 0$. Then

$$\gcd(x, y) = \gcd(y, x \bmod y)$$

Euclid's Algorithm

How do we compute gcd of two numbers x and y ?

Theorem (Euclid's Algorithm)

Let $x \geq y > 0$. Then

$$\gcd(x, y) = \gcd(y, x \bmod y)$$

Example

Euclid's Algorithm

How do we compute gcd of two numbers x and y ?

Theorem (Euclid's Algorithm)

Let $x \geq y > 0$. Then

$$\gcd(x, y) = \gcd(y, x \bmod y)$$

Example

Compute $\gcd(16, 10)$:

Euclid's Algorithm

How do we compute gcd of two numbers x and y ?

Theorem (Euclid's Algorithm)

Let $x \geq y > 0$. Then

$$\gcd(x, y) = \gcd(y, x \bmod y)$$

Example

Compute $\gcd(16, 10)$:

$$\begin{aligned}\gcd(16, 10) &= \gcd(10, 6) \\ &= \gcd(6, 4) \\ &= \gcd(4, 2) \\ &= \gcd(2, 0) \\ &= 2.\end{aligned}$$

Extended Euclid's algorithm

Extended Euclid's algorithm

How to compute the multiplicative inverse?

Extended Euclid's algorithm

How to compute the multiplicative inverse?

- Need an algorithm that returns integers a and b such that:

$$\gcd(x, y) = ax + by.$$

Extended Euclid's algorithm

How to compute the multiplicative inverse?

- Need an algorithm that returns integers a and b such that:

$$\gcd(x, y) = ax + by.$$

Theorem (Bézout's Identity)

Extended Euclid's algorithm

How to compute the multiplicative inverse?

- Need an algorithm that returns integers a and b such that:

$$\gcd(x, y) = ax + by.$$

Theorem (Bézout's Identity)

For nonzero integers x and y , let d be the greatest common divisor such that $d = \gcd(x, y)$. Then, there exist integers a and b such that

$$ax + by = d.$$

Extended Euclid's algorithm

How to compute the multiplicative inverse?

- Need an algorithm that returns integers a and b such that:

$$\gcd(x, y) = ax + by.$$

Theorem (Bézout's Identity)

For nonzero integers x and y , let d be the greatest common divisor such that $d = \gcd(x, y)$. Then, there exist integers a and b such that

$$ax + by = d.$$

- When $\gcd(x, y) = 1$, we can deduce that b is an inverse of $y \bmod x$.

Extended Euclid's algorithm

How to compute the multiplicative inverse?

- Need an algorithm that returns integers a and b such that:

$$\gcd(x, y) = ax + by.$$

Theorem (Bézout's Identity)

For nonzero integers x and y , let d be the greatest common divisor such that $d = \gcd(x, y)$. Then, there exist integers a and b such that

$$ax + by = d.$$

- When $\gcd(x, y) = 1$, we can deduce that b is an inverse of $y \bmod x$.
- This uses back substitutions repetitively so that the final expression is in terms of x and y .

Functions

Definition (Function)

Definition (Function)

Let A and B be nonempty sets. A **function** f from A to B is an assignment of exactly one element of B to each element of A . (vertical line test)

Definition (Function)

Let A and B be nonempty sets. A **function** f from A to B is an assignment of exactly one element of B to each element of A . (vertical line test)

- To denote such a function, we write $f : A \rightarrow B$ (f maps A to B).

Definition (Function)

Let A and B be nonempty sets. A **function** f from A to B is an assignment of exactly one element of B to each element of A . (vertical line test)

- To denote such a function, we write $f : A \rightarrow B$ (f maps A to B).
- A is the **domain** and B is the **co-domain**.

Definition (Function)

Let A and B be nonempty sets. A **function** f from A to B is an assignment of exactly one element of B to each element of A . (vertical line test)

- To denote such a function, we write $f : A \rightarrow B$ (f maps A to B).
- A is the **domain** and B is the **co-domain**.
- Pre-image is a **subset** of domain, and the image/range is the **subset** of co-domain.

Definition (Function)

Let A and B be nonempty sets. A **function** f from A to B is an assignment of exactly one element of B to each element of A . (vertical line test)

- To denote such a function, we write $f : A \rightarrow B$ (f maps A to B).
- A is the **domain** and B is the **co-domain**.
- Pre-image is a **subset** of domain, and the image/range is the **subset** of co-domain.
 - If $f(a) = b$, where $a \in A$ and $b \in B$, then we say that b is the image of a and a is the pre-image of b .

Bijection

Definition (One-to-one)

Definition (One-to-one)

A function f is said to be **one-to-one** if and only if $f(a) = f(a')$ implies that $a = a'$ for all $a, a' \in A$. A function is said to be **injective** if it is **one-to-one**.

Definition (One-to-one)

A function f is said to be **one-to-one** if and only if $f(a) = f(a')$ implies that $a = a'$ for all $a, a' \in A$. A function is said to be **injective** if it is **one-to-one**.

- To show that a function is *one-to-one*, we show that $a \neq a' \implies f(a) \neq f(a')$. (Why?)

Bijection

Definition (One-to-one)

A function f is said to be **one-to-one** if and only if $f(a) = f(a')$ implies that $a = a'$ for all $a, a' \in A$. A function is said to be **injective** if it is **one-to-one**.

- To show that a function is *one-to-one*, we show that $a \neq a' \implies f(a) \neq f(a')$. (Why?)

Definition (Onto)

Definition (One-to-one)

A function f is said to be **one-to-one** if and only if $f(a) = f(a')$ implies that $a = a'$ for all $a, a' \in A$. A function is said to be **injective** if it is **one-to-one**.

- To show that a function is *one-to-one*, we show that $a \neq a' \implies f(a) \neq f(a')$. (Why?)

Definition (Onto)

A function f is called **onto**, or a surjection, if and only if for every element $b \in B$ there is an element $a \in A$ such that $f(a) = b$. We also say that f is **surjective** if it's onto.

Bijection

Definition (One-to-one)

A function f is said to be **one-to-one** if and only if $f(a) = f(a')$ implies that $a = a'$ for all $a, a' \in A$. A function is said to be **injective** if it is **one-to-one**.

- To show that a function is *one-to-one*, we show that $a \neq a' \implies f(a) \neq f(a')$. (Why?)

Definition (Onto)

A function f is called **onto**, or a surjection, if and only if for every element $b \in B$ there is an element $a \in A$ such that $f(a) = b$. We also say that f is **surjective** if it's onto.

- To show that a function is *onto*, choose $a = f^{-1}(b)$ and so $f(f^{-1}(b)) = b$.

Definition (Bijection)

Definition (Bijection)

A function f is a **bijection** if and only if it is both *one-to-one* and *onto*. We also say that f is bijective.

Definition (Bijection)

A function f is a **bijection** if and only if it is both *one-to-one* and *onto*. We also say that f is bijective.

- If $f : A \rightarrow B$ is a bijection, it will have an **inverse** function (a lemma from notes), and $|A| = |B|$.

Fermat's Little Theorem

Fermat's Little Theorem

Theorem (Fermat's Little Theorem)

Fermat's Little Theorem

Theorem (Fermat's Little Theorem)

For any prime p and any $a \in \{1, 2, \dots, p-1\}$, we have

$$a^{p-1} \equiv 1 \pmod{p}.$$

Fermat's Little Theorem

Theorem (Fermat's Little Theorem)

For any prime p and any $a \in \{1, 2, \dots, p-1\}$, we have

$$a^{p-1} \equiv 1 \pmod{p}.$$

Proof:

Fermat's Little Theorem

Theorem (Fermat's Little Theorem)

For any prime p and any $a \in \{1, 2, \dots, p-1\}$, we have

$$a^{p-1} \equiv 1 \pmod{p}.$$

Proof:

- Consider $S = \{1, 2, \dots, p-1\}$ and $S' = \{a \bmod p, 2a \bmod p, \dots, (p-1)a \bmod p\}$.

Fermat's Little Theorem

Theorem (Fermat's Little Theorem)

For any prime p and any $a \in \{1, 2, \dots, p-1\}$, we have

$$a^{p-1} \equiv 1 \pmod{p}.$$

Proof:

- Consider $S = \{1, 2, \dots, p-1\}$ and $S' = \{a \bmod p, 2a \bmod p, \dots, (p-1)a \bmod p\}$.
- They are the same set under mod p (different order).

Fermat's Little Theorem

Theorem (Fermat's Little Theorem)

For any prime p and any $a \in \{1, 2, \dots, p-1\}$, we have

$$a^{p-1} \equiv 1 \pmod{p}.$$

Proof:

- Consider $S = \{1, 2, \dots, p-1\}$ and $S' = \{a \bmod p, 2a \bmod p, \dots, (p-1)a \bmod p\}$.
- They are the same set under mod p (different order).

$$\prod_{k=1}^{p-1} k \equiv \prod_{k=1}^{p-1} ka \pmod{p}$$

Fermat's Little Theorem

Theorem (Fermat's Little Theorem)

For any prime p and any $a \in \{1, 2, \dots, p-1\}$, we have

$$a^{p-1} \equiv 1 \pmod{p}.$$

Proof:

- Consider $S = \{1, 2, \dots, p-1\}$ and $S' = \{a \bmod p, 2a \bmod p, \dots, (p-1)a \bmod p\}$.
- They are the same set under mod p (different order).

$$\prod_{k=1}^{p-1} k \equiv \prod_{k=1}^{p-1} ka \pmod{p}$$

$$(p-1)! \equiv a^{p-1}(p-1)! \pmod{p}$$

Fermat's Little Theorem

Theorem (Fermat's Little Theorem)

For any prime p and any $a \in \{1, 2, \dots, p-1\}$, we have

$$a^{p-1} \equiv 1 \pmod{p}.$$

Proof:

- Consider $S = \{1, 2, \dots, p-1\}$ and $S' = \{a \bmod p, 2a \bmod p, \dots, (p-1)a \bmod p\}$.
- They are the same set under mod p (different order).

$$\prod_{k=1}^{p-1} k \equiv \prod_{k=1}^{p-1} ka \pmod{p}$$

$$(p-1)! \equiv a^{p-1}(p-1)! \pmod{p}$$

$$a^{p-1} \equiv 1 \pmod{p}$$

Chinese Remainder Theorem

Chinese Remainder Theorem

Theorem (Chinese Remainder Theorem)

Chinese Remainder Theorem

Theorem (Chinese Remainder Theorem)

Let n_1, n_2, \dots, n_k be positive integers that are coprime to each other. Then, for any integers a_i , the system of simultaneous congruences

Chinese Remainder Theorem

Theorem (Chinese Remainder Theorem)

Let n_1, n_2, \dots, n_k be positive integers that are coprime to each other. Then, for any integers a_i , the system of simultaneous congruences

$$x \equiv a_1 \pmod{n_1}, x \equiv a_2 \pmod{n_2}, \dots, x \equiv a_k \pmod{n_k}$$

Chinese Remainder Theorem

Theorem (Chinese Remainder Theorem)

Let n_1, n_2, \dots, n_k be positive integers that are coprime to each other. Then, for any integers a_i , the system of simultaneous congruences

$$x \equiv a_1 \pmod{n_1}, x \equiv a_2 \pmod{n_2}, \dots, x \equiv a_k \pmod{n_k}$$

has a unique solution

$$x = \left(\sum_{i=1}^k a_i b_i \right) \pmod{N}$$

Chinese Remainder Theorem

Theorem (Chinese Remainder Theorem)

Let n_1, n_2, \dots, n_k be positive integers that are coprime to each other. Then, for any integers a_i , the system of simultaneous congruences

$$x \equiv a_1 \pmod{n_1}, x \equiv a_2 \pmod{n_2}, \dots, x \equiv a_k \pmod{n_k}$$

has a unique solution

$$x = \left(\sum_{i=1}^k a_i b_i \right) \bmod N$$

where $N = \prod_{i=1}^k n_i$ and $b_i = \frac{N}{n_i} \left(\frac{N}{n_i} \right)^{-1}_{n_i}$ where $\left(\frac{N}{n_i} \right)^{-1}_{n_i}$ denotes the multiplicative inverse $\pmod{n_i}$ of the integer $\frac{N}{n_i}$.

Chinese Remainder Theorem

Chinese Remainder Theorem

Proof:

To see why x is a solution, notice that for each $i = 1, 2, \dots, k$, we have

$$x \equiv a_1 y_1 z_1 + a_2 y_2 z_2 + \cdots + a_k y_k z_k \pmod{n_i}$$

Chinese Remainder Theorem

Proof:

To see why x is a solution, notice that for each $i = 1, 2, \dots, k$, we have

$$\begin{aligned} x &\equiv a_1 y_1 z_1 + a_2 y_2 z_2 + \cdots + a_k y_k z_k \pmod{n_i} \\ &\equiv a_i y_i z_i \pmod{n_i} \end{aligned}$$

Chinese Remainder Theorem

Proof:

To see why x is a solution, notice that for each $i = 1, 2, \dots, k$, we have

$$\begin{aligned} x &\equiv a_1 y_1 z_1 + a_2 y_2 z_2 + \cdots + a_k y_k z_k && (\text{mod } n_i) \\ &\equiv a_i y_i z_i && (\text{mod } n_i) \\ &\equiv a_i && (\text{mod } n_i). \end{aligned}$$

Chinese Remainder Theorem

Proof:

To see why x is a solution, notice that for each $i = 1, 2, \dots, k$, we have

$$\begin{aligned} x &\equiv a_1 y_1 z_1 + a_2 y_2 z_2 + \cdots + a_k y_k z_k && (\text{mod } n_i) \\ &\equiv a_i y_i z_i && (\text{mod } n_i) \\ &\equiv a_i && (\text{mod } n_i). \end{aligned}$$

- The second line follows since $y_j \equiv 0 \pmod{n_i}$ for each $j \neq i$.

Chinese Remainder Theorem

Proof:

To see why x is a solution, notice that for each $i = 1, 2, \dots, k$, we have

$$\begin{aligned}x &\equiv a_1 y_1 z_1 + a_2 y_2 z_2 + \cdots + a_k y_k z_k \pmod{n_i} \\&\equiv a_i y_i z_i \pmod{n_i} \\&\equiv a_i \pmod{n_i}.\end{aligned}$$

- The second line follows since $y_j \equiv 0 \pmod{n_i}$ for each $j \neq i$.
- The third line follows since $y_i z_i \equiv 1 \pmod{n_i}$.

Chinese Remainder Theorem

Proof:

To see why x is a solution, notice that for each $i = 1, 2, \dots, k$, we have

$$\begin{aligned} x &\equiv a_1 y_1 z_1 + a_2 y_2 z_2 + \cdots + a_k y_k z_k \pmod{n_i} \\ &\equiv a_i y_i z_i \pmod{n_i} \\ &\equiv a_i \pmod{n_i}. \end{aligned}$$

- The second line follows since $y_j \equiv 0 \pmod{n_i}$ for each $j \neq i$.
- The third line follows since $y_i z_i \equiv 1 \pmod{n_i}$.

Now, to prove uniqueness, suppose there are two solutions x and y .

Chinese Remainder Theorem

Proof:

To see why x is a solution, notice that for each $i = 1, 2, \dots, k$, we have

$$\begin{aligned} x &\equiv a_1 y_1 z_1 + a_2 y_2 z_2 + \cdots + a_k y_k z_k \pmod{n_i} \\ &\equiv a_i y_i z_i \pmod{n_i} \\ &\equiv a_i \pmod{n_i}. \end{aligned}$$

- The second line follows since $y_j \equiv 0 \pmod{n_i}$ for each $j \neq i$.
- The third line follows since $y_i z_i \equiv 1 \pmod{n_i}$.

Now, to prove uniqueness, suppose there are two solutions x and y .

- Then $n_1 \mid (x - y), n_2 \mid (x - y), \dots, n_k \mid (x - y)$.

Chinese Remainder Theorem

Proof:

To see why x is a solution, notice that for each $i = 1, 2, \dots, k$, we have

$$\begin{aligned}x &\equiv a_1 y_1 z_1 + a_2 y_2 z_2 + \cdots + a_k y_k z_k && (\text{mod } n_i) \\&\equiv a_i y_i z_i && (\text{mod } n_i) \\&\equiv a_i && (\text{mod } n_i).\end{aligned}$$

- The second line follows since $y_j \equiv 0 \pmod{n_i}$ for each $j \neq i$.
- The third line follows since $y_i z_i \equiv 1 \pmod{n_i}$.

Now, to prove uniqueness, suppose there are two solutions x and y .

- Then $n_1 \mid (x - y)$, $n_2 \mid (x - y)$, \dots , $n_k \mid (x - y)$.
- Since n_1, n_2, \dots, n_k are relatively prime, we have that $n_1 n_2 \cdots n_k$ divides $x - y$, or

Chinese Remainder Theorem

Proof:

To see why x is a solution, notice that for each $i = 1, 2, \dots, k$, we have

$$\begin{aligned}x &\equiv a_1 y_1 z_1 + a_2 y_2 z_2 + \cdots + a_k y_k z_k \pmod{n_i} \\&\equiv a_i y_i z_i \pmod{n_i} \\&\equiv a_i \pmod{n_i}.\end{aligned}$$

- The second line follows since $y_j \equiv 0 \pmod{n_i}$ for each $j \neq i$.
- The third line follows since $y_i z_i \equiv 1 \pmod{n_i}$.

Now, to prove uniqueness, suppose there are two solutions x and y .

- Then $n_1 \mid (x - y)$, $n_2 \mid (x - y)$, \dots , $n_k \mid (x - y)$.
- Since n_1, n_2, \dots, n_k are relatively prime, we have that $n_1 n_2 \cdots n_k$ divides $x - y$, or

$$x \equiv y \pmod{N}.$$

Chinese Remainder Theorem

Proof:

To see why x is a solution, notice that for each $i = 1, 2, \dots, k$, we have

$$\begin{aligned}x &\equiv a_1 y_1 z_1 + a_2 y_2 z_2 + \cdots + a_k y_k z_k \pmod{n_i} \\&\equiv a_i y_i z_i \pmod{n_i} \\&\equiv a_i \pmod{n_i}.\end{aligned}$$

- The second line follows since $y_j \equiv 0 \pmod{n_i}$ for each $j \neq i$.
- The third line follows since $y_i z_i \equiv 1 \pmod{n_i}$.

Now, to prove uniqueness, suppose there are two solutions x and y .

- Then $n_1 \mid (x - y)$, $n_2 \mid (x - y)$, \dots , $n_k \mid (x - y)$.
- Since n_1, n_2, \dots, n_k are relatively prime, we have that $n_1 n_2 \cdots n_k$ divides $x - y$, or

$$x \equiv y \pmod{N}.$$

Thus, the solution is unique modulo N . □

Chinese Remainder Theorem

General construction:

Chinese Remainder Theorem

General construction:

- 1 Compute $N = n_1 \times n_2 \times \cdots \times n_k$.

Chinese Remainder Theorem

General construction:

- 1 Compute $N = n_1 \times n_2 \times \cdots \times n_k$.
- 2 For each $i = 1, 2, \dots, k$, compute

$$y_i = \frac{N}{n_i} = n_1 n_2 \cdots n_{i-1} n_{i+1} \cdots n_k.$$

Chinese Remainder Theorem

General construction:

- 1 Compute $N = n_1 \times n_2 \times \cdots \times n_k$.
- 2 For each $i = 1, 2, \dots, k$, compute

$$y_i = \frac{N}{n_i} = n_1 n_2 \cdots n_{i-1} n_{i+1} \cdots n_k.$$

- 3 For each $i = 1, 2, \dots, k$, compute $z_i \equiv y_i^{-1} \pmod{n_i}$ (z_i exists since n_1, n_2, \dots, n_k are pairwise coprime).

Chinese Remainder Theorem

General construction:

① Compute $N = n_1 \times n_2 \times \cdots \times n_k$.

② For each $i = 1, 2, \dots, k$, compute

$$y_i = \frac{N}{n_i} = n_1 n_2 \cdots n_{i-1} n_{i+1} \cdots n_k.$$

③ For each $i = 1, 2, \dots, k$, compute $z_i \equiv y_i^{-1} \pmod{n_i}$ (z_i exists since n_1, n_2, \dots, n_k are pairwise coprime).

④ Compute

$$x = \sum_{i=1}^k a_i y_i z_i$$

and $x \bmod N$ is the unique solution modulo N .

Chinese Remainder Theorem

Intuitive way to solve for CRT:

Chinese Remainder Theorem

Intuitive way to solve for CRT:

- 1 Begin with the congruence with the largest modulus, $x \equiv a_k \pmod{n_k}$.

Chinese Remainder Theorem

Intuitive way to solve for CRT:

- 1 Begin with the congruence with the largest modulus, $x \equiv a_k \pmod{n_k}$.
- 2 Re-write this modulus as an equation, $x = j_k n_k + a_k$, for some positive integer j_k .

Chinese Remainder Theorem

Intuitive way to solve for CRT:

- 1 Begin with the congruence with the largest modulus, $x \equiv a_k \pmod{n_k}$.
- 2 Re-write this modulus as an equation, $x = j_k n_k + a_k$, for some positive integer j_k .
- 3 Substitute the expression for x into the congruence with the next largest modulus, $x \equiv a_k \pmod{n_k} \implies j_k n_k + a_k \equiv a_{k-1} \pmod{n_{k-1}}$.

Chinese Remainder Theorem

Intuitive way to solve for CRT:

- 1 Begin with the congruence with the largest modulus, $x \equiv a_k \pmod{n_k}$.
- 2 Re-write this modulus as an equation, $x = j_k n_k + a_k$, for some positive integer j_k .
- 3 Substitute the expression for x into the congruence with the next largest modulus, $x \equiv a_k \pmod{n_k} \implies j_k n_k + a_k \equiv a_{k-1} \pmod{n_{k-1}}$.
- 4 Solve this congruence for j_k .

Chinese Remainder Theorem

Intuitive way to solve for CRT:

- 1 Begin with the congruence with the largest modulus, $x \equiv a_k \pmod{n_k}$.
- 2 Re-write this modulus as an equation, $x = j_k n_k + a_k$, for some positive integer j_k .
- 3 Substitute the expression for x into the congruence with the next largest modulus, $x \equiv a_{k-1} \pmod{n_{k-1}} \implies j_k n_k + a_k \equiv a_{k-1} \pmod{n_{k-1}}$.
- 4 Solve this congruence for j_k .
- 5 Write the solved congruence as an equation, and then substitute this expression for j_k into the equation for x .

Chinese Remainder Theorem

Intuitive way to solve for CRT:

- 1 Begin with the congruence with the largest modulus, $x \equiv a_k \pmod{n_k}$.
- 2 Re-write this modulus as an equation, $x = j_k n_k + a_k$, for some positive integer j_k .
- 3 Substitute the expression for x into the congruence with the next largest modulus, $x \equiv a_{k-1} \pmod{n_{k-1}} \implies j_k n_k + a_k \equiv a_{k-1} \pmod{n_{k-1}}$.
- 4 Solve this congruence for j_k .
- 5 Write the solved congruence as an equation, and then substitute this expression for j_k into the equation for x .
- 6 Continue substituting and solving congruences until the equation for x implies the solution to the system of congruences.

Chinese Remainder Theorem

Example:

Chinese Remainder Theorem

Example:

$$\begin{cases} x \equiv 1 & (\text{mod } 3) \\ x \equiv 4 & (\text{mod } 5) \\ x \equiv 6 & (\text{mod } 7) \end{cases}$$

Chinese Remainder Theorem

Example:

$$\begin{cases} x \equiv 1 & (\text{mod } 3) \\ x \equiv 4 & (\text{mod } 5) \\ x \equiv 6 & (\text{mod } 7) \end{cases}$$

Solution:

Chinese Remainder Theorem

Example:

$$\begin{cases} x \equiv 1 & (\text{mod } 3) \\ x \equiv 4 & (\text{mod } 5) \\ x \equiv 6 & (\text{mod } 7) \end{cases}$$

Solution:

- Start with mod 7. Write $x = 7k + 6$.

Chinese Remainder Theorem

Example:

$$\begin{cases} x \equiv 1 & (\text{mod } 3) \\ x \equiv 4 & (\text{mod } 5) \\ x \equiv 6 & (\text{mod } 7) \end{cases}$$

Solution:

- Start with mod 7. Write $x = 7k + 6$.
- Then we have $7k + 6 \equiv 4 \pmod{5} \implies k \equiv 4 \pmod{5}$.

Chinese Remainder Theorem

Example:

$$\begin{cases} x \equiv 1 & (\text{mod } 3) \\ x \equiv 4 & (\text{mod } 5) \\ x \equiv 6 & (\text{mod } 7) \end{cases}$$

Solution:

- Start with mod 7. Write $x = 7k + 6$.
- Then we have $7k + 6 \equiv 4 \pmod{5} \implies k \equiv 4 \pmod{5}$.
- Then solving for k gives $5j + 4$.

Chinese Remainder Theorem

Example:

$$\begin{cases} x \equiv 1 & (\text{mod } 3) \\ x \equiv 4 & (\text{mod } 5) \\ x \equiv 6 & (\text{mod } 7) \end{cases}$$

Solution:

- Start with mod 7. Write $x = 7k + 6$.
- Then we have $7k + 6 \equiv 4 \pmod{5} \implies k \equiv 4 \pmod{5}$.
- Then solving for k gives $5j + 4$.
- Now we have $x = 7k + 6 = 7(5j + 4) + 6 = 35j + 34$.

Chinese Remainder Theorem

Example:

$$\begin{cases} x \equiv 1 & (\text{mod } 3) \\ x \equiv 4 & (\text{mod } 5) \\ x \equiv 6 & (\text{mod } 7) \end{cases}$$

Solution:

- Start with mod 7. Write $x = 7k + 6$.
- Then we have $7k + 6 \equiv 4 \pmod{5} \implies k \equiv 4 \pmod{5}$.
- Then solving for k gives $5j + 4$.
- Now we have $x = 7k + 6 = 7(5j + 4) + 6 = 35j + 34$.
- Then $35j + 34 \equiv 1 \pmod{3} \implies j \equiv 0 \pmod{3} \implies j = 3t$.

Chinese Remainder Theorem

Example:

$$\begin{cases} x \equiv 1 & (\text{mod } 3) \\ x \equiv 4 & (\text{mod } 5) \\ x \equiv 6 & (\text{mod } 7) \end{cases}$$

Solution:

- Start with mod 7. Write $x = 7k + 6$.
- Then we have $7k + 6 \equiv 4 \pmod{5} \implies k \equiv 4 \pmod{5}$.
- Then solving for k gives $5j + 4$.
- Now we have $x = 7k + 6 = 7(5j + 4) + 6 = 35j + 34$.
- Then $35j + 34 \equiv 1 \pmod{3} \implies j \equiv 0 \pmod{3} \implies j = 3t$.
- Finally, we have $x = 35(3t) + 34 = 105t + 34 \implies x \equiv \boxed{34} \pmod{105}$.

Problem Time!