

CS70 Counting

Kelvin Lee

kelvinlee@berkeley.edu

October 7, 2020

Overview

- 1 Rules of Counting
- 2 Stars and Bars
- 3 Binomial Theorem
- 4 Combinatorial Proofs
- 5 Principle of Inclusion-Exclusion
- 6 Problems
- 7 Summary/Tips

First Rule of Counting

First Rule of Counting

First Rule of Counting(Product Rule):

First Rule of Counting

First Rule of Counting(Product Rule):

If there are n ways of doing something, and m ways of doing another thing after that, then there are $n \times m$ ways to perform both of these actions.

First Rule of Counting

First Rule of Counting(Product Rule):

If there are n ways of doing something, and m ways of doing another thing after that, then there are $n \times m$ ways to perform both of these actions.

- **Order matters** (permutations).

First Rule of Counting

First Rule of Counting(Product Rule):

If there are n ways of doing something, and m ways of doing another thing after that, then there are $n \times m$ ways to perform both of these actions.

- **Order matters** (permutations).
- Sampling k elements from n items:

First Rule of Counting

First Rule of Counting(Product Rule):

If there are n ways of doing something, and m ways of doing another thing after that, then there are $n \times m$ ways to perform both of these actions.

- **Order matters** (permutations).
- Sampling k elements from n items:
 - **With replacement:**

First Rule of Counting

First Rule of Counting(Product Rule):

If there are n ways of doing something, and m ways of doing another thing after that, then there are $n \times m$ ways to perform both of these actions.

- **Order matters** (permutations).
- Sampling k elements from n items:
 - **With replacement:** n^k .

First Rule of Counting

First Rule of Counting(Product Rule):

If there are n ways of doing something, and m ways of doing another thing after that, then there are $n \times m$ ways to perform both of these actions.

- **Order matters** (permutations).
- Sampling k elements from n items:
 - **With replacement:** n^k .
 - **Without replacement:**

First Rule of Counting

First Rule of Counting(Product Rule):

If there are n ways of doing something, and m ways of doing another thing after that, then there are $n \times m$ ways to perform both of these actions.

- **Order matters** (permutations).
- Sampling k elements from n items:
 - **With replacement:** n^k .
 - **Without replacement:** $\frac{n!}{(n-k)!}$

Second Rule of Counting

Second Rule of Counting

Second Rule of Counting:

Second Rule of Counting

Second Rule of Counting:

If order doesn't matter count ordered objects and then divide by number of orderings.

Second Rule of Counting

Second Rule of Counting:

If order doesn't matter count ordered objects and then divide by number of orderings.

- **Without replacement** and **ordering doesn't matter** (combinations).

Second Rule of Counting

Second Rule of Counting:

If order doesn't matter count ordered objects and then divide by number of orderings.

- **Without replacement** and **ordering doesn't matter** (combinations).
- Number of ways of choosing k -element subsets out of a set of size n :

Second Rule of Counting

Second Rule of Counting:

If order doesn't matter count ordered objects and then divide by number of orderings.

- **Without replacement** and **ordering doesn't matter** (combinations).
- Number of ways of choosing k -element subsets out of a set of size n :

$$\binom{n}{k} = \frac{n!}{(n-k)!k!}.$$

Starts and Bars

Starts and Bars

Problem: Consider the equation $a + b + c + d = 12$ where a, b, c, d are non-negative integers. How many solutions are there to this equation?

Starts and Bars

Problem: Consider the equation $a + b + c + d = 12$ where a, b, c, d are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 and 3 bars.

Starts and Bars

Problem: Consider the equation $a + b + c + d = 12$ where a, b, c, d are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 and 3 bars.

★ ★ | ★ ★ | ★ ★ ★ | ★ ★ ★ ★

Starts and Bars

Problem: Consider the equation $a + b + c + d = 12$ where a, b, c, d are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 and 3 bars.

★ ★ | ★ ★ | ★ ★ ★ | ★ ★ ★ ★

- How many ways can we arrange them?

Starts and Bars

Problem: Consider the equation $a + b + c + d = 12$ where a, b, c, d are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 and 3 bars.

★ ★ | ★ ★ | ★ ★ ★ | ★ ★ ★ ★

- How many ways can we arrange them? $\binom{12+3}{3}$

Starts and Bars

Problem: Consider the equation $a + b + c + d = 12$ where a, b, c, d are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 and 3 bars.

★ ★ | ★ ★ | ★ ★ ★ | ★ ★ ★ ★

- How many ways can we arrange them? $\binom{12+3}{3} = \binom{15}{3}$

Starts and Bars

Problem: Consider the equation $a + b + c + d = 12$ where a, b, c, d are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 and 3 bars.

★ ★ | ★ ★ | ★ ★ ★ | ★ ★ ★ ★

- How many ways can we arrange them? $\binom{12+3}{3} = \binom{15}{3}$
- This is the answer to our original problem! Do you see the bijection between the two problems?

Starts and Bars

Problem: Consider the equation $a + b + c + d = 12$ where a, b, c, d are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 and 3 bars.

$$\star\star \mid \star\star \mid \star\star\star \mid \star\star\star\star$$

- How many ways can we arrange them? $\binom{12+3}{3} = \binom{15}{3}$
- This is the answer to our original problem! Do you see the bijection between the two problems?

Zeroth Rule of Counting:

Starts and Bars

Problem: Consider the equation $a + b + c + d = 12$ where a, b, c, d are non-negative integers. How many solutions are there to this equation?

- Let's simplify this problem a little bit. Suppose we have 12 and 3 bars.

★★ | ★★ | ★★★ | ★★★★★

- How many ways can we arrange them? $\binom{12+3}{3} = \binom{15}{3}$
- This is the answer to our original problem! Do you see the bijection between the two problems?

Zeroth Rule of Counting:

If a set A has a bijection relationship with a set B , then $|A| = |B|$.

Starts and Bars

Stars and Bars

Stars and Bars:

Starts and Bars

Stars and Bars:

How many ways can one distribute n indistinguishable objects into k distinguishable bins?

Stars and Bars:

How many ways can one distribute n indistinguishable objects into k distinguishable bins?

$$\binom{n + k - 1}{k - 1}$$

Stars and Bars

Stars and Bars:

How many ways can one distribute n indistinguishable objects into k distinguishable bins?

$$\binom{n + k - 1}{k - 1}$$

- Useful for **with replacement** but **order doesn't matter** type of problems.

Binomial Theorem

Binomial Theorem

Binomial Theorem:

Binomial Theorem

Binomial Theorem:

For all $n \in \mathbb{N}$,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

Binomial Theorem

Binomial Theorem:

For all $n \in \mathbb{N}$,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

Proof:

Binomial Theorem

Binomial Theorem:

For all $n \in \mathbb{N}$,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

Proof:

See notes. □

Binomial Theorem

Binomial Theorem:

For all $n \in \mathbb{N}$,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

Proof:

See notes. □

Corollary:

Binomial Theorem

Binomial Theorem:

For all $n \in \mathbb{N}$,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

Proof:

See notes. □

Corollary:

$$\sum_{k=0}^n (-1)^k \binom{n}{k} = 0.$$

Binomial Theorem

Binomial Theorem:

For all $n \in \mathbb{N}$,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

Proof:

See notes. □

Corollary:

$$\sum_{k=0}^n (-1)^k \binom{n}{k} = 0.$$

Proof:

Binomial Theorem

Binomial Theorem:

For all $n \in \mathbb{N}$,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

Proof:

See notes. □

Corollary:

$$\sum_{k=0}^n (-1)^k \binom{n}{k} = 0.$$

Proof:

Plug in $a = -1$ and $b = 1$ for the binomial theorem.

Binomial Theorem

Binomial Theorem:

For all $n \in \mathbb{N}$,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}.$$

Proof:

See notes. □

Corollary:

$$\sum_{k=0}^n (-1)^k \binom{n}{k} = 0.$$

Proof:

Plug in $a = -1$ and $b = 1$ for the binomial theorem. □

Combinatorial Proofs

Combinatorial Proofs

- Intuitive counting arguments. No tedious algebraic manipulation.

Combinatorial Proofs

- Intuitive counting arguments. No tedious algebraic manipulation.
- Proofs by stories: same story from multiple perspectives.

Combinatorial Proofs

- Intuitive counting arguments. No tedious algebraic manipulation.
- Proofs by stories: same story from multiple perspectives.
- Proving an identity by counting the same thing in two different ways.

Combinatorial Proofs

- Intuitive counting arguments. No tedious algebraic manipulation.
- Proofs by stories: same story from multiple perspectives.
- Proving an identity by counting the same thing in two different ways.
- Useful identity:

Combinatorial Proofs

- Intuitive counting arguments. No tedious algebraic manipulation.
- Proofs by stories: same story from multiple perspectives.
- Proving an identity by counting the same thing in two different ways.
- Useful identity:

$$\binom{n}{k} = \binom{n}{n-k}.$$

Combinatorial Proofs

- Intuitive counting arguments. No tedious algebraic manipulation.
- Proofs by stories: same story from multiple perspectives.
- Proving an identity by counting the same thing in two different ways.
- Useful identity:

$$\binom{n}{k} = \binom{n}{n-k}.$$

- Choosing k objects to include is equivalent to choosing $n - k$ objects to exclude.

Example

Combinatorial Identity:

Combinatorial Identity:

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

Example

Combinatorial Identity:

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

Proof:

Example

Combinatorial Identity:

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

Proof:

Although we can use binomial theorem by letting $a = b = 1$, we use combinatorial argument to prove this.

Example

Combinatorial Identity:

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

Proof:

Although we can use binomial theorem by letting $a = b = 1$, we use combinatorial argument to prove this.

RHS:

Example

Combinatorial Identity:

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

Proof:

Although we can use binomial theorem by letting $a = b = 1$, we use combinatorial argument to prove this.

RHS: Total number of subsets of a set of size n .

Example

Combinatorial Identity:

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

Proof:

Although we can use binomial theorem by letting $a = b = 1$, we use combinatorial argument to prove this.

RHS: Total number of subsets of a set of size n .

LHS:

Example

Combinatorial Identity:

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

Proof:

Although we can use binomial theorem by letting $a = b = 1$, we use combinatorial argument to prove this.

RHS: Total number of subsets of a set of size n .

LHS: The number of ways to choose a subset of size i is $\binom{n}{i}$. To find the total number of subsets, we simply add all the cases when $i = 0, 1, 2, \dots, n$.

□

Example

Principle of Inclusion-Exclusion(General):

Example

Principle of Inclusion-Exclusion(General):

Let A_1, \dots, A_n be arbitrary subsets of the same finite set A . Then,

$$|A_1 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k-1} \sum_{S \subseteq \{1, \dots, n\}: |S|=k} |\cap_{i \in S} A_i|.$$

Example

Principle of Inclusion-Exclusion(General):

Let A_1, \dots, A_n be arbitrary subsets of the same finite set A . Then,

$$|A_1 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k-1} \sum_{S \subseteq \{1, \dots, n\}: |S|=k} |\cap_{i \in S} A_i|.$$

Proof:

Example

Principle of Inclusion-Exclusion(General):

Let A_1, \dots, A_n be arbitrary subsets of the same finite set A . Then,

$$|A_1 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k-1} \sum_{S \subseteq \{1, \dots, n\}: |S|=k} |\cap_{i \in S} A_i|.$$

Proof:

See notes.



Example

Principle of Inclusion-Exclusion(General):

Let A_1, \dots, A_n be arbitrary subsets of the same finite set A . Then,

$$|A_1 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k-1} \sum_{S \subseteq \{1, \dots, n\}: |S|=k} |\cap_{i \in S} A_i|.$$

Proof:

See notes.



Principle of Inclusion-Exclusion(Simplified):

Example

Principle of Inclusion-Exclusion(General):

Let A_1, \dots, A_n be arbitrary subsets of the same finite set A . Then,

$$|A_1 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k-1} \sum_{S \subseteq \{1, \dots, n\}: |S|=k} |\cap_{i \in S} A_i|.$$

Proof:

See notes.



Principle of Inclusion-Exclusion(Simplified):

$$|A \cup B| = |A| + |B| - |A \cap B|.$$

Problems

SP19 MT2 6.3

SP19 MT2 6.3

How many permutations of the numbers 1 through n are there such that 1 comes before 2 and after 3? Assume $n > 3$.

SP19 MT2 6.3

How many permutations of the numbers 1 through n are there such that 1 comes before 2 and after 3? Assume $n > 3$.

Hint:

SP19 MT2 6.3

How many permutations of the numbers 1 through n are there such that 1 comes before 2 and after 3? Assume $n > 3$.

Hint: how many ways to choose the positions for the three numbers? What do we do with the remaining numbers?

SP19 MT2 6.3

How many permutations of the numbers 1 through n are there such that 1 comes before 2 and after 3? Assume $n > 3$.

Hint: how many ways to choose the positions for the three numbers? What do we do with the remaining numbers?

Solution:

SP19 MT2 6.3

How many permutations of the numbers 1 through n are there such that 1 comes before 2 and after 3? Assume $n > 3$.

Hint: how many ways to choose the positions for the three numbers? What do we do with the remaining numbers?

Solution:

- There are $\binom{n}{3}$ ways to pick positions for 1, 2, 3. For the positions picked, we place the three numbers in a way such that the conditions are met, i.e, we place them in the order of 3, 1, 2.

SP19 MT2 6.3

How many permutations of the numbers 1 through n are there such that 1 comes before 2 and after 3? Assume $n > 3$.

Hint: how many ways to choose the positions for the three numbers? What do we do with the remaining numbers?

Solution:

- There are $\binom{n}{3}$ ways to pick positions for 1, 2, 3. For the positions picked, we place the three numbers in a way such that the conditions are met, i.e, we place them in the order of 3, 1, 2.
- Now for the remaining numbers, there are $(n - 3)!$ to arrange them.

SP19 MT2 6.3

How many permutations of the numbers 1 through n are there such that 1 comes before 2 and after 3? Assume $n > 3$.

Hint: how many ways to choose the positions for the three numbers? What do we do with the remaining numbers?

Solution:

- There are $\binom{n}{3}$ ways to pick positions for 1, 2, 3. For the positions picked, we place the three numbers in a way such that the conditions are met, i.e, we place them in the order of 3, 1, 2.
- Now for the remaining numbers, there are $(n - 3)!$ to arrange them.
- Finally, by the **first rule of counting**, we have $\boxed{\frac{n!}{6}}$ permutations.

Problems

SP18 MT2 5

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

Hint:

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

Hint: how many edges are there? How do we choose them?

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

Hint: how many edges are there? How do we choose them?

Solution:

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

Hint: how many edges are there? How do we choose them?

Solution:

- If every vertex has degree 1, then we can only have 3 edges.

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

Hint: how many edges are there? How do we choose them?

Solution:

- If every vertex has degree 1, then we can only have 3 edges.
- Each edge requires 2 vertices, so $\binom{6}{2} = 15$ ways to choose an edge.

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

Hint: how many edges are there? How do we choose them?

Solution:

- If every vertex has degree 1, then we can only have 3 edges.
- Each edge requires 2 vertices, so $\binom{6}{2} = 15$ ways to choose an edge.
- After choosing the first edge, we have 4 vertices remaining, so there are $\binom{4}{2} = 6$ ways to choose the second edge and similarly $\binom{2}{2} = 1$ way to choose the final edge.

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(a) How many such graphs are there such that all vertices have degree 1?

Hint: how many edges are there? How do we choose them?

Solution:

- If every vertex has degree 1, then we can only have 3 edges.
- Each edge requires 2 vertices, so $\binom{6}{2} = 15$ ways to choose an edge.
- After choosing the first edge, we have 4 vertices remaining, so there are $\binom{4}{2} = 6$ ways to choose the second edge and similarly $\binom{2}{2} = 1$ way to choose the final edge.
- However, since order doesn't matter, by the **second rule of counting**, we divide by $3! = 6$. So our final answer is 15.

Problems

SP18 MT2 5

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(b) How many ways can we form two disjoint cycles of length 3 with 6 vertices?

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(b) How many ways can we form two disjoint cycles of length 3 with 6 vertices?

Hint:

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(b) How many ways can we form two disjoint cycles of length 3 with 6 vertices?

Hint: how many ways to pick two groups?

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(b) How many ways can we form two disjoint cycles of length 3 with 6 vertices?

Hint: how many ways to pick two groups?

Solution:

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(b) How many ways can we form two disjoint cycles of length 3 with 6 vertices?

Hint: how many ways to pick two groups?

Solution:

- We are choosing two sets of 3 vertices. There are $\binom{6}{3} \binom{3}{3} = 20$ ways.

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(b) How many ways can we form two disjoint cycles of length 3 with 6 vertices?

Hint: how many ways to pick two groups?

Solution:

- We are choosing two sets of 3 vertices. There are $\binom{6}{3} \binom{3}{3} = 20$ ways.
- But order doesn't matter here again. So we divide by $2!$. Thus, the answer is 10.

Problems

SP18 MT2 5

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

Hint:

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

Hint: how many ways can we permute the vertices?

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

Hint: how many ways can we permute the vertices?

Solution:

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

Hint: how many ways can we permute the vertices?

Solution:

- We think of the cycle as a permutation of the vertices, which has $6!$ possibilities.

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

Hint: how many ways can we permute the vertices?

Solution:

- We think of the cycle as a permutation of the vertices, which has $6!$ possibilities.
- However, it doesn't matter where we start, so divide by 6.

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

Hint: how many ways can we permute the vertices?

Solution:

- We think of the cycle as a permutation of the vertices, which has $6!$ possibilities.
- However, it doesn't matter where we start, so divide by 6.
- The direction in which we travel along the cycle also doesn't matter, so divide by 2.

SP18 MT2 5

We wish to count how many undirected graphs on 6 vertices with equal degrees there are.

(c) How many ways can we form a long cycle of length 6?

Hint: how many ways can we permute the vertices?

Solution:

- We think of the cycle as a permutation of the vertices, which has $6!$ possibilities.
- However, it doesn't matter where we start, so divide by 6.
- The direction in which we travel along the cycle also doesn't matter, so divide by 2.
- Thus, our answer is $\frac{6!}{2 \cdot 6} = \boxed{60}$.

Problems

SP17 MT2 4

SP17 MT2 4

What are the number of ways to divide m dollar bills among z people?

SP17 MT2 4

What are the number of ways to divide m dollar bills among z people?

Hint:

SP17 MT2 4

What are the number of ways to divide m dollar bills among z people?

Hint: Stars and bars.

SP17 MT2 4

What are the number of ways to divide m dollar bills among z people?

Hint: Stars and bars.

Solution:

SP17 MT2 4

What are the number of ways to divide m dollar bills among z people?

Hint: Stars and bars.

Solution:

- This is a stars and bars problem where we have $z - 1$ bars and m stars.

SP17 MT2 4

What are the number of ways to divide m dollar bills among z people?

Hint: Stars and bars.

Solution:

- This is a stars and bars problem where we have $z - 1$ bars and m stars.
- So $n = m$ and $k = z$ in this case.

SP17 MT2 4

What are the number of ways to divide m dollar bills among z people?

Hint: Stars and bars.

Solution:

- This is a stars and bars problem where we have $z - 1$ bars and m stars.
- So $n = m$ and $k = z$ in this case.

- Thus, the answer is $\binom{n+k-1}{k-1} = \boxed{\binom{m+z-1}{z-1}}.$

Problems

SP16 MT2 6

SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

Hint:

SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

Hint: Ternary strings.

SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

Hint: Ternary strings.

Solution:

SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

Hint: Ternary strings.

Solution:

- **LHS:**

SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

Hint: Ternary strings.

Solution:

- **LHS:** the number of ternary strings of length n .

SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

Hint: Ternary strings.

Solution:

- **LHS:** the number of ternary strings of length n .
- **RHS:**

SP16 MT2 6

Give a combinatorial proof for

$$3^n = \sum_{i=0}^n \binom{n}{i} 2^{n-i}$$

Hint: Ternary strings.

Solution:

- **LHS:** the number of ternary strings of length n .
- **RHS:** There are $\binom{n}{i}$ positions of the 2's, and there are 2^{n-i} possible patterns of 0 and 1's in the remaining positions. The sum gives you all the ternary strings.

Summary/Tips

Summary:

Summary:

Summary/Tips

Summary:

	with replacement	w/o replacement
order matters	n^k	$\frac{n!}{(n-k)!}$
order doesn't matter	$\binom{n+k-1}{k-1}$	$\binom{n}{k}$

Tips:

Summary/Tips

Summary:

	with replacement	w/o replacement
order matters	n^k	$\frac{n!}{(n-k)!}$
order doesn't matter	$\binom{n+k-1}{k-1}$	$\binom{n}{k}$

Tips:

- Don't memorize formulas. Understand them by counting.

Summary/Tips

Summary:

	with replacement	w/o replacement
order matters	n^k	$\frac{n!}{(n-k)!}$
order doesn't matter	$\binom{n+k-1}{k-1}$	$\binom{n}{k}$

Tips:

- Don't memorize formulas. Understand them by counting.
- Identify which categories does the problem fall under.

Summary/Tips

Summary:

	with replacement	w/o replacement
order matters	n^k	$\frac{n!}{(n-k)!}$
order doesn't matter	$\binom{n+k-1}{k-1}$	$\binom{n}{k}$

Tips:

- Don't memorize formulas. Understand them by counting.
- Identify which categories does the problem fall under.
- Double check answers by using two different counting approaches.

Summary/Tips

Summary:

	with replacement	w/o replacement
order matters	n^k	$\frac{n!}{(n-k)!}$
order doesn't matter	$\binom{n+k-1}{k-1}$	$\binom{n}{k}$

Tips:

- Don't memorize formulas. Understand them by counting.
- Identify which categories does the problem fall under.
- Double check answers by using two different counting approaches.
- Check for overcounting.

Summary/Tips

Summary:

	with replacement	w/o replacement
order matters	n^k	$\frac{n!}{(n-k)!}$
order doesn't matter	$\binom{n+k-1}{k-1}$	$\binom{n}{k}$

Tips:

- Don't memorize formulas. Understand them by counting.
- Identify which categories does the problem fall under.
- Double check answers by using two different counting approaches.
- Check for overcounting.
- Relax and have fun!