
STOCHASTIC PROCESSES

STAT 150

Instructor: Benson Au

KELVIN LEE

UC BERKELEY

Contents

1	Probability Review	3
1.1	Basic Definitions	3
1.2	Overview	3
1.3	Useful Properties	4
1.4	Conditional Probability	4
1.5	Random Variables	5
1.5.1	Discrete Random Variables	5
1.5.2	Indicator Random Variable	5
1.5.3	Continuous Random Variables	5
1.5.4	Cumulative Distribution Function (CDF)	6
1.5.5	Expectation	6
1.5.6	Variance	6
1.5.7	Moments	7
1.5.8	Joint Distribution	7
1.5.9	Independence	7
1.5.10	Linearity of Expectation	7
1.5.11	Convolution	8
1.5.12	Gamma Distribution	9
1.5.13	Moment Generating Function	9
1.6	Conditional Probability (Cont'd)	10
1.6.1	Conditional Expectation	11
2	Random Sums	13

1 Probability Review

1.1 Basic Definitions

Definition 1.1.1 (Probability Space). A *probability space* $(\Omega, \mathcal{F}, \mathbb{P})$ is a triple consisting of a set Ω called the *sample space*, a set $\mathcal{F} \subseteq \Omega$ satisfying certain closure properties, and a function $\mathbb{P} : \mathcal{F} \rightarrow [0, 1]$ that assigns probabilities to events in a coherent way.

Requirements for \mathcal{F} :

- (i) $\Omega \in \mathcal{F}$.
- (ii) If $E \in \mathcal{F}$, then $E^c \in \mathcal{F}$.
- (iii) If $\{E_i\}_{i=1}^{\infty} \subseteq \mathcal{F}$, then

$$\bigcap_{i=1}^{\infty} E_i \in \mathcal{F}.$$

Requirements for \mathbb{P} :

- (i) $\mathbb{P}(\Omega) = 1$.
- (ii) If $\{E_i\}_{i=1}^{\infty} \subseteq \mathcal{F}$ are pairwise disjoint (meaning $E_i \cap E_j = \emptyset$ for $i \neq j$), then

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(E_i)$$

Definition 1.1.2 (Random Variable). A *random variable* is a function $X : \Omega \rightarrow \mathbb{R}$ such that $X^{-1}(B) \in \mathcal{F}$ whenever B is a "nice" subset of \mathbb{R} .

Example 1.1.3. $\Omega = \{H, T\}$, $\mathcal{F} = 2^{\Omega}$, $\mathbb{P}(\{H\}) = \frac{1}{2}$. $X(H) = 1$, $X(T) = 0$.

$$\mathbb{P}(X = 1) = \mathbb{P}(\{H\}) = \frac{1}{2}, \quad \mathbb{P}(X = 0) = \mathbb{P}(\{T\}) = \frac{1}{2}.$$

1.2 Overview

Definition 1.2.1 (Stochastic Process). A *stochastic process* is a collection $\{X_t : t \in T\}$ of random variables $X_t : \Omega \rightarrow S \subseteq \mathbb{R}$ all defined on the same probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Here T is some index set (typically representing time) and S is the *state space*. One writes this as

$X : \Omega \times T \rightarrow S$, $(\omega, t) \mapsto X_t(\omega)$. For a given outcome $\omega \in \Omega$, we get a sample path trajectory $X(\omega) : T \rightarrow S, t \mapsto X_t(\omega)$. A stochastic process can then be thought of as a random function.

The theme of this course is what can we say about the distribution of trajectories?

Example 1.2.2 (Branching Process (DTDS)). $X_0 = 1$, one individual in the 0th generation individuals produce a random number of offspring, i.i.d. $(\xi_i^{(n)})_{i \in \mathbb{N}, n \in \mathbb{N}_0}$.

$$X_{n+1} = \sum_{i=1}^{X_n} \xi_i^{(n)}.$$

One interesting question would be what is $\mathbb{P}(X_n = 0 \text{ eventually})$, the probability of dying out?

Example 1.2.3 (Poisson Process (CTDS)). Recall that the Poisson distribution is used to model the number of occurrences of a rare event in some fixed period of time. The Poisson process $(N_t)_{t \geq 0}$ models the number of occurrences throughout time. $N_t = \#$ of occurrences by time t .

1.3 Useful Properties

(i) (*DeMorgan*)

$$(E \cup F)^c = E^c \cap F^c, \quad (E \cap F)^c = E^c \cup F^c.$$

(ii) (*Complementation*)

$$\mathbb{P}(E) = 1 - \mathbb{P}(E^c).$$

(iii) (*Inclusion-exclusion*)

$$\begin{aligned} \mathbb{P}(E \cup F) &= \mathbb{P}(E) + \mathbb{P}(F) - \mathbb{P}(E \cap F) \\ \mathbb{P}\left(\bigcup_{i=1}^n E_i\right) &= \sum_{j=1}^n (-1)^{j-1} \sum_{S \in [n]: |S|=j} \mathbb{P}\left(\bigcap_{i \in S} E_i\right). \end{aligned}$$

(iv) (*Partitioning*) If $\bigsqcup_{i=1}^{\infty} E_i = \Omega$, then

$$\mathbb{P}(F) = \mathbb{P}\left(\bigsqcup_{i=1}^{\infty} (F \cap E_i)\right) = \sum_{i=1}^{\infty} \mathbb{P}(F \cap E_i)$$

1.4 Conditional Probability

Conditioning: For $\mathbb{P}(F) > 0$,

$$\mathbb{P}(E | F) = \frac{\mathbb{P}(E \cap F)}{\mathbb{P}(F)}.$$

$\mathbb{P}(\cdot | F)$ defines a new probability measure on (Ω, \mathcal{F}) .

Multiplication rule:

$$\mathbb{P}(E \cap F) = \mathbb{P}(F) \mathbb{P}(E | F).$$

If $\bigcup_{i=1}^{\infty} F_i = \Omega$, then

$$\mathbb{P}(E) = \sum_{i=1}^{\infty} \mathbb{P}(E \cap F_i) = \sum_{i=1}^{\infty} \mathbb{P}(F_i) \mathbb{P}(E | F_i).$$

Bayes' rule:

$$\mathbb{P}(F_j | E) = \frac{\mathbb{P}(F_j) \mathbb{P}(E | F_j)}{\sum_{i=1}^{\infty} \mathbb{P}(F_i) \mathbb{P}(E | F_i)}$$

1.5 Random Variables

1.5.1 Discrete Random Variables

If $X : \Omega \rightarrow S \subseteq \mathbb{R}$ is discrete,

$$\mathbb{P}(X \in E) = \sum_{x \in E} \mathbb{P}(X = x) = \sum_{x \in E \cap S} \mathbb{P}(X = x).$$

1.5.2 Indicator Random Variable

$$X(\omega) = \mathbb{I}_E(\omega) = \begin{cases} 1 & \omega \in E \\ 0 & \omega \notin E. \end{cases}$$

1.5.2.1 Binomial Random Variable

$$X = \sum_{i=1}^n \mathbb{I}_{E_i}, \quad \mathbb{P}(E_i) = p.$$

$$p_X(x) = \mathbb{P}(X = x) = \binom{n}{x} p^x (1-p)^{n-x}.$$

1.5.3 Continuous Random Variables

If X continuous,

$$\mathbb{P}(X \in E) = \int_E f_X(x) dx.$$

$$\mathbb{P}(X \in [a, b]) = \int_a^b f_X(x) dx.$$

1.5.3.1 Exponential Random Variable

$$f_X(x) = \lambda e^{-\lambda x} \mathbb{I}_{x \geq 0}.$$

1.5.3.2 Gaussian Random Variable

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(x-\mu)^2/2\sigma^2}.$$

1.5.4 Cumulative Distribution Function (CDF)

$$F_X : \mathbb{R} \rightarrow [0, 1],$$

$$F_X(r) = \mathbb{P}(X \leq r) = \mathbb{P}(X \in (-\infty, r]).$$

If X is discrete,

$$F_X(r) = \sum_{x_i \leq r} p_X(x_i).$$

If X is continuous,

$$F_X(r) = \int_{-\infty}^r f_X(x) dx.$$

$$\frac{d}{dr} F_X(r) = f_X(r).$$

1.5.5 Expectation

1.5.5.1 Discrete case

$$\mathbb{E}[X] = \sum_{x_i \in S} x_i \mathbb{P}(X = x_i).$$

$$\mathbb{E}[X] = \sum_{n=1}^{\infty} \mathbb{P}(X \geq n).$$

$$\mathbb{E}[g(X)] = \sum_{x_i \in S} g(x_i) \mathbb{P}(X = x_i).$$

1.5.5.2 Continuous case

$$\mathbb{E}[X] = \int_{-\infty}^{\infty} x f_X(x) dx.$$

$$\mathbb{E}[X] = \int_{-\infty}^{\infty} x \mathbb{P}(X \geq x) dx.$$

$$\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx.$$

1.5.6 Variance

$$\text{var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2.$$

1.5.7 Moments

$$\mathbb{E}[X^m] = \int_0^\infty mx^{m-1}\mathbb{P}(X \geq x)dx.$$

1.5.8 Joint Distribution

1.5.8.1 Discrete

$$p_{X,Y}(x,y) = \mathbb{P}(X = x, Y = y)$$

1.5.8.2 Continuous

$$\mathbb{P}((X,Y) \in E) = \int \int_E f_{X,Y}(x,y)dx dy$$

1.5.8.3 Marginal Distribution

$$p_X(x) = \sum_{y \in S_Y} p_{X,Y}(x,y).$$

$$f_X(x) = \int_{y \in S_y} f_{X,Y}(x,y)dy$$

1.5.9 Independence

$$p_{X,Y}(x,y) = p_X(x)p_Y(y).$$

$$f_{X,Y}(x,y) = f_X(x)f_Y(y).$$

$$\mathbb{P}(X \leq x, Y \leq y) = F_X(x)F_Y(y).$$

1.5.10 Linearity of Expectation

$$\mathbb{E} \left[\sum_{i=1}^n c_i X_i \right] = \sum_{i=1}^n c_i \mathbb{E}[X_i]$$

If $(X_i)_{i=1}^n$ independent,

$$(g(X_i))_{i=1}^n$$

independent.

$$\mathbb{E} \left[\prod_{i=1}^n g(X_i) \right] = \prod_{i=1}^n \mathbb{E} [g(x_i)]$$

$$\text{Var} \left(\sum_{i=1}^n x_i \right) = \sum_{i=1}^n \text{Var}(x_i)$$

In general,

$$\text{Var} \left(\sum_{i=1}^n x_i \right) = \sum_{i,j=1}^n \text{Cov}(x_i, x_j)$$

1.5.11 Convolution

Discrete case: X, Y discrete $X \perp\!\!\!\perp Y$

$$\begin{aligned} \mathbb{P}(X + Y = z) &= \sum_Y \mathbb{P}(X + Y = z, Y = y) \\ &= \sum_y \mathbb{P}(X = z - y, Y = y) \\ &= \sum_y \mathbb{P}(X = z - y) \mathbb{P}(Y = y) \quad (= \sum_x \mathbb{P}(X = x) \mathbb{P}(Y = z - x)). \end{aligned}$$

If X, Y are \mathbb{Z} -valued, this becomes

$$\begin{aligned} \mathbb{P}(X + Y = n) &= \sum_{k=-\infty}^{\infty} \mathbb{P}(X = n - k) \mathbb{P}(Y = k) \\ &= \sum_{k=-\infty}^{\infty} \mathbb{P}(X = k) \mathbb{P}(Y = n - k) \\ &= (\mathbb{P}_X * \mathbb{P}_Y)(n) \end{aligned}$$

Example 1.5.1 (Poisson). $X \sim \text{Poisson}(\lambda)$, $Y \sim \text{Poisson}(\mu)$, $X + Y \sim \text{Poisson}(\lambda + \mu)$

$$\begin{aligned} \mathbb{P}(X + Y = n) &= \sum_{k=0}^n \mathbb{P}(X = k) \mathbb{P}(Y = n - k) \\ &= \sum_{k=0}^n e^{-\lambda} \frac{\lambda^k}{k!} e^{-\mu} \frac{\mu^{n-k}}{(n-k)!} \\ &= e^{-(\lambda+\mu)} \sum_{k=0}^n \frac{n!}{k!(n-k)!} \lambda^k \mu^{n-k} \\ &= \frac{e^{-(\lambda+\mu)}}{n!} (\lambda + \mu)^n \\ &= \mathbb{P}(Z = n) \end{aligned}$$

where $Z \sim \text{Poisson}(\lambda + \mu)$.

Continuous case: X, Y continuous

$$\begin{aligned} \mathbb{P}(X + Y \leq z) &= \int_{-\infty}^{\infty} \int_{-\infty}^{z-x} f_X(x) f_Y(y) dy dx \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^z f_X(x) f_Y(y - x) dy dx \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_X(x) f_Y(y - x) dx dy \end{aligned}$$

$$f_{X+Y}(z) = \int_{-\infty}^{\infty} f_X(x) f_Y(z - x) dx = \int_{-\infty}^{\infty} f_Y(y) f_X(z - y) dy = f_X * f_Y.$$

Example 1.5.2 (Convolution in uniform distributions). $X, Y \sim U[0, 1]$, $X \perp\!\!\!\perp Y$.

$$f_{X+Y}(z) = \int_{-\infty}^{\infty} f_X(x)f_Y(z-x)dx.$$

$$f_X(x) = \mathbb{I}_{[0,1]}(x) \quad f_Y(y) = \mathbb{I}_{[0,1]}(y)$$

so

$$\begin{aligned} f_{X+Y}(z) &= \int_{x \in [0,1], z-x \in [0,1]} 1dx \\ &= \int_{x \in [0,1], x \in [-1+z, z]} 1dx \\ &= \int_{\max(0, -1+z)}^{\min(1, z)} 1dx \\ &= \min(1, z) - \max(0, -1+z). \end{aligned}$$

1.5.12 Gamma Distribution

Definition 1.5.3 (Gamma function). Let $\alpha > 0$. The *gamma function* $\Gamma : (0, \infty) \rightarrow (0, \infty)$ is defined by

$$\Gamma(\alpha) = \int_0^{\infty} x^{\alpha-1} e^{-x} dx = \mathbb{E}[X^{\alpha-1}]$$

where $X \sim \text{Exp}(1)$. Let $\alpha, \lambda > 0$. The $\text{Gamma}(\alpha, \lambda)$ distribution is defined by

$$f(x) = \frac{\lambda^{\alpha} x^{\alpha-1}}{\Gamma(\alpha)} e^{-\lambda x} \mathbb{I}_{x \geq 0}.$$

Exercise 1.5.4. $\Gamma(n) = (n-1)!$ for $n \in \mathbb{N}$. (Hint: use induction)

1.5.13 Moment Generating Function

Definition 1.5.5 (MGF). For a random variable X , the *moment generating function* (MGF) is the function $M_X : \mathbb{R} \rightarrow \mathbb{R}_{\geq 0} \cup \{+\infty\}$,

$$M_X(t) = \mathbb{E}[e^{tX}].$$

If $M_X(t) < +\infty$ for $t \in (-\epsilon, \epsilon)$, then

$$M_X(t) = \sum_{k=0}^{\infty} \frac{t^k \mathbb{E}[x^k]}{k!} \text{ for } |t| < \epsilon$$

For independent RVs $(X_i)_{i=1}^n$,

$$M_{\sum_{i=1}^n X_i}(t) = \prod_{i=1}^n M_{X_i}(t)$$

Exercise 1.5.6. If $X \sim \text{Exp}(\lambda)$, then $M_X(t) = \frac{\lambda}{\lambda-t}$ if $t < \lambda$, $+\infty$ otherwise.

If $X \sim \text{Gamma}(n, \lambda)$, then

$$M_X(t) = \left(\frac{\lambda}{\lambda-t} \right)^n.$$

If $X \sim \text{Gamma}(\alpha, \lambda)$, then

$$M_X(t) = \left(\frac{\lambda}{\lambda-t} \right)^{\alpha}.$$

1.6 Conditional Probability (Cont'd)

Exercise 1.6.1 (Generalization). $(X_i)_{i=1}^n, (Y_j)_{j=1}^m$

$$p_{X_1, \dots, X_n | Y_1, \dots, Y_m} (x_1, \dots, x_n | y_1, \dots, y_m) = ?$$

Example 1.6.2. Let $M \in \mathbb{N}$ and $p, q \in (0, 1)$. Consider $N \sim \text{Bin}(M, q)$ and $X \sim \text{Bin}(N, p)$. What is the distribution of X ?

$$\begin{aligned} \mathbb{P}(X = k) &= \sum_{n=0}^M \mathbb{P}(N = n) \mathbb{P}(X = k | N = n) \\ &= \sum_{n=0}^M \binom{M}{n} q^n (1-q)^{M-n} \binom{n}{k} p^k (1-p)^{n-k} \\ &= \frac{p^k}{k!} \sum_{n=k}^M \frac{M!}{(M-n)!(n-k)!} q^n (1-q)^{M-n} (1-p)^{n-k} \\ &= \frac{p^k}{k!(M-k)!} \sum_{n=k}^M \frac{M!(M-k)!}{(M-n)!(n-k)!} q^n (1-q)^{M-n} (1-p)^{n-k} \\ &= \binom{M}{k} p^k q^k \sum_{n=k}^M \binom{M-k}{n-k} q^{n-k} (1-q)^{M-j} (1-p)^{n-k} \\ &= \binom{M}{k} p^k q^k \sum_{t=0}^{M-k} \binom{M-k}{t} (q(1-p))^t (1-q)^{M-k-t} \\ &= \binom{M}{k} (pq)^k (q(1-p) + (1-q))^{M-k} \\ &= \binom{M}{k} (pq)^k (1-pq)^{M-k}. \end{aligned}$$

Thus, $X \sim \text{Bin}(M, pq)$.

Remark. What if $k > n$ in $\mathbb{P}(X = k | N = n)$ above in the first line? The probability is simply 0.

Question. Why does this answer make sense?

Answer. Think about retesting whenever we succeeded for the first M trials. Then X is simply the number of trials with double successes, thus we have the pq parameter.

Exercise 1.6.3. Consider $N \sim \text{Poisson}(\lambda)$, $X \sim \text{Bin}(N, p)$. What is the distribution of X ?

Answer. $X \sim \text{Poisson}(\lambda p)$.

Question. How can we interpret this?

Answer. We can interpret X as the number of customers visiting a store who purchase something.

1.6.1 Conditional Expectation

For X, Y discrete, $g : \mathbb{R} \rightarrow \mathbb{R}$. Assume $\mathbb{E}[|g(X)|] = \sum_x |g(x)p_X(x)| < \infty$.

Definition 1.6.4 (Conditional expectation). The *conditional expectation* is defined as

$$\mathbb{E}[g(X) | Y = y] = \sum_x g(x)p_{X|Y}(x|y)$$

if $p_Y(y) \neq 0$.

Remark. Note that $\mathbb{E}[g(X) | Y = y]$ is a real number, whereas $\mathbb{E}[g(X) | Y]$ is a random variable.

1.6.1.1 Tower Property

$$\begin{aligned} \mathbb{E}[\mathbb{E}[g(X) | Y]] &= \mathbb{E}\left[\sum_y \mathbb{E}[g(X) | Y = y]p_Y(y)\right] \\ &= \sum_y \mathbb{E}[g(X) | Y = y]p_Y(y) \\ &= \sum_y \sum_x g(x)p_{X|Y}(x|y)p_Y(y) \\ &= \sum_x g(x) \sum_y p_{X|Y}(x|y)p_Y(y) \\ &= \sum_x g(x)p_X(x) \\ &= \mathbb{E}[g(X)]. \end{aligned}$$

Remark. One intuitive example would be considering the averages of heights of students from a classroom. We divide it into several groups and let Y denote the whichever group we select and let $\mathbb{E}[g(X) | Y]$ be the average of those from group Y . Then the average height of the entire classroom $\mathbb{E}[g(X)]$ is equivalent to the average of the average of heights of each group, which is $\mathbb{E}[\mathbb{E}[g(X) | Y]]$.

Properties of conditional expectations:

1. $\mathbb{E}[c_1g(x_1) + c_2h(x_2) | Y = y] = c_1\mathbb{E}[g(X_1) | Y = y] + c_2\mathbb{E}[h(X_2) | Y = y]$
2. If $g \geq 0$, then $\mathbb{E}[g(x) | Y = y] \geq 0$.
3. $\mathbb{E}[f(X, Y) | Y = y] = \mathbb{E}[f(X, y) | Y = y]$.
4. If $X \perp\!\!\!\perp Y$, $\mathbb{E}[g(X) | Y = y] = \mathbb{E}[g(X)]$
5. $\mathbb{E}[g(x)h(y) | Y = y] = h(y)\mathbb{E}[g(x) | Y = y]$
6. $\mathbb{E}[g(x)h(y)] = \sum_y h(y)\mathbb{E}[g(x) | Y = y]p_Y(y) = \mathbb{E}[h(Y)\mathbb{E}[g(X) | Y]]$

Proof of 3.

$$\begin{aligned}\mathbb{E}[f(X, Y) \mid Y = y] &= \sum_{x, z} f(x, z) p_{X, Y \mid Y}(x, z \mid y) \\ &= \sum_{x, z} f(x, z) \frac{p_{X, Y, Y}(x, z, y)}{p_Y(y)} \\ &= \sum_x f(x, y) \frac{p_{X, Y}(x, y)}{p_Y(y)} \\ &= \mathbb{E}[f(X, y) \mid Y = y].\end{aligned}$$

□

Remark. $\mathbb{E}[f(X, y)] \neq \mathbb{E}[f(X, y) \mid Y = y]$.

2 Random Sums

Definition 2.0.1. Let $(\xi_i)_{i=1}^{\infty}$ be i.i.d random variables, N be a \mathbb{N}_0 -valued random variable, $N \perp\!\!\!\perp (\xi_i)_{i=1}^{\infty}$. The *random sum* is defined as

$$X = \sum_{i=1}^N \xi_i = \sum_{n=0}^{\infty} \left(\sum_{i=1}^n \xi_i \right) \mathbf{1}_{N=n} = \begin{cases} \sum_{i=1}^n \xi_i & \text{if } N = n \geq 1 \\ 0 & \text{if } N = 0. \end{cases}$$

Question. What is the distribution of X ?

Let X, N be random variables. N is \mathbb{N}_0 -valued. The condition CDF is

$$F_{X|N}(x|n) = \mathbb{P}(X \leq x \mid N = n)$$

if $\mathbb{P}(N = n) \neq 0$. This is an actual CDF, but for the random variable $X \mid N = n$.

Suppose that X is continuous and $F_{X|N}(x|n)$ is a differentiable function of x for each n such that $p_N(n) > 0$. The conditional PDF is

$$f_{X|N}(x|n) = \frac{d}{dx} F_{X|N}(x|n).$$

$$\int_a^b f_{X|N}(x|n) dx = F_{X|N}(b|n) - F_{X|N}(a|n).$$