

Improving Precision in Estimating Marginal Treatment Effects in Randomized Trials

Michael Rosenblum, Joshua Betz, Kelly Van Lancker

Johns Hopkins Bloomberg School of Public Health

<https://mrosenblumbiostat.wordpress.com>

September 3, 2023

Disclaimer

The opinions in this presentation are of the authors (Michael Rosenblum, Joshua Betz, Kelly Van Lancker) and do not necessarily represent Johns Hopkins University, the FDA/HHS, or anyone else.

Overview

Covariate adjustment in randomized trials for estimating marginal (average) treatment effects:

- Preplanned adjustment for baseline variables when estimating average treatment effect in primary efficacy analysis.
- Target of inference (estimand) is same as when using unadjusted estimator.
- Can improve precision and reduce required sample size to achieve desired power.
- Goal: avoid making any model assumptions beyond what's assumed for unadjusted estimator (robustness to model misspecification).

Use of Covariate Adjustment in Randomized Trials: Two Surveys

Pocock et al. (2002) surveyed 50 randomized clinical trial reports. Findings: “The statistical emphasis on covariate adjustment is quite complex and often poorly understood, and there remains confusion as to what is an appropriate statistical strategy.”

Austin et al. (2010) surveyed 114 randomized trial articles. Findings: only 39 presented an adjusted analysis. Paper title: “A substantial and confusing variation exists in handling of baseline covariates in randomized controlled trials: a review of trials published in leading medical journals.”

FDA Guidance Documents on Covariate Adjustment

- ① ICH E9 Statistical Principles for Clinical Trials (1998):
“Pretrial deliberations should identify those covariates and factors expected to have an important influence on the primary variable(s), and should consider how to account for these in the analysis to improve precision...”
- ② FDA guidance (2023): Adjusting for Covariates in Randomized Clinical Trials for Drugs and Biological Products
“Covariate adjustment leads to efficiency gains when the covariates are prognostic for the outcome of interest in the trial. Therefore, FDA recommends that sponsors adjust for covariates that are anticipated to be most strongly associated with the outcome of interest.”

Goal of Covariate Adjustment for Marginal Treatment Effects

- Target of Inference (Estimand) is Population Average Treatment Effect, i.e., a contrast between (marginal) outcome distributions if all were assigned to treatment versus all assigned to control. (Intention To Treat)
- **Goal: Estimation of Average Treatment Effect in a Randomized Trial.**
- **If baseline variables prognostic for outcome, Covariate adjustment has potential to substantially improve precision (shorter CI's), reduce sample size, and reduce trial duration compared to unadjusted estimator.**
- Require consistent, interpretable, model-robust estimators.
- Intuition: Gain precision by adjusting for chance imbalances in prognostic baseline variables between study arms.

Some related work on covariate adjustment

Yang and Tsiatis, 2001, Zhang et al. 2008; Tsiatis et al. 2008, Rubin and van der Laan, 2008, Moore and van der Laan, 2009, Zhang and Gilbert 2010, Moore et al. 2011, Tian et al. 2012, Zheng et al. 2015, Vermuelen et al. 2015, Wager et al. 2016, Wu and Gagnon-Bartsch 2018, Zhang and Ma, 2019, Jiang et al. 2019, Wang et al. 2019.

Some material presented below is from:

Benkeser, Diaz, Luedtke, Segal, Scharfstein, Rosenblum (2020)
Improving Precision and Power in Randomized Trials for COVID-19
Treatments Using Covariate Adjustment, for Binary, Ordinal, or
Time to Event Outcomes. Biometrics.
<https://doi.org/10.1111/biom.13377>

Estimands (Targets of Inference)

Estimands (contrasts between marginal distributions under treatment and control):

- For continuous outcomes: difference in means.
- For binary outcomes: risk difference, relative risk, odds ratio.
- For ordinal outcomes: difference in means, the Mann-Whitney estimand= $P(\text{random individual assigned to treatment has better outcome than random individual assigned to control})$, average of cumulative log odds ratios over outcome levels.
- For time-to-event outcomes: difference in survival probabilities, ratio of survival probabilities, difference in restricted mean survival times.

Estimators:

- For each estimand, there exist corresponding covariate adjusted estimators that leverage information in baseline variables to improve precision and reduce required sample size to achieve desired power.

Continuous Outcomes: Example of Planning Alzheimer's Disease Trial

Context: Vitamin E and Donepezil for the Treatment of Mild Cognitive Impairment (MCI) phase 3 randomized trial (Petersen et al. 2005)

- Primary Outcome Y : Change in Clinical Dementia Rating Sum of Boxes (CDR-SB) at 1.5 years vs. baseline.
- Study arms A : new drug vs. placebo.
- Baseline variables B : age, gender, Alzheimer's Disease Assessment Scale (ADAS)-cognitive score, Mini-Mental State Examination (MMSE) score, Activities of Daily Living total score, Global Deterioration scale, and CDR-SB. .

Goal: Estimate Avg. Treatment Effect (Estimand)

$$E(Y|A = 1) - E(Y|A = 0).$$

- 23% sample size reduction comparing covariate-adjusted estimator to unadjusted.

Goals of Covariate Adjustment for Marginal Treatment Effect

- **Goal: Estimation of Average Treatment Effect in a Randomized Trial.** Require consistent, interpretable, model-robust estimators.
- **Not Goal:** Estimation of Conditional (within stratum of **baseline variables** B) Treatment Effects, e.g., $E(Y|A = 1, B) - E(Y|A = 0, B)$.
Conditional Treatment Effects would be useful to know, but typically require model assumptions (such as logistic regression model) and uninterpretable under model misspecification.
- **Not Goal:** Finding subpopulations who benefit or patient-specific optimal treatment policy. (Also useful goal in its own right.)

Assumptions

We assume:

- Treatment Randomized (A independent of B) by design.
- Participant data vectors (B_i, A_i, Y_i) , for $i = 1$ to n , independent, identically distributed draws from unknown distribution P .


These assumptions (or similar) are needed for standard, unadjusted estimator to be consistent (converge to average treatment effect).

No assumptions on the relationship among B, A, Y except randomization (A, B independent).

We Do Not Make Any Parametric Model Assumptions

- Population distribution of Y given A, B may differ arbitrarily from, e.g., linear regression model
$$E(Y|A, B) = \beta_0 + \beta_1 A + \beta_2 B.$$
- True relationships among B, A, Y may be much more complex than this.
- We require consistent estimators under arbitrary model misspecification.

Hypothetical Example of Misspecification:



Example: ANCOVA estimator for Mean Difference

For continuous Y , to estimate $\mu_1 - \mu_0$ where $\mu_a = E(Y|A = a)$:

- Fit linear regression model $E(Y = 1|A, B) = \beta_0 + \beta_1 A + \beta_2 B$.
- Compute standardized estimators for μ_0, μ_1 :
 - $\hat{\mu}_0 = \frac{1}{n} \sum_{i=1}^n \hat{\beta}_0 + \hat{\beta}_2 B_i$ (i.e., set $A = 0$ in model fit)
 - $\hat{\mu}_1 = \frac{1}{n} \sum_{i=1}^n \hat{\beta}_0 + \hat{\beta}_1 + \hat{\beta}_2 B_i$ (i.e., set $A = 1$ in model fit)
- ANCOVA estimator of $\mu_1 - \mu_0$ is mean difference $\hat{\mu}_1 - \hat{\mu}_0$.

Some remarkable properties of ANCOVA estimator (Yang and Tsiatis, 2001):

- Consistent (converges to average treatment effect) **under arbitrary model misspecification**.
- Equal or better precision (asymptotically) than unadjusted estimator (difference between sample means).

Note: ANCOVA estimator simplifies to $\hat{\beta}_1$, but above construction applies more generally (as shown below).

ANCOVA vs. Unadjusted Estimator Example

TADS Clinical Trial

- Treatment for Adolescents with Depression Study.
- Primary outcome Y : change of Childrens Depression Rating Scale-Revised (CDRS-R) score over 12 weeks.
- Treatment assignment A : Fluoxetine (107) and placebo (112).
- Baseline variables B : baseline CDRS-R score and 8 others on next slide.

ANCOVA Estimator: Impact of each Baseline Variable in TADS trial

Regress outcome (Y =change score) on arm A , baseline vars B :

$$E(Y|A, B) = \beta_0 + \beta_A A + \beta_1 B_1 + \cdots + \beta_9 B_9.$$

All baseline variables B_1, \dots, B_9 standardized (centered and divided by standard error).

Estimated Coefficients from Regression Model:

ARM	age	sex	CDRS-R	CGI	CGAS	RADS	Suicide ideation	Depr. episode	Comor.
$\hat{\beta}_A$	$\hat{\beta}_1$	$\hat{\beta}_2$	$\hat{\beta}_3$	$\hat{\beta}_4$	$\hat{\beta}_5$	$\hat{\beta}_6$	$\hat{\beta}_7$	$\hat{\beta}_8$	$\hat{\beta}_9$
-4.4	1.5	-0.1	-8.0	-0.6	0.1	-1.0	-1.6	0.8	-0.4

Estimator of average treatment effect is $\hat{\beta}_A = -4.4$.

Clinical Trial Applications

Trial Name	Unadjusted Estimator (95% CI)	ANCOVA Estimator (95% CI)	Impact of Adjustment	Variance Reduction (\hat{R}^2)
TADS	-1.4 (-6.0, 3.1)	-4.4 (-8.1, -0.7)	-2.9	32%

Table: Summary of clinical trial data analyses. Negative estimates are in direction of treatment benefit. Variance reduction refers to reduction due to adjusting for baseline variables by ANCOVA.

ANCOVA Estimator: Impact of each Baseline Variable in TADS trial

Regress outcome (Y =change score) on arm A , baseline vars B :

$$E(Y|A, B) = \beta_0 + \beta_A A + \beta_1 B_1 + \cdots + \beta_9 B_9.$$

All baseline variables B_1, \dots, B_9 standardized (centered and divided by standard error).

Estimated Coefficients from Regression Model:

ARM	age	sex	CDRS-R	CGI	CGAS	RADS	Suicide ideation	Depr. episode	Comor.
$\hat{\beta}_A$	$\hat{\beta}_1$	$\hat{\beta}_2$	$\hat{\beta}_3$	$\hat{\beta}_4$	$\hat{\beta}_5$	$\hat{\beta}_6$	$\hat{\beta}_7$	$\hat{\beta}_8$	$\hat{\beta}_9$
-4.4	1.5	-0.1	-8.0	-0.6	0.1	-1.0	-1.6	0.8	-0.4
Imbal. l_j	0.1	0.0	-0.2	-0.3	-0.1	-0.3	-0.2	0.0	-0.2
Correction i.e., $\hat{\beta}_j l_j$	0.1	0	1.9	0.2	0	0.3	0.4	0	0.1

Estimator of average treatment effect is $\hat{\beta}_A = -4.4$.

Binary Outcomes: Example of Planning MISTIE Phase III Stroke Trial

Problem: Confirmatory trial of new surgical treatment for intracerebral hemorrhage (PI: Daniel Hanley).

- Primary Outcome Y : modified Rankin Scale ≤ 3 at 180 days from enrollment.
- Study arms A : surgery vs. standard of care.
- Baseline variables B : NIH Stroke Scale, clot volume, and location.

Goal: Estimate Avg. Treatment Effect

$$P(Y = 1|A = 1) - P(Y = 1|A = 0).$$

Simulated trials based on resampling participants from MISTIE Phase II data.

- 38% precision gain from adjusted estimator compared to unadjusted.
- Equivalent to 28% $(1 - \frac{1}{1.38})$ reduction in required sample size.

Binary Outcomes: Estimands

Notation: outcome Y ; study arm $A = 0$ (control) and $A = 1$ (treatment).

$\mu_1 = P(Y = 1 | A = 1)$ and $\mu_0 = P(Y = 1 | A = 0)$.

Estimand (target of inference) is contrast between μ_1, μ_0 :

Estimand 1: Risk Difference. Difference between probability of good outcome comparing treatment to control arms, that is, $P(Y = 1 | A = 1) - P(Y = 1 | A = 0) = \mu_1 - \mu_0$.

Estimand 2: Relative Risk. Ratio of probability of good outcome comparing treatment to control arms, that is, $P(Y = 1 | A = 1) / P(Y = 1 | A = 0) = \mu_1 / \mu_0$.

Estimand 3: Odds Ratio. Ratio of odds of good outcome, comparing treatment to control arms, that is, $\{\mu_1 / (1 - \mu_1)\} / \{\mu_0 / (1 - \mu_0)\}$.

Binary Outcomes: Covariate Adjusted Estimators

For dichotomous Y :

- Fit logistic regression model for $P(Y = 1|A, B) = \text{logit}^{-1}(\beta_0 + \beta_1 A + \beta_2 B)$.
- Compute standardized estimators for treatment specific means μ_0, μ_1 :
 - $\hat{\mu}_0 = \frac{1}{n} \sum_{i=1}^n \text{logit}^{-1}(\hat{\beta}_0 + \hat{\beta}_2 B_i)$
 - $\hat{\mu}_1 = \frac{1}{n} \sum_{i=1}^n \text{logit}^{-1}(\hat{\beta}_0 + \hat{\beta}_1 + \hat{\beta}_2 B_i)$
- Estimator is contrast of interest between μ_1, μ_0 , e.g., risk difference $\hat{\mu}_1 - \hat{\mu}_0$.

Estimator $\hat{\mu}_1 - \hat{\mu}_0$ consistent **under arbitrary model**

misspecification (Scharfstein, Rotnitzky, Robins, 1999; Moore and van der Laan, 2009, Ge et al. 2011).

Same holds for other (unconditional) estimands, e.g., relative risk

Ordinal Outcome: Estimands

Outcome Y ; study arm $A = 0$ (control) and $A = 1$ (treatment).

Estimand 1: Difference in means (DIM). For $u(\cdot)$ a pre-specified, real-valued transformation of an outcome, the estimand is defined as:

$$\text{DIM: } E\{u(Y) \mid A = 1\} - E\{u(Y) \mid A = 0\}.$$

Estimand 2: Mann-Whitney (MW) estimand.

$$\text{MW: } P\left(\tilde{Y} > Y \mid \tilde{A} = 1, A = 0\right) + \frac{1}{2}P\left(\tilde{Y} = Y \mid \tilde{A} = 1, A = 0\right),$$

for (A, Y) and (\tilde{A}, \tilde{Y}) independent treatment-outcome pairs.

Estimand 3: Log-odds ratio (LOR).

$$\text{LOR: } \frac{1}{K-1} \sum_{j=1}^{K-1} \log \left\{ \frac{\text{odds}(Y \leq j \mid A = 1)}{\text{odds}(Y \leq j \mid A = 0)} \right\}.$$

Ordinal Outcome: Covariate Adjusted Estimator

Notation: baseline variables \mathbf{B} ; study arm A ; outcome Y .

- First estimate CDF of Y given study arm $A = a$ for each $a \in \{0, 1\}$, and then substitute into desired estimand.
- To estimate CDF for study arm a , **using only data with** $A = a$, we fit proportional odds working model

$$\text{logit} \{P(Y \leq j | A = a, \mathbf{B})\} = \alpha_j + \beta^\top \mathbf{B},$$

for each $j = 1, \dots, K - 1$ with parameters $\alpha_1, \dots, \alpha_{K-1}, \beta$.

- Then estimator of CDF is $\frac{1}{n} \sum_{i=1}^n \hat{P}(Y \leq j | A = a, \mathbf{B}_i)$ where sum is over baseline variables \mathbf{B}_i for **all participants** $i = 1, \dots, n$ pooled across study arms.
- Estimator consistent and asymptotically normal under arbitrary model misspecification. Also can be generalized to handle outcomes missing at random, and then double robust. We use superpopulation inference framework assuming $(\mathbf{B}_i, A_i, Y_i), i = 1, \dots, n$ i.i.d. draws from unknown distribution.

Time-to-Event Outcome: Estimands

Notation: study arm $A = 0$ (control) and $A = 1$ (treatment); T is event time.

Estimand 1: Difference in restricted mean survival times (RMSTs). The RMST is the expected value of a survival time that is truncated at a specified time τ , that is,

$$\text{RMST: } E(\min\{T, \tau\} | A = 1) - E(\min\{T, \tau\} | A = 0).$$

Estimand 2: Survival probability difference (also called risk difference, RD). Difference between arm-specific probabilities of survival to a specified time t^* , that is,

$$\text{RD: } P(T \leq t^* | A = 1) - P(T \leq t^* | A = 0).$$

Estimand 3: Relative risk (RR). Ratio of the arm-specific probabilities of survival to a specified time t^* , that is,

$$\text{RR: } \frac{P(T \leq t^* | A = 1)}{P(T \leq t^* | A = 0)}.$$

Time-to-Event Outcome: Covariate Adjusted Estimators

- Adjusted Kaplan-Meier estimator for survival curves: Moore and van der Laan 2009, Zhang 2014
- Restricted Mean Survival Time: Moore and van der Laan 2009, Diaz et al. 2019
- If assume proportional hazards model (marginally): Lu and Tsiatis et al. 2011.

Time-to-Event Example: Population, Baseline Variables, and Outcomes in COVID-19 context

- Population: hospitalized, COVID-19 positive patients
- Outcomes: intensive care unit (ICU) admission, intubation with ventilation, and death.
- Baseline variables: age, sex, required supplemental oxygen at ED presentation, dyspnea, hypertension, bilateral infiltrates on the chest x-ray

Data Generating Distributions for Simulations (Survival Outcomes)

- Patient data re-sampled with replacement from 500 patients hospitalized at Weill Cornell Medicine New York Presbyterian Hospital prior to March 28, 2020.
- Simulated sample sizes $n = 100, 200, 500$, and 1000.
- Hypothetical treatment variable drawn independent of all other data
- To simulate positive treatment effect: add independent draw from a χ^2 with 4 d.f. to each treatment arm participant's outcome
- Censoring: 5% censored completely at random; censoring time from uniform distribution on $\{1, \dots, 14\}$.

Results: difference in restricted mean survival times (RMST) 14 days after hospitalization

Table: Results when treatment effect is 1 day. n =sample size;
RE=relative efficiency (ratio of adjusted vs. unadj. MSE).

n	Estimator	Power	$n \times \text{MSE}$	RE
100	Unadjusted	0.09	53.7	1.00
100	Adjusted	0.15	51.0	0.95
200	Unadjusted	0.33	62.7	1.00
200	Adjusted	0.40	56.4	0.90
500	Unadjusted	0.74	72.9	1.00
500	Adjusted	0.82	62.2	0.85
1000	Unadjusted	0.96	76.5	1.00
1000	Adjusted	0.98	63.5	0.83

R Packages

- Ordinal Outcomes: R package, drord,
<https://github.com/benkeser/drord>.
- Time-to-Event Outcomes: R package survtmle
<https://github.com/idiast/survtmle>
- Continuous, Binary, Time-to-Event: R package speff2trial
<https://cran.r-project.org/web/packages/speff2trial/speff2trial.pdf>

Stratified Randomization and Covariate Adjustment

Wang, B., Susukida, R., Mojtabai, R., Masoumeh, A.-E.; and Rosenblum, M. (2019) Model-Robust Inference for Clinical Trials that Improve Precision by Stratified Randomization and Adjustment for Additional Baseline Variables.

<https://arxiv.org/abs/1910.13954>

- Two commonly used methods for improving precision: **stratified randomization and covariate adjustment.**
- Many trials do not fully capitalize on the combined precision gains from these two methods.
- 70% of confirmatory trials use stratified randomization (Lin et al. 2015).
- Gives general formula for variance of estimator (unadjusted and adjusted) under stratified randomization.
- Handles many estimators used to analyze randomized trials: ANCOVA, standardized logistic regression, MMRM, Kaplan-Meier estimator.

Recommendations for Primary Efficacy Analysis

- ① **Estimand when the outcome is ordinal.** Recommend: difference between means or the Mann-Whitney estimand. Don't recommend log odds ratios.
- ② **Covariate adjustment.** Adjust for prognostic baseline variables to improve precision and power.
 - ① Baseline variables should be specified before the trial is started (or selected using prespecified algorithm, e.g., with cross-validation).
- ③ **Confidence intervals (CI) and hypothesis testing.** Nonparametric bootstrap (BCa), 10000 replicates for CI.
 - ① Entire estimation procedure repeated in each replicate data set.
 - ② Hypothesis tests: invert confidence interval or use permutation methods (latter especially for smaller sample sizes)

Recommendations for Primary Efficacy Analysis (con't)

1 Use Information Monitoring

- 1 Final analysis time based on the information accrued (1/estimator variance).
- 2 Precision gains from covariate adjustment translate into faster information accrual and shorter trial duration.

2 Plotting the CDF and the probability mass function (PMF) when the outcome is ordinal.

- 1 Covariate adjusted estimate of the PMF and/or CDF of primary outcome plotted for each study arm.
- 2 Pointwise and simultaneous confidence intervals displayed

3 Missing covariates. Impute based only on data from those covariates that were observed.

4 Missing outcomes. Use doubly robust methods and sensitivity analyses of robustness to assumptions.

Bottom Line: Pros/Cons in Using Covariate Adjustment

- **Pro:** Covariate adjustment as described above gives consistent estimator of average treatment effect (same quantity estimated by unadjusted estimator); does not require parametric model assumptions.
- **Pro:** If baseline variable(s) strongly prognostic for outcome, covariate adjustment can substantially improve precision + power (or reduce sample size) vs. unadjusted estimator.
- **Pro:** Covariate adjustment useful even in large trials; that's where biggest equivalent sample size reduction
- **Con:** Can lose efficiency (at small sample size) if all baseline variables pure noise, but losses small.
In simulations, 2% loss at sample size 100; < 1% loss at sample size 1000 (Colantuoni and Rosenblum 2015).

Some resources for optimizing trial design and analysis

Available at **<http://rosenblum.jhu.edu>**:

- Papers and Open-Source Software for Covariate-Adjustment
- Adaptive Enrichment Design Trial Planning/Optimization Software
- FDA short-course video-recording
- Plain Language Document Outlining Advantages and Limitations of Adaptive Enrichment Designs
- Papers with Statistical Methods and Case Studies

- Austin, P. C., A. Manca, M. Zwarenstein, D. N. Juurlink, and M. B. Stanbrook (2010). A substantial and confusing variation exists in handling of baseline covariates in randomized controlled trials: a review of trials published in leading medical journals. *Journal of Clinical Epidemiology* 63(2), 142–153.
- Benkeser, D., M. Carone, and P. B. Gilbert (2018). Improved estimation of the cumulative incidence of rare outcomes. *Statistics in Medicine* 37(2), 280–293.
- Benkeser, D., I. Diaz, A. Luedtke, J. Segal, D. Scharfstein, and M. Rosenblum (2020). Improving Precision and Power in Randomized Trials for COVID-19 Treatments Using Covariate Adjustment, for Ordinal or Time to Event Outcomes Biometrics. <https://doi.org/10.1111/biom.13377> .
- Benkeser, D., P. B. Gilbert, and M. Carone (2019). Estimating and testing vaccine sieve effects using machine learning. *Journal of the American Statistical Association* 114(527), 1038–1049.
- Brooks, J. C., M. J. van der Laan, D. E. Singer, and A. S. Go (2013). Targeted minimum loss-based estimation of causal

effects in right-censored survival data with time-dependent covariates: Warfarin, stroke, and death in atrial fibrillation. *Journal of Causal Inference* 1(2), 235–254.

CDC COVID-19 Response Team (2020). Severe Outcomes Among Patients with Coronavirus Disease 2019 (COVID-19) United States, February 12 to March 16, 2020.
<https://www.cdc.gov/mmwr/volumes/69/wr/mm6912e2.htm> .
MMWR Morb Mortal Wkly Rep 69(12), 343–346.

Chaisinankul, N., O. Adeoye, R. J. Lewis, J. C. Grotta, J. Broderick, T. G. Jovin, R. G. Nogueira, J. J. Elm, T. Graves, S. Berry, K. R. Lees, A. D. Barreto, J. L. Saver, null null, A. Furlan, B. Baxter, H. L. Lutsep, M. Ribo, O. Jansen, R. Gupta, and V. M. Pereira (2015). Adopting a patient-centered approach to primary outcome analysis of acute stroke trials using a utility-weighted modified rankin scale. *Stroke* 46(8), 2238–2243.

Chen, P.-Y. and A. A. Tsiatis (2001). Causal inference on the

- difference of the restricted mean lifetime between two groups. *Biometrics* 57(4), 1030–1038.
- Díaz, I., E. Colantuoni, D. F. Hanley, and M. Rosenblum (2019). Improved precision in the analysis of randomized trials with survival outcomes, without assuming proportional hazards. *Lifetime Data Analysis* 25(3), 439–468.
- Díaz, I., E. Colantuoni, and M. Rosenblum (2016). Enhanced precision in the analysis of randomized trials with ordinal outcomes. *Biometrics* 72(2), 422–431.
- Efron, B. and R. J. Tibshirani (1994). *An introduction to the bootstrap*. CRC press.
- EMA (2015). Guideline on adjustment for baseline covariates in clinical trials. Reference number EMA/CHMP/295050/2013. Committee for Medicinal Products for Human Use (CHMP).
- FDA (2019). Adjusting for Covariates in Randomized Clinical Trials for Drugs and Biologics with Continuous Outcomes. Draft Guidance for Industry.
<https://www.fda.gov/media/123801/download>.

- FDA (2020, May). COVID-19: Developing Drugs and Biological Products for Treatment or Prevention. Guidance for Industry. <https://www.fda.gov/media/137926/download>.
- FDA and EMA (1998). E9 statistical principles for clinical trials. *U.S. Food and Drug Administration: CDER/CBER. European Medicines Agency: CPMP/ICH/363/96*.
- Ge, M., L. K. Durham, R. D. Meyer, W. Xie, and N. Thomas (2011). Covariate-adjusted difference in proportions from clinical trials using logistic regression and weighted risk differences. *Drug Information Journal* 45(4), 481–493.
- Jiang, F., L. Tian, H. Fu, T. Hasegawa, and L. J. Wei (2019). Robust alternatives to ancova for estimating the treatment effect via a randomized comparative study. *Journal of the American Statistical Association* 114(528), 1854–1864.
- Kaplan, E. L. and P. Meier (1958). Nonparametric estimation from incomplete observations. *Journal of the American Statistical Association* 53(282), 457–481.

- Lu, X. and A. A. Tsiatis (2011). Semiparametric estimation of treatment effect with time-lagged response in the presence of informative censoring. *Lifetime Data Analysis* 17(4), 566–593.
- McCullagh, P. (1980). Regression models for ordinal data. *Journal of the Royal Statistical Society: Series B (Methodological)* 42(2), 109–127.
- Moore, K. L., R. Neugebauer, T. Valappil, and M. J. van der Laan (2011). Robust extraction of covariate information to improve estimation efficiency in randomized trials. *Statistics in Medicine* 30(19), 2389–2408.
- Moore, K. L. and M. J. van der Laan (2009). Covariate adjustment in randomized trials with binary outcomes: Targeted maximum likelihood estimation. *Statistics in Medicine* 28(1), 39.
- Moore, K. L. and M. J. van der Laan (2009). Increasing power in randomized trials with right censored outcomes through covariate adjustment. *Journal of Biopharmaceutical Statistics* 19(6), 1099–1131. PMID: 20183467.

- Nogueira, R. G., A. P. Jadhav, D. C. Haussen, A. Bonafe, R. F. Budzik, P. Bhuva, D. R. Yavagal, M. Ribo, C. Cognard, R. A. Hanel, C. A. Sila, A. E. Hassan, M. Millan, E. I. Levy, P. Mitchell, M. Chen, J. D. English, Q. A. Shah, F. L. Silver, V. M. Pereira, B. P. Mehta, B. W. Baxter, M. G. Abraham, P. Cardona, E. Veznedaroglu, F. R. Hellinger, L. Feng, J. F. Kirmani, D. K. Lopes, B. T. Jankowitz, M. R. Frankel, V. Costalat, N. A. Vora, A. J. Yoo, A. M. Malik, A. J. Furlan, M. Rubiera, A. Aghaebrahim, J.-M. Olivot, W. G. Tekle, R. Shields, T. Graves, R. J. Lewis, W. S. Smith, D. S. Liebeskind, J. L. Saver, and T. G. Jovin (2018). Thrombectomy 6 to 24 hours after stroke with a mismatch between deficit and infarct. *New England Journal of Medicine* 378(1), 11–21. PMID: 29129157.
- Parast, L., L. Tian, and T. Cai (2014). Landmark estimation of survival and treatment effect in a randomized clinical trial. *Journal of the American Statistical Association* 109(505), 384–394.

- Royston, P. and M. K. Parmar (2011). The use of restricted mean survival time to estimate the treatment effect in randomized clinical trials when the proportional hazards assumption is in doubt. *Statistics in Medicine* 30(19), 2409–2421.
- Scharfstein, D. O., A. Rotnitzky, and J. M. Robins (1999). Adjusting for nonignorable drop-out using semiparametric nonresponse models. *Journal of the American Statistical Association* 94(448), 1096–1120.
- Stitelman, O. M., V. De Gruttola, and M. J. van der Laan (2011). A general implementation of tmle for longitudinal data applied to causal inference in survival analysis. *The International Journal of Biostatistics* 8(1).
- Tsiatis, A. A., M. Davidian, M. Zhang, and X. Lu (2008). Covariate adjustment for two-sample treatment comparisons in randomized clinical trials: A principled yet flexible approach. *Statistics in Medicine* 27(23), 4658–4677.
- van der Vaart, A. W. (1998). *Asymptotic Statistics*. Cambridge: Cambridge University Press.

- Vermeulen, K., O. Thas, and S. Vansteelandt (2015). Increasing the power of the mann-whitney test in randomized experiments through flexible covariate adjustment. *Statistics in Medicine* 34(6), 1012–1030.
- Yang, L. and A. A. Tsiatis (2001). Efficiency study of estimators for a treatment effect in a pretest–posttest trial. *The American Statistician* 55(4), 314–321.
- Zhang, M. (2014). Robust methods to improve efficiency and reduce bias in estimating survival curves in randomized clinical trials. *Lifetime Data Analysis* 21(1), 119–137.
- Zhang, M. and P. B. Gilbert (2010). Increasing the efficiency of prevention trials by incorporating baseline covariates. *Statistical Communications in Infectious Diseases* 2(1).
- Zhang, M., A. A. Tsiatis, and M. Davidian (2008). Improving efficiency of inferences in randomized clinical trials using auxiliary covariates. *Biometrics* 64(3), 707–715.

- Austin, Peter C., et al. A substantial and confusing variation exists in handling of baseline covariates in randomized controlled trials: a review of trials published in leading medical journals. *J. Clin Epi* 63.2 (2010): 142-153.
- Colantuoni, E. and Rosenblum, M. (2015) Leveraging Prognostic Baseline Variables to Gain Precision in Randomized Trials. *Statistics in Medicine*. 34(18), 2602-2617. <http://goo.gl/evGHF6>
- Diaz, I., Colantuoni, E., Rosenblum, M. Enhanced Precision in the Analysis of Randomized Trials with Ordinal Outcomes. *Biometrics*. (2016)
- Jarskog, L. F., Hamer, R. M., Catellier, D. J., Stewart, D. D., LaVange, L., Ray, N., Golden, L. H., Lieberman, J. A. and Stroup, T. S. (2013). Metformin for weightloss and metabolic control in overweight outpatients with schizophrenia and schizoaffective disorder. *American Journal of Psychiatry* 170(9), 1032-1040.

- Jiang, F., L. Tian, H. Fu, T. Hasegawa, M. A. Pfeffer, and L. J. Wei (2016). Robust alternatives to ANCOVA for estimating the treatment effect via a randomized comparative study, Harvard University Biostatistics Working Paper. Series Working paper 209.
- Moore K, van der Laan MJ. Covariate adjustment in randomized trials with binary outcomes: targeted maximum likelihood estimation. *Statistics in Medicine* 2009; 28(1):39-64.
- Moore, K. L., R. Neugebauer, T. Valappil, and M. J. van der Laan (2011). Robust extraction of covariate information to improve estimation efficiency in randomized trials. *Statistics in Medicine* 30 (19), 2389-2408.
- Petersen, R. C., Thomas, R. G., Grundman, M., Bennett, D., Doody, R., Ferris, S., Galasko, D., Jin, S., Kaye, J., Levey, A., Pfeiffer, E., Sano, M., van Dyck, C. H. and others. (2005). Vitamin E and Donepezil for the Treatment of Mild Cognitive Impairment. *New England Journal of Medicine* 352(23),

- Pocock, S. J., Assmann, S. E., Enos, L. E., and Kasten, L. E. (2002). Subgroup analysis, covariate adjustment and baseline comparisons in clinical trial reporting: current practice and problems. *Statistics in medicine*, 21(19), 2917-2930.
- Robins JM, Sued M, Lei-Gomez Q, Rotnitzky A. Double-robustness with improved efficiency in missing and causal inference models. Technical Report, Harvard School of Public Health, 2007
- Rotnitzky A, Lei Q, Sued M, Robins JM. Improved double-robust estimation in missing data and causal inference models. *Biometrika* 2012; 99(2):439-456.
- Rubin D, van der Laan MJ. Empirical efficiency maximization: improved locally efficient covariate adjustment in randomized experiments and survival analysis. *International Journal of Biostatistics* 2008; 4(1):Article 5.

- Scharfstein, D.O., Rotnitzky, A., and Robins, J.M. Adjusting for nonignorable drop-out using semiparametric nonresponse models. *Journal of the American Statistical Association* 94, 1096–1120 (with Rejoinder, 1135–1146), 1999.
- Steingrimsdóttir, Jon Arni; Hanley, Daniel F.; and Rosenblum, Michael. Improving precision by adjusting for prognostic baseline variables in randomized trials with binary outcomes, without regression model assumptions, *Contemporary Clinical Trials*, 2017, <http://doi.org/10.1016/j.cct.2016.12.026>
- Treatment for Adolescents With Depression Study (TADS) Team. (2004). Fluoxetine, cognitive-behavioral therapy, and their combination for adolescents with depression: Treatment for adolescents with depression study (TADS) randomized controlled trial. *JAMA* 292(7), 807–820.
- Tan Z. Bounded, efficient and doubly robust estimating equations for marginal and nested structural models. *Biometrika* 2010; 97:661–682.

- Lu Tian, Tianxi Cai, Lihui Zhao, Lee-Jen Wei, On the covariate-adjusted estimation for an overall treatment difference with data from a randomized comparative clinical trial, Biostatistics, Volume 13, Issue 2, April 2012, Pages 256–273, <https://doi.org/10.1093/biostatistics/kxr050>
- Stefan Wager, Wenfei Du, Jonathan Taylor, Robert J. Tibshirani. High-dimensional randomized experiments. Proceedings of the National Academy of Sciences Nov 2016, 113 (45) 12673-12678; DOI: 10.1073/pnas.1614732113
- Wu, E. and J. A. Gagnon-Bartsch (2018). The LOOP Estimator: Adjusting for Covariates in Randomized Experiments. Evaluation Review 42(4), 458–488. <https://doi.org/10.1177/0193841X18808003>.
- ZHANG, M., TSIATIS, A. AND DAVIDIAN, M. (2008). Improving efficiency of inferences in randomized clinical trials using auxiliary covariates. Biometrics 64, 707–715.
- Zhang, Z, Ma, S. Machine learning methods for leveraging baseline covariate information to improve the efficiency of

- Wang, B., Susukida, R., Mojtabai, R., Masoumeh, A.-E.; and Rosenblum, M. (2019) Model-Robust Inference for Clinical Trials that Improve Precision by Stratified Randomization and Adjustment for Additional Baseline Variables.
<https://arxiv.org/abs/1910.13954>

Acknowledgments

MR was supported by the Johns Hopkins Center of Excellence in Regulatory Science and Innovation, which is funded by the Food and Drug Administration (FDA) of the U.S. Department of Health and Human Services (HHS) as part of a financial assistance award (U01FD005942). The contents are those of the author(s) and do not necessarily represent the official views of, nor an endorsement, by any of the aforementioned organizations, the FDA/HHS, nor the U.S. Government.