

# EpiCO19 Final Report

Epidemiological survey and active screening  
for COVID-19:

Investigating the circulation and impact of SARS-CoV-2 on  
the population of the health district of Cité Verte in  
Yaounde



Jan 04, 2020



**Supported by:**

ASCRES • CIRES •  
MINSANTE Cité Verte •  
CIRCB • GIZ

# Epidemiological survey and active screening for COVID-19: Investigating the circulation and impact of SARS-CoV-2 on the population of the health district of Cité Verte in Yaounde

Principal Investigators	Affiliation
Dr Laura Ciaffi	<b>ASCRES</b> - Association de soutien aux centres de recherches, d'enseignements et de soins Tél : 00237 694 92 67 86 Mail : <a href="mailto:lauraciaffi2002@yahoo.fr">lauraciaffi2002@yahoo.fr</a>
Dr Franck Wanda	<b>CIRES</b> – Centre International de Recherches, d'Enseignements et de Soins Tél : 00237 699 73 46 89 Mail : <a href="mailto:franck.wanda@cires.solutions">franck.wanda@cires.solutions</a>

Investigators	Affiliation
Dr Lucien Mama District medical Officer	<b>Ministry of Public health</b> District de Sante Cité Verte -Yaounde
Dr Joseph Fokam Virologist	<b>CIRCB</b> - Centre International de recherche Chantal Biya Yaoundé
Dr Eric Comte Public Health Specialist	<b>ASCRES</b> - Association de soutien aux centres de recherches, d'enseignements et de soins

Project Staff	Role
Eric Mimbe	Data Manager
Jeanine Meke	Interviewers Team Leader
Armel Tassegnig	Interviewers Team Leader
Thomas Eyinga	Interviewer
Evelyne Boyomo	Interviewer
Assana Gondio A Tiati	Interviewer
Yolande Maled	Interviewer
Josphine Embolo Manga	Interviewer
Michel Ntsama	Interviewer

Sponsor	
GIZ Cameroun	Mr Josselin Guillebert director ProPASSaR/GIZ Mme Patricia Ndong

## Table of Contents

Background .....	3
Methods .....	4
Design .....	4
Population .....	4
Sampling .....	4
Testing .....	5
Questionnaire .....	6
Field work .....	6
Data management .....	7
Data analysis .....	7
Ethical considerations .....	8
Results .....	8
Participation rate .....	8
Respondent characteristics .....	9
Seroprevalence estimates .....	9
Individual-level findings .....	<b>Error! Bookmark not defined.</b>
Conclusions .....	18
Bibliography .....	19

## Background

Coronavirus disease 2019 (COVID-19) is caused by the severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2), which was identified for the first time in China at the end of 2019<sup>1</sup>. The virus quickly spread globally, and the World Health Organization declared COVID-19 to be a pandemic on March 11, 2020. As at December 1, 2020, there have been over 67,000,000 notified cases, with over 1,500,000 reported deaths.<sup>2</sup>

Given the rapid spread of the epidemic in Europe and the Americas, and the handicapped response by countries with the richest health systems, the outlook for less developed countries, and sub-Saharan Africa in particular, seemed dire. High numbers of deaths were expected due to weaknesses in health systems, difficulties in enforcing hygiene measures, and perceived health vulnerabilities of the population.<sup>3,4</sup> But the trajectory of the epidemic on the continent appears to have gone against expectation. Despite having over 2,200,200 infections as of December 1,<sup>2</sup> Africa remains the least affected region and the mortality rate, even if not well documented, remains lower than expected.<sup>4</sup>

Multiple hypotheses have been advanced to explain the seemingly mild trajectory of the COVID-19 epidemic in Africa: researchers have pointed to the continent's warm climate, young population, or cross-reactive immunity from other infections as possible mitigating factors.<sup>4</sup> But an informed explanation of the epidemic trajectory requires accurate numbers on the actual extent of population infection. And, as is the case elsewhere,<sup>5</sup> the officially reported case counts in Africa significantly underestimate the extent of spread.<sup>4</sup>

In this context, the use of serological antibody tests to detect exposure to SARS-CoV-2 is valuable. A number of validated SARS-CoV-2 antibody tests now exist on the market,<sup>6</sup> and some of these are point-of-care lateral-flow immunoassays, which are affordable, easy to use and provide quick results. Although concerns about sensitivity and specificity remain, these antibody tests offer the opportunity to more accurately assess the prior infection rate of populations in regions where PCR-based testing has been uncommon.<sup>5</sup>

This report presents the protocol and results of our study using a lateral-flow immunoassay to assess the seroprevalence of anti-SARS-CoV-2 IgG and IgM antibodies in a region of Yaounde, the capital of Cameroon.

In addition to examining seroprevalence, our study also hopes to shed light on the broader impact of the epidemic on the socioeconomic wellbeing of the Cité Verte population. Soon after its first notified COVID-19 case (on March 6, 2020), Cameroon closed its borders and implemented significant social distancing measures, including mandatory bar closures after 6 PM, limitations on public transport occupancy, bans on gatherings of more than 50 people and the closing of in-person schools<sup>7</sup>. Similar measures were taken in other African countries.

The effects of these measures on the wellbeing of the Cameroonian population has not been well examined. Studies in other contexts have noted that during the pandemic period, access to routine health care services has been disrupted. We designed and implemented a survey to assess whether such disruptions were common in the studied population. This survey was performed concurrently with the seroprevalence assays.

## Methods

The EpiCO 19 study is a community-based survey to estimate the circulation and the impact of SARS-CoV-2 and the COVID-19 pandemic on a district in Yaounde, which is one of the cities most affected by COVID-19 in Cameroon.

### Design

EpiCO 19 is a cross-sectional clinico-epidemiological and socio-economic study.

### Population

The Cité Verte district, located in the Mfoundi department of the Center Region of Cameroon, covers a surface area of 19 square kilometers in the administrative area of Yaounde II. The total estimated population is 432,858, of which the majority are young. It is located in the oldest part of the city of Yaounde and residents are from several different regions of Cameroon.

The dwellings are of two types: 60% are permanent built with standard building materials while 40% are temporary structures. Some neighbourhoods, notably Tsinga and Cité Verte (the Cité Verte neighbourhood of the Cité Verte district), are better equipped in services and roads than others.

Cité Verte is divided into nine health zones with a total of 63 health structures, of which eight are public and three are run by private, faith-based providers. In each of the health areas there is a referent health structure and a number of community health workers supporting public health activities that are carried out by the Ministry of health or other partners in the community. These personnel have been key for the study to facilitate access to the population.

### Sampling

The following formula was used to calculate the size of the sample for the study:

$$n = Z^2 P(1 - P)/d^2$$

where n = sample size, Z = Z statistic for a level of confidence, P = expected prevalence or proportion (in proportion of one; if 20%, P = 0.2) and d = precision (in proportion of one; if 5%, d = 0.05).

With this formula, a level of confidence of 95% ( $Z=1.96$ ), a precision of 5%, and an estimated prevalence of 20% in the general population, a sample of 245 people was calculated to be sufficient, but the sample was increased to 1000 people to provide for improved precision and better representativeness of the different parts of the community.

Houses were randomly selected from a geographic grid, and geo-localisation software was used to guide the interviewers to the locations of selected households. A standard replacement procedure was used when the selected buildings were not residential.

Households, which were the sampling unit of the study, were defined as a group of people living in the same residence and sharing the kitchen.

In each household, all individuals between 5 and 80 years were included if they a), were living in the selected household, b) had been present in the household for at least 14 days and c) could give written informed consent (or had an adult guardian who could give consent). Children of less than 5 years and people with severe psychiatric illness were excluded.

Two hundred and fifty households were targeted for a total population of 1000 respondents living in the Cité Verte District.

## Testing

The EpiCO19 study used the Abbott Panbio™ COVID-19 IgG/IGM Rapid Test Device, an immunochromatographic test for the qualitative detection of IgG and IgM antibodies to SARS-CoV-2. The manufacturer-estimated sensitivity and specificity of the test are 95.8% and 94% respectively.

The serological tests were performed on capillary blood which was collected from a finger prick from all the consenting participants. All waste material was eliminated following safety procedures by the study team.

Additional steps were taken when active COVID-19 was suspected. (Such cases were identified if they matched the WHO COVID-19 criteria of symptoms, updated as at August 2020.) The national algorithm for further testing was applied to these suspected cases. First, a rapid antigen test was performed. If the result of the antigen test was negative, then a nasopharyngeal swab was taken. The swab was sent to the virological reference laboratory for an RNA PCR test for SARS-CoV-2. This PCR test was also proposed to all participants with a positive IgM result in the serological essay, since an IgM response can sometimes be observed concurrent with active infection.

The PCR tests were carried out at the virology laboratory of the CIRCB, the reference laboratory for diagnostic procedures in the Cité Verte District, following standard national procedures.

During sample collection all biosafety precautions were followed. The person in charge of swab collection was provided with full personal protection equipment (apron, gloves, face shield and FFP2 mask).

## Questionnaire

A questionnaire was created with KoboCollect©, a free software for epidemiological surveys.

The first few sections of the questionnaire focused on household composition, household income and recent deaths within the household. Then individual information about each survey respondent was surveyed. These sections included questions about the respondents'

- experience of symptoms compatible with COVID-19;
- health-seeking behavior during the pandemic;
- financing of healthcare during the pandemic;
- attitudes towards COVID-19; and
- respect of hygiene and physical distancing recommendations;

among others.

The questionnaire was field tested and modified following the feedback of the tests in the field.

## Field work

All the interviewers were trained over three days on research practice and ethics, and the specific study protocol.

In the first two weeks of study implementation, while waiting for the administrative authorization, a part of the team had time to conduct the geo-localisation of the majority of the households, supported by the health workers of each health area.

To increase receptiveness to the survey, awareness talks were carried out the day before, or a few hours before, the interviewers' visit. On the interview days, study teams visited each household to interview them immediately or to arrange an appointment for a future interview.

Once residents were gathered, the study was presented and a signature of informed consent was given out to be signed. For anyone below 21 years, a signature was requested of their parent or legal guardian; but consent was still solicited from anyone above 15 years. The survey questionnaire was then administered to each household member.

In a quiet and private space, the material for the tests was set up and all the procedures for the test were respected. A picture of the test was taken and a paper result was handed out to the participant.



When active COVID-19 was suspected, a nasopharyngeal swab was done, information about prevention measures was given and the surveillance system of the district was alerted.

## Data management

Once the questionnaire was completed, it was uploaded to a secure server. Data were regularly monitored for completeness and coherence, and for agreement between the responses of different members of each household. Where doubts remained, the relevant interviewer was contacted for clarification.

## Data analysis

For population-weighting, seroprevalence values were first calculated within each age-sex stratum. Weighted means were then obtained based on the stratum's share of the Yaounde population. The age-sex distribution for Yaounde was sourced from the 2018 cameroon DHS<sup>1</sup>.

Seroprevalence values were adjusted for test performance using the Rogan-Gladen formula<sup>2</sup>, which allows the estimation of true prevalence in the presence of a test with imperfect sensitivity and specificity. Test sensitivity and specificity were taken from Batra and others' validation study<sup>3</sup>. In that study, the IgG test correctly diagnosed 75 out of 82 serum samples of COVID-19 patients (91.5% sensitivity), and 149 out of 150 pre-pandemic samples (99.3% specificity).

We did not apply corrections for the performance of the IgM portion of the Abbott test; since IgM antibodies decline very rapidly, becoming undetectable a few weeks after infection<sup>4</sup>, a steady estimate of sensitivity for the IgM test cannot be obtained. Sensitivity estimates will vary very widely with time since infection. As we made no assumptions about the time of infection of those who tested positive, we are unable to conclude about the contextual sensitivity of the test. This caveat also applies to the IgG test, but the slower decline of IgG antibodies means the assumption of stable test performance is more plausible in that case.

Confidence intervals for test-adjusted estimates are Lang-Reiczigel intervals<sup>5</sup>, which take into account the sample size of the antibody test validation study. Other confidence intervals for prevalences are Wilson score intervals, applied using the DescTools R package<sup>6</sup>.

Household-level statistics were calculated based on the answers of household representatives. The household representative was taken to be:

- the first interviewed household head; and, if no head is present,
- the first interviewed adult



Data were processed and analyzed using R version 3.6.1 and RStudio. We will use direct standardization on the age and population weights and a multilevel regression which adjusts for the specificity and sensitivity of the test. Bayesian analysis will be done using the rjags package.

The current preliminary report only includes descriptive statistics and crude estimates. Bayesian adjustments will be included in the final report.

## Ethical considerations

The study protocol obtained the ethical clearance and the administrative authorization of the Ministry of Health of Cameroon. Every adult participant (21 years or above) signed an informed consent. For minors, a person with parental authority was asked to sign the consent form and, if the age was equal to or above 15 years, an assent was also requested. Questionnaires were coded and names of participants were recorded in a confidential list available only to the study team. Before starting the study, all the team members were trained on research ethics, good clinical practices and study protocol and procedures.

## Results

### Participation rate

From 14, October to 26, November 2020, 256 households were visited, of which 63 (24%) refused to give any information, one was excluded because not in the selected sample. In the end, 192 households were included for a total of 1007 participants.

Of the 192 included households, 128 were the originally sampled by the random system, while the remaining 64 (33%) were replaced through standard procedures because the identified buildings were non-residential.

**Table 1: Households and individuals expected and included in the study.**

	Inhabitants	N of household sampled	N of inclusions expected	N of households included	N of people included
Briqueterie	52 850	31	124	24 (77%)	109 (88%)
Carrière	104 785	61	244	45 (73%)	240 (98%)
Cité verte	38 832	22	88	24 (109%)	88 (100%)
Messa	18 899	11	44	9 (81%)	48 (109%)
Mokolo	46 960	27	108	17 (63%)	96 (89%)
Nkomkana	41 953	24	96	15 (62%)	81 (84%)
Tsinga	35 349	20	80	16 (80%)	83 (103%)
Tsinga Oliga	37 572	22	88	14 (63%)	71 (80%)
Ekoudou	55 659	32	128	28 (87%)	191 (149%)
<b>Total</b>	<b>432 859</b>	<b>250</b>	<b>1000</b>	<b>192</b>	<b>1007</b>

All the health areas of the district were surveyed, but participation varied between the neighbourhoods (Table 1). All participants were to be tested for SARS-CoV2-antibodies, but in 35 cases (3%), some members of the household, despite responding to the questionnaire, refused the test.

## Respondent characteristics

Participants had a mean age of 29 years and 570 (56.6%) were women. The demographic characteristics of respondents are summarized in Table 2.

Households had a mean of 4.6 adult inhabitants (15 years or above) and 1.8 children. In the majority (92%) of the visited households, the head of the family was surveyed and in 136 households (70.8%), we accessed the person providing the main source of income. This was most often the father (55.7% of households) or the mother (36.5%).

Household members were most commonly in school (39.5%), working “small jobs” (21%) or working as traders (12.4%). This explains the high percentage of households who said their income arrives through irregular and informal sources (41%). Interestingly, despite only 5.7% of respondents reported regular waged work, formal regular income was the main source of revenues for 23% of the households, showing the possible contribution of other family members.

Only six households reported a prior positive COVID-19 test among their members, but 67 (7%) respondents reported having been tested for SARS-CoV-2 before, with nearly half of these as part of governmental mass testing campaigns.

## Seroprevalence estimates

Of the 970 respondents tested for IgG and IgM antibodies, 340 (35.1%) tested were seropositive for at least one antibody type (Figure 1). IgM seropositivity was quite low, and the overlap between IgG and IgM seropositivity was minimal; among the 32 individuals who were IgM positive, only 6 were also IgG positive.




Figure 1: Seropositivity of respondents by test type

**Table 2:** Sociodemographic characteristics of the final sample

Characteristic	Female	Male	Total
<b>n</b>	570	437	1007
Num. household heads	77	101	178
Mean Age (SD)	29.6 (17.4)	28.5 (17.6)	29.1 (17.5)
Mean BMI (SD)	27.3 (26.7)	23.2 (6.26)	25.5 (20.6)
<b>Age groups, n (% of sex in each group)</b>			
5-15	137 (24%)	128 (29.3%)	265 (26.3%)
16-29	186 (32.6%)	136 (31.1%)	322 (32%)
30-45	140 (24.6%)	89 (20.4%)	229 (22.7%)
46-65	79 (13.9%)	66 (15.1%)	145 (14.4%)
>65	28 (4.9%)	18 (4.1%)	46 (4.6%)
<b>Education Level, n (% of sex in each group)</b>			
Secondary	258 (45.3%)	184 (42.1%)	442 (43.9%)
Primary	191 (33.5%)	137 (31.4%)	328 (32.6%)
University	76 (13.3%)	81 (18.5%)	157 (15.6%)
No formal instruction	35 (6.1%)	18 (4.1%)	53 (5.3%)
Doctorate	7 (1.2%)	13 (3%)	20 (2%)
Other	3 (0.5%)	4 (0.9%)	7 (0.7%)
<b>Profession, n (% of sex in each group)</b>			
Student	216 (36.5%)	202 (43.3%)	418 (39.5%)
Small jobs	86 (14.5%)	136 (29.1%)	222 (21%)
Business/Trade	81 (13.7%)	50 (10.7%)	131 (12.4%)
Housewife	74 (12.5%)	0 (0%)	74 (7%)
Unemployed	52 (8.8%)	21 (4.5%)	73 (6.9%)
Salaried worker	38 (6.4%)	22 (4.7%)	60 (5.7%)
Retired	18 (3%)	17 (3.6%)	35 (3.3%)
Other	17 (2.9%)	15 (3.2%)	32 (3%)
Farmer	10 (1.7%)	4 (0.9%)	14 (1.3%)


**Figure 2:** Seropositivity in each age-sex stratum. Positive cases include individuals positive for either IgG or IgM antibodies

Figure 2 shows overall antibody seroprevalence (either IgG or IgM) in each age-sex stratum. Seroprevalence is higher in men, and increases with age. The highest values were seen in men above 65, where the seropositivity was 50%, although the sample size for that group is notably small.

Tables 3 and 4 show the adjustments for the age-sex distribution of the Yaounde population and for diagnostic test performance. Since women were oversampled relative to their share of the population (56.6% of sample was female), and women showed a lower seroprevalence, the crude estimates were downwardly biased. Thus, population weighting increased the overall estimate of seropositivity in nearly all age categories for both the IgG and IgM assays. Adjustments for specificity and sensitivity also increased the estimates slightly.

Only one of the 21 RNA PCR tests was positive.

**Table 3: Population-weighted and test-adjusted seroprevalence estimates for anti-SARS-CoV-2 IgG antibodies**


	Sample size, n	Seropos., n	Seroprevalence (95% Confidence Interval)		
			Crude	Population-weighted	Population-weighted, test-adjusted
<b>Total</b>	971	302	31.1% (28.3 - 34.1)	31.3% (28.4 - 34.3)	33.7% (29.5 - 38.0)
<b>Female</b>	549	154	28.1% (24.5 - 32.0)	28.0% (24.4 - 31.9)	30.1% (25.4 - 35.3)
<b>Male</b>	422	148	35.1% (30.7 - 39.7)	34.6% (30.2 - 39.3)	37.4% (32.4 - 44.2)
<b>5 - 14</b>	241	69	28.6% (23.3 - 34.6)	28.7% (23.3 - 34.7)	30.8% (24.3 - 38.1)
<b>15 - 29</b>	325	98	30.2% (25.4 - 35.4)	30.7% (25.9 - 35.9)	33.1% (26.6 - 39.0)
<b>30 - 44</b>	212	69	32.5% (26.6 - 39.1)	32.7% (26.7 - 39.3)	35.3% (28.0 - 43.2)
<b>45 - 64</b>	153	51	33.3% (26.4 - 41.1)	34.1% (27.0 - 41.9)	36.8% (27.8 - 45.4)
<b>65 +</b>	40	15	37.5% (24.2 - 53.0)	39.4% (25.8 - 54.8)	42.6% (25.6 - 58.6)

**Table 4: Population-weighted seroprevalence estimates for anti-SARS-CoV-2 IgM antibodies**

	Sample size, n	Seropos., n	Seroprevalence (95% Confidence Interval)	
			Crude	Population-weighted
<b>Total</b>	953	32	3.4% (2.39 - 4.7)	3.4% (2.43 - 4.8)
<b>Female</b>	549	17	3.1% (1.94 - 4.9)	3.0% (1.90 - 4.8)
<b>Male</b>	404	15	3.7% (2.26 - 6.0)	3.8% (2.32 - 6.1)
<b>5 - 14</b>	241	5	2.1% (0.89 - 4.8)	2.1% (0.89 - 4.8)
<b>15 - 29</b>	325	12	3.7% (2.12 - 6.3)	3.6% (2.07 - 6.2)
<b>30 - 44</b>	211	7	3.3% (1.62 - 6.7)	3.9% (2.02 - 7.5)
<b>45 - 64</b>	152	6	3.9% (1.82 - 8.3)	4.1% (1.94 - 8.6)
<b>65 +</b>	24	2	8.3% (2.32 - 25.8)	8.3% (2.32 - 25.8)

## COVID-19 related symptoms


Three hundred and two (30%) respondents reported having at least one symptom compatible with SARS-CoV-2 infection (frequency of symptoms is reported in Figure 1).



**Figure 3: Common symptomatologies reported by respondents**

Among those who were seropositive, the most common symptoms reported were fever (18.3%), headache (18%) and cough (18%), and all three were significantly more common in seropositive and seronegative individuals (Figure 4). Surprisingly, agnosia or ageusia was only experienced by 4% of the seropositive respondents (and 2% of the seronegative respondents).

## Impact on households



Concerning the impact of the epidemic on the households, 163 households (85%) reported that their income had fallen since March 1st. Only in eight households (4%) was someone reported to have completely lost the source of income, and 24 (12%) reported no change at all.

**Figure 4: Frequency of symptoms among seropositive and seronegative individuals.** \* p

Of the included households, 11 reported a death in the household during the period of the epidemic. The mean age of people who died was 69.5 years and none of these deaths was reported to be COVID-related.

Most of these (83%) were considered to be of moderate severity while for 44 (15%) they were severe, but only 75 (7%) sought medical attention; the majority (62%) used self-medication on the advice of a member of the family or a significant other. Paracetamol, traditional medicines and antibiotics were the most commonly used remedies for those who had symptoms (Figure 2).




Figure 2. Use of drugs in respondents with symptoms

Among those who actually consulted formal health services, none went to a specialized COVID-19 care center; 40% visited private clinics and 20% visited general public hospitals. Forty-seven participants reported hospitalization, but only one hospitalization was in relation to COVID-19. Very few individuals reported additional health expenditure due to COVID-19; in cases when this was reported, the additional expenses were mainly for the purchase of personal protective equipment (7 participants paid for transport, 2 for the test and 2 for consultation).

One hundred eight participants (11%) reported suffering from a chronic condition, mainly hypertension (34%), respiratory conditions (16%) or diabetes (10%). With a mean BMI of 27.3 (versus 22.5 for men), women showed a higher frequency of overweight. Only 2 respondents reported HIV infection.

In the group of people with chronic conditions, 14.8% experienced disruption in the continuity of care and 11% had difficulties in the access to their regular treatment.

In seeking care either for an acute event or for a chronic condition, a number of obstacles were reported. Fifty-four individuals (15%) answered that they had encountered financial difficulty in paying for care since March 1st, and 45 (12%) individuals said financial difficulty caused them to delay care. Notably, only 41 respondents (4%) have medical insurance.


Despite the fact that 57.8% reported complete or partial compliance with preventive measures (see Figure 3), the 972 serological tests done showed an IgG positivity rate of nearly 30% (302), while 3% (32) were positive for IgM antibodies. When seroprevalence is broken down by age and sex (Figure 4), the group with the highest prevalence is men between age 50 and age 70.

Adjustments for test performances will need to be done to take into account test specificity and sensitivity.



**Do you follow the basic hygiene protocols (wash hands frequently, sneeze in your elbow, use a disposable tissue, etc.)**


Definitely yes Partly Definitely not




**Do you follow social distancing rules (avoid handshakes or hugs, avoid unnecessary outings, etc.)**



**Since March 1st, have you stayed confined?**



**Figure 3:** Proportion of respondents following prevention measures according to serology test result



**Figure 4:** Positive serological test by sex and age.

There were no major differences in sociodemographic characteristics between exposed and not exposed participants (Table 3); but further statistical analysis is necessary to test for significance. There may be slightly lower risk of exposure in the group with no formal education. Surprisingly, seropositive individuals do not show a much greater prevalence of symptoms than seronegative individuals.

Concerning the impact of the epidemic on the daily living of the studied communities, 560 of the respondents (56%) reported having to reduce daily activities, including 492 (49%) who reported reduced work hours because of confinement. About 82% of the households reported a reduction of living income.

Participants attitudes towards the COVID-19 were notable: around 825 (82%) reported fear of contamination, but only 48% said they fully respected the basic prevention measures like washing hands or cough etiquette, and only 23% said they fully followed the rules of physical distancing. Nearly half of the participants believed that people with COVID-19 are stigmatized, and 7% reported that they would conceal their illness to avoid this stigma.

Notably, more than 50% (543) of respondents reported an increase in stress and pressure in the family environment with 343 (34%) notifying having been victims of psychological or physical violence.

Facing all these constraints, external support was very limited, only 94 (9%) report having received external help. This help was primarily monetary, or in the form of free personal protective equipment. And the most common sources of these were the government and non-governmental organizations (53%) or other members of the family (40%).

**Table 3:** Demographics and symptomatology of seropositive (IgG or IgM) and seronegative respondents.

Characteristic	Seropositive	Seronegative
n	319	623
Mean Age (SD)	30.6 (17.9)	28.2 (17)
Mean BMI (SD)	27.4 (33)	24.6 (11)
<b>Sex groups, n (% of all positive/negatives that are of each sex)</b>		
Female	164 (51.4%)	371 (59.6%)
Male	155 (48.6%)	252 (40.4%)
<b>Age groups, n (% of all positive/negatives that are in each age group)</b>		
5-15	76 (23.8%)	175 (28.1%)
16-29	100 (31.3%)	196 (31.5%)
30-45	76 (23.8%)	143 (23%)
46-65	50 (15.7%)	87 (14%)
above 65	17 (5.3%)	22 (3.5%)
<b>Education Level, n (% of all positive/negatives that are in each education class)</b>		
Secondary	135 (42.3%)	287 (46.1%)
Primary	114 (35.7%)	195 (31.3%)
University	52 (16.3%)	86 (13.8%)
No formal instruction	11 (3.4%)	40 (6.4%)
Doctorate	6 (1.9%)	10 (1.6%)
Other	1 (0.3%)	5 (0.8%)
<b>Profession, n (% of all positive/negatives that are in each profession class)</b>		
Student	124 (38.9%)	265 (42.5%)
Small trader	78 (24.5%)	137 (22%)
Businessperson	42 (13.2%)	80 (12.8%)
Home-maker	26 (8.2%)	46 (7.4%)
Unemployed	25 (7.8%)	44 (7.1%)
Salaried worker	21 (6.6%)	31 (5%)
Retired	9 (2.8%)	22 (3.5%)
Other	8 (2.5%)	20 (3.2%)
Farmer	5 (1.6%)	9 (1.4%)
<b>Symptoms, n (% of all positive/negatives that reported each symptom)</b>		
No symptoms	205 (64.3%)	456 (73.2%)
Fever	58 (18.2%)	74 (11.9%)
Headache	59 (18.5%)	69 (11.1%)
Cough	58 (18.2%)	65 (10.4%)
Runny or stuffy nose	38 (11.9%)	45 (7.2%)
Sneezing	23 (7.2%)	20 (3.2%)
Fatigue	14 (4.4%)	24 (3.9%)
Muscle pain	18 (5.6%)	17 (2.7%)
Smell or taste loss	13 (4.1%)	12 (1.9%)
Sore throat	8 (2.5%)	10 (1.6%)
Stomach ache	8 (2.5%)	8 (1.3%)
Diarrhoea	2 (0.6%)	11 (1.8%)
Nausea, Vomiting	4 (1.3%)	9 (1.4%)
Shortness of breath	1 (0.3%)	4 (0.6%)
Other	1 (0.3%)	3 (0.5%)

## Conclusions

SARS-CoV-2 appears to have circulated quite broadly in the Cité Verte district of Yaoundé. Nearly one person in three was found to be IgG or IgM positive, but more than 60% of these people had no symptoms between March and the date of survey, confirming that the asymptomatic form of the disease is more frequent than severe forms.

Survey results do not suggest significant health service disruption during the pandemic period. However, this may be explained by the fact that study participants did not seek out formal care when they had COVID-19 symptoms, turning instead to family and neighbors for advice on treatment, and using easily-acquired drugs and traditional remedies.

While the epidemic did not seem affect the functioning of health services, it caused significant disruption to household incomes. Such disruptions are unsurprising, given that the majority of households acquire income from the informal sector, whose vibrance depends on the daily activities of the entire public.

Social and family activities were also significantly disrupted and this could have created a situation of conflict and stress, sometimes resulting in violence.

## Bibliography

1. Lu H, Stratton CW, Tang Y. Outbreak of pneumonia of unknown etiology in Wuhan, China: The mystery and the miracle. *J Med Virol*. 2020;92(4):401-402. doi:10.1002/jmv.25678
2. World Health Organization. Weekly epidemiological update - 1 December 2020. Accessed December 8, 2020. <https://www.who.int/publications/m/item/weekly-epidemiological-update---1-december-2020>
3. AFP. WHO warns Africa is ill-equipped to deal with coronavirus due to “weaker health systems.” *TheJournal.ie*. Accessed December 8, 2020. <https://www.thejournal.ie/world-health-organisation-african-coronavirus-5017867-Feb2020/>
4. Umvilighozo G, Mupfumi L, Sonela N, et al. Sub-Saharan Africa preparedness and response to the COVID-19 pandemic: A perspective of early career African scientists. *Wellcome Open Res*. 2020;5. doi:10.12688/wellcomeopenres.16070.2
5. Stringhini S, Wisniak A, Piumatti G, et al. Seroprevalence of anti-SARS-CoV-2 IgG antibodies in Geneva, Switzerland (SEROCoV-POP): a population-based study. *The Lancet*. 2020;396(10247):313-319. doi:10.1016/S0140-6736(20)31304-0
6. Dortet L, Ronat J-B, Vauloup-Fellous C, et al. Evaluating Ten Commercially-Available SARS-CoV-2 Rapid Serological Tests Using the STARD (Standards for Reporting of Diagnostic Accuracy Studies) Method. *J Clin Microbiol*. Published online November 25, 2020. doi:10.1128/JCM.02342-20
7. Chazai Partners. *Recueil Des Mesures Prises Par Le Gouvernement Camerounais Dans Le Cadre de La Lutte Contre La COVID-19*. Accessed December 8, 2020. <https://www.chazai-partners.com/wp-content/uploads/2020/05/Chazai-Partners-Recueil-des-mesures-prises-par-le-gouvernement-Camerounais-dans-la-lutte-contre-la-COVID-19-29-05-20.pdf>