

| Population of interest | sample_frame_age | denominator_value | seroprevalence_95_percent_ci | sampling_method | test |
|--------------------------|----------------------|-------------------|------------------------------|------------------------|------|
| | Adults (18-64 years) | 66 | 26% (NR) | Unclear | He |
| and caregivers | Adults (18-64 years) | 74 | 12.2% (NR) | Convenience | Ar |
| and caregivers | Multiple groups | 77 | | Stratified probability | Gu |
| | Multiple groups | 99 | 3% (0.6-8.6%) | Convenience | Ab |
| for non-COVID-19 reasons | Multiple groups | 99 | 3% (0.6-8.6%) | Convenience | Ab |
| | Multiple groups | 142 | 4.2% (NR) | Simplified probability | Gu |
| community samples | Multiple groups | 185 | 25.4% (NR) | Stratified probability | Ca |
| and caregivers | Adults (18-64 years) | 203 | 18.2% (NR) | Convenience | Ar |
| community samples | Adults (18-64 years) | 301 | 8.8% (5.5-11.6%) | Simplified probability | Zh |
| care workers | Adults (18-64 years) | 405 | 23.7% (NR) | Simplified probability | Ab |
| and caregivers | Adults (18-64 years) | 500 | 12.3% (8.2-16.5%) | Convenience | On |
| community samples | Adults (18-64 years) | 754 | 19.7% (NR) | Self-referral | W |
| s and unemployed persons | Adults (18-64 years) | 955 | 0.9% (0.4-1.8%) | Convenience | "H |
| | Adults (18-64 years) | 1.69e+03 | 25.1% (NR) | Convenience | AA |
| | Adults (18-64 years) | 2.79e+03 | 44.6% (42.6-46.6%) | Convenience | Ro |
| | Multiple groups | 3.1e+03 | 5.2% (3.7-7.1%) | Convenience | Kr |
| | Adults (18-64 years) | 3.1e+03 | 4.3% (2.9-5.8%) | Sequential | N/ |
| and caregivers | Multiple groups | 4.04e+03 | 1.3% (NR) | Convenience | Ar |