

Cell culture:

I have reviewed and understand [Tissue Culture Basics](#)

I have reviewed and understand [Minimizing tissue culture contaminations](#)

**HEK293T (human; overexpression studies)**

Media: DMEM + 1% L-glutamine + 10% FCS

Subculture: 1:5 – 1:10

\*cells can be detached by rinsing with media; be sure to mix well to avoid cell clumps

**Jurkats (human, T cell line)**

Media: RPMI1640 + 1% L-glutamine + 10% FCS

Subculture: suspension cells so culture between  $2 \times 10^5$  to  $1 \times 10^6$  cells

**A549 (human, NSCLC)**

Media: DMEM + 1% L-glutamine + 10% FCS

Subculture: 1:5 – 1:10

**DM93 (human, melanoma)**

Media: RPMI1640 + 1% L-glutamine + 10% FCS

Subculture: 1:3 – 1:5

**H82 (human, SCLC)**

Media: RPMI1640 + 1% L-glutamine + 10% FCS

Subculture: suspension cells so culture between  $2 \times 10^5$  to  $1 \times 10^6$  cells

**Min6 (mouse; immortalized  $\beta$ -islet cells)**

Media: DMEM + 1% L-glutamine + 10% FCS + 25  $\mu$ M  $\beta$ ME

Subculture: 1:2 – 1:3; detach using 0.25% trypsin

**RAW264.7 (mouse; macrophage-like cells)**

Media: RPMI1640 + 1% L-glutamine + 10% FCS

Subculture: 1:5 – 1:10; detach by washing 1X with sterile PBS followed by incubation with PBS for 5 min

\*avoid trypsin on these cells since it will compromise viability

**THP1 (human; monocytes)**

Media: RPMI1640 + 1% L-glutamine + 10% FCS + 50  $\mu$ M  $\beta$ ME

Subculture: suspension cells so culture between  $2 \times 10^5$  to  $1 \times 10^6$  cells

**IC-21 (mouse; immortalized peritoneal macrophages)**

Media: RPMI1640 + 1% L-glutamine + 10% FCS

Subculture: 1:2 – 1:5; detach by washing 1X with sterile PBS followed by incubation with PBS for 5 min

\*avoid trypsin on these cells since it will compromise viability

\* Centrifuge cells at 400 x g for 5 min to pellet for both subculture and isolation for biological studies.

**Revised: 08/02/2022**