

C-1.



MASENO UNIVERSITY
UNIVERSITY EXAMINATIONS 2017/2018

**FIRST YEAR END OF YEAR EXAMINATION FOR THE DEGREE OF
BACHELOR OF MEDICINE AND BACHELOR OF SURGERY WITH
INFORMATION TECHNOLOGY**

MAIN CAMPUS

MCS 100: NURSING SKILLS

Date: 22nd November, 2017

Time: 2.00 - 5.00 pm

INSTRUCTIONS:

- Follow the instructions as attached.



NURSING SKILLS**(3 HOURS)****INSTRUCTIONS**

1. The exam consists of **THREE (3)** sections. (Total 100 Marks)
2. Answer **ALL** questions in sections **A, B and C**
3. Answer questions in the **ANSWER SHEET and BOOKLET** provided
4. **SECTION A** has Thirty **(30)** Multiple Choice Questions (MCQs). (30 Marks)
5. **SECTION B** has Six **(6)** Short Answer Questions (SAQs). (30 Marks)
6. **SECTION C** has two **(2)** Long Essay Questions (LEQs). (40 Marks)



Section A: Multiple Choice Questions (MCQs)**(30Marks)**

Select the most appropriate response to the stem question. A correct response earns you one (1) Mark

1. Cidex solution used for high level disinfection should be discarded after:-
 - A. 1 day
 - B. 14 days
 - C. 7 days
 - D. 28 days
 - E. 10 days
2. _____ initiated formal nursing training using a curriculum.
 - A. Virginia Henderson
 - B. Patterson Kussman
 - C. Florence Nightingale
 - D. Laura Johns
 - E. Callista Roy
3. Formal nursing training in Kenya at enrolled and registered levels was started in the year:-
 - A. 1945
 - B. 1950
 - C. 1935
 - D. 1955
 - E. 1970
4. The first university in Kenya to offer Bachelor of Science in Nursing was:-
 - A. Nairobi
 - B. Moi
 - C. Kenyatta
 - D. University of East Africa Baraton
 - E. Uzima
5. The following are some of the ways of overcoming the factors that hinder team collaboration environment:-
 - A. An open attitude, feelings of mutual respect
 - B. Trust, coercion



- C. Mistrust among team members, open attitude
 - D. Coercion, open attitude
 - E. Assertiveness, mistrust
6. The criteria of a profession includes:-
- A. Strong scientific based knowledge, strong service orientation
 - B. A recognized authority by the professional group and with community sanction, no code of ethics
 - C. Professional organization that sets standards, limited research
 - D. Ongoing research, no autonomy
 - E. Autonomy, poor service orientation
7. Performing an act that a reasonably prudent person under similar circumstances would not do is referred to as:-
- A. Negligence
 - B. Battery
 - C. Assault
 - D. Intentional tort
 - E. Malpractice
8. Patient X, 24 years old is to receive 1 gram of oral paracetamol tid. The container is labeled 500 mg (milligrams) for each tab; the expiry date is November, 2015. How many such tablets will you administer today at 9 am:-
- A. 2
 - B. 4
 - C. 1
 - D. 5
 - E. The drug should not be administered
9. The standing order **BD** Means:-
- A. Twice daily / 12 hourly
 - B. Twice daily
 - C. Thrice daily
 - D. Thrice/8 hourly
 - E. Administer when necessary

10. The rationale for adjusting the scale against the client's height is to:-
- A. Avoid zero error
 - B. To enable measurement of client's full height
 - C. To promote cooperation
 - D. To help in building up client's self esteem
 - E. To minimize false reading
11. Calculate the body mass index (BMI) of a client who weighs 60 kg, with a height of 166cm:-
- A. 40 KG/M²
 - B. 21 KG/M²
 - C. 36.14 KG/M²
 - D. 21.77 KG/M²
 - E. 43.58 KG/M²
12. The normal blood pressure range of a 1-2 years old child is:-
- A. 75-100/50-70
 - B. 80-110/50-80
 - C. 80-110/50-80
 - D. 85-120/50-80
 - E. 95-120/60-80
13. During resuscitation of an adult whereby there is only one rescuer, 1 cycle of CPR consists of:-
- A. 15 compressions to two breaths
 - B. 15 compressions to one breath
 - C. 30 compressions to one breath
 - D. 30 compressions to two breaths
 - E. 5 compressions to 1 breath
14. The manoeuvre that is used to open the airway of an unconscious patient with suspected neck injury is:-
- A. Head tilt-chin lift
 - B. Jaw thrust
 - C. Chest thrust
 - D. Chin lift
 - E. Log roll

15. The correct position of your hands when giving abdominal thrusts to a conscious adult who is choking is:-
- A. On the centre of the chest
 - B. In the middle of the abdomen just above the navel
 - C. Below the navel
 - D. Anywhere on the abdomen
 - E. The left anterior chest
16. A pregnant woman, 30 weeks gestation is choking but is conscious and cannot cough or speak. What should the rescuer do?
- A. Lower the casualty to the floor and remove the object, if visible at the back of the throat.
 - B. Give chest thrusts until the object is dislodged or the casualty becomes unconscious
 - C. Encourage the casualty to cough to try and dislodge the object.
 - D. Give abdominal thrusts until the object is dislodged or she becomes unconscious.
 - E. Take the casualty to the emergency department
17. If a foreign body blocks a child's airway and the child is coughing forcefully and able to speak, what should the rescuer do?
- A. Give abdominal thrusts
 - B. Slap her on the back until she coughs up the object
 - C. Encourage her to continue coughing
 - D. Lower her to the floor, check for and remove any visible object
 - E. Give chest compressions to dislodge the object.
18. One of your friends has cut his arm and is bleeding severely. The immediate action is to:-
- A. Put the injured limb in some cold water
 - B. Apply direct pressure over the wound
 - C. Give him paracetamol 1 gram per oral
 - D. Apply a tourniquet above the wound
 - E. Take him to the emergency department
19. The correct treatment for a choking infant (under 12 months) is:-
- A. Alternate back blows with abdominal thrusts
 - B. Up to 5 back blows with up to 5 chest thrusts

- C. Chest thrusts only
 - D. Back blows only
 - E. Immediate surgery
20. The immediate action for a child who appears to have broken his/her arm is to:-
- A. Wash the arm in cold water to reduce swelling and take him/her to see a doctor
 - B. Tightly bandage the arm to reduce swelling and call an ambulance
 - C. Steady and support the arm, reassure the child and seek medical aid
 - D. Wait for 30 minutes to see if pain subsides – it may only be a sprain
 - E. Try to realign the fractured bone manually
21. Which is the first step when caring for bleeding wounds?
- A. Apply direct pressure with a clean or sterile dressing.
 - B. Apply pressure at the pressure point.
 - C. Add bulky dressings to reinforce blood soaked bandages.
 - D. Elevate the wound.
 - E. Use a tourniquet to stop all blood flow
22. The planning phase of the nursing process involves:-
- A. Assessing and diagnosing
 - B. Evaluating goal achievement and documenting
 - C. Performing nursing actions and documenting
 - D. Setting goals and selecting interventions
 - E. Implementation and evaluation
23. Collaborative interventions are therapies that require:-
- A. Physician and nurse intervention
 - B. Nurse and client interventions
 - C. Client and physician interventions
 - D. Multiple health care professionals
 - E. Client and family interventions
24. Core body temperature is the measurement of:-
- A. The skin surface
 - B. The forehead



- C. The body's internal environment
 - D. The axilla
 - E. The extremities
25. Sweating helps the body to:-
- A. Warm the body up
 - B. Cool the body
 - C. Get rid of toxins
 - D. Maintain constant heart rate
 - E. Get rid of excess water
26. Pulse measured at the apex of the heart is referred to as:-
- A. Apical
 - B. Radial
 - C. Apex
 - D. Temporal
 - E. Femoral
27. Radial artery is found:-
- A. Behind the knee
 - B. Besides the thighs
 - C. Either sides of the neck
 - D. Lateral side of the legs
 - E. Lateral side of the forearm
28. Intra dermal means:-
- A. Beneath the skin
 - B. In the vein
 - C. Within the vein
 - D. Through the skin
 - E. Beneath the muscle layer
29. _____ considered the 5th vital sign.
- A. Height and weight
 - B. Respiration
 - C. Body mass index
 - D. Pain
 - E. Pulse
30. The normal respiratory rate for an adult is:-
- A. 10 – 15 breaths per minute
 - B. 8 – 12 breaths per minute
 - C. 12 – 20 breaths per minute
 - D. 14 – 24 breaths per minute
 - E. 15 – 30 breaths per minute

Section B: Short Answer Questions (SAQ's)**30 Marks**Answer **ALL** Questions in this section

1. Define the following terms:
 - a. Nursing (1 Mark)
 - b. Professionalism (1 Mark)
2. State four (4) considerations observed in order to prevent infection during administration of an injection (4 Marks)
3. Outline four (4) principles of management of soft tissue injuries. (4 Marks)
4. a. Define the term Euthanasia. (1 Mark)
b. Explain the two types of Euthanasia. (4 Marks)
5. Compare and contrast medical diagnosis and nursing diagnosis giving relevant examples (10 Marks)
6. The patient's prescription reads, 1g of Ceftriaxone IV, O.D, 1/52. The pharmacy stock has 250mg/2mls dosage of Ceftriaxone.
 - a. Interpret the above prescription (2 Marks)
 - b. Calculate the amount in volume (3 Marks)



LONG Essay Questions (LEQ's)**40 Marks****Answer ALL Questions in this section.**

1. Patient O, a known hypertensive, comes to the facility where you are attached as a medical student with complaints of headache, dizziness, restlessness and fatigue. You are asked by the in-charge to take his blood pressure.
 - a. Outline the procedure of measuring arterial blood pressure using an aneroid sphygmomanometer giving a rationale for each step (14 Marks)
 - b. State any six (6) factors that influence blood pressure of a client (6 Marks)
2. A man, carrying out repairs on a building while on a ladder falls from the ladder to the ground and appears to be unconscious. You and other passers-by rush to his aid.
 - a. Explain six (6) steps you will take in management of this casualty. (12 Marks)
 - b. Describe two ways of opening the airway and indicate which would be most suitable for this casualty. (4 Marks)
 - c. Outline four differences when giving CPR to an infant compared to an adult. (4 Marks)





MASENO UNIVERSITY
UNIVERSITY EXAMINATIONS 2017/2018

**FOURTH YEAR EXAMINATION FOR THE DEGREE OF
BACHELOR OF MEDICINE & BACHELOR OF SURGERY
WITH INFORMATION TECHNOLOGY**

CITY CAMPUS – SUPPLEMENTARY EXAM

MCS 400: CLINICAL PHARMACOLOGY II

Date: 22nd May, 2018

Time: 9.00 - 12.00pm

INSTRUCTIONS:

- As per the attached.



CLINICAL PHARMACOLOGY II**(3 HOURS)****GENERAL INSTRUCTIONS**

1. Write your University Registration Number on every piece of paper you use.
2. Do not write your name on any piece of paper you use.
3. The questions are set out in **THREE (3)** sections headed Section A: MCQs,
Section B: SAQs and Section C : MEQs (120 Marks)
4. **SECTION A** has sixty (**60**) Multiple Choice Questions (MCQs). (60 Marks)
5. **SECTION B** has SEVEN (7) Short Answer Questions (SAQs). (40 Marks)
6. **SECTION C** has TWO (2) Modified Essay Questions (MEQs). (20 Marks)
7. Answer MCQs only in this question PAPER. SAQs and MEQs are to be answered in the ANSWER BOOKLET provided.
8. Read carefully the additional instructions preceding each section.

SECTION A: MULTIPLE CHOICE QUESTIONS (MCQs)**(60 MARKS)****INSTRUCTIONS:-**

i. EACH OF THE QUESTIONS IN THIS SECTION CONSISTS OF A STATEMENT/QUESTION WHOSE MOST APPROPRIATE ANSWER IS PROVIDED AMONG THE FIVE OPTIONS NUMBERED a) to e).

ii. CIRCLE THE MOST APPROPRIATE ANSWER.

1. The following antiemetic used in cancer chemotherapy acts by blocking both dopamine and serotonin receptors:
 - a. Ondansetron
 - b. Domperidone
 - c. Metoclopramide
 - d. Dexamethasone
 - e. Cyclizine
2. The toxic effects of methotrexate on normal cells may be reduced by administering:
 - a. N-acetyl-L-cysteine
 - b. Vitamin K
 - c. Penicillamine
 - d. Leucovorin
 - e. Deferoxamine
3. Which of the following chemotherapeutic agents is a potent vesicant?
 - a. Doxorubicin
 - b. Etoposide

- c. Cyclosporin
 - d. Cyclophosphamide
 - e. Dacarbazine
4. The following chemotherapeutic agent is effective in the management of brain tumours:
- a. Methotrexate
 - b. Chlorambucil
 - c. Nitrosoureas
 - d. Ifosphamide
 - e. Azathioprine
5. A keratolytic agent used in management of psoriasis:
- a. Crotamiton
 - b. Sodium sulfacetamide
 - c. Tolnaftate
 - d. Propylene glycol
 - e. Selenium sulfide
6. All the following drugs may cause a drug-induced photosensitivity reaction on the skin EXCEPT:
- a. Chlorpromazine
 - b. Amitriptyline
 - c. Methotrexate
 - d. Sulfadoxine
 - e. Tetracycline

7. Which of the following dermatological agent does NOT match its mechanism of action?
- a. Pimecrolimus - Inhibits T-lymphocyte activation
 - b. Doxepin – Inhibits H_1 receptors
 - c. Eflornithine - Inhibits tyrosine decarboxylase
 - d. Mequinol – Inhibits tyrosinase
 - e. Naftifine – Inhibits squalene epoxidase
8. A class of agents indicated in treatment of most anxiety states is:
- a. Barbiturates
 - b. Benzodiazepines
 - c. Lithium salts
 - d. Phenothiazines
 - e. Chloral hydrate
9. The mechanism of hypnotic benzodiazepine action:
- a. Increasing the duration of the GABA-gated Cl^- channel openings
 - b. NMDA (N-methyl D aspartate) receptor blocker
 - c. Directly activating the chloride channels
 - d. Increasing the frequency of Cl^- channel opening events
 - e. N- and L-calcium channel blocker
10. The following statement does not limit using benzodiazepines as anti-anxiety agents?
- a. Tendency to develop psychologic dependence
 - b. A high risk of drug interactions based on liver enzyme induction
 - c. Synergic CNS depression with concomitant use of other drugs

- d. Formation of active metabolites
- e. Muscle relaxing effects

11. The following drug is not suitable for the management of Status Epilepticus:

- a. Diazepam
- b. Lamotrigine
- c. Lorazepam
- d. Phenobarbital
- e. Phenytoin

12. Which of the following anti-convulsants is a prodrug?

- a. Gabapentin
- b. Primidone
- c. Topiramate
- d. Ethosuximide
- e. Tiagabine

13. Which of the following statements is INCORRECT concerning sodium valproate?

- a. Increases GABA synthesis and decreases degradation
- b. Causes weight gain among patients
- c. Is hepatotoxic
- d. May prolong bleeding time
- e. Decreases serum level of Phenobarbital

14. The following are non biologic disease modifying anti-rheumatic drugs EXCEPT?

- a. Anakinra

- b. Gold salts
- c. Azathioprine
- d. D-Penicillamine
- e. Minocycline

15. Rituximab's mechanism of action:

- a. Inhibits pyrimidine synthesis
- b. Blocks the pro inflammatory cytokine TNF-alpha
- c. Blocks interleukin-1 (IL-1)
- d. Depletes peripheral B cells
- e. Inhibits formation of RNA and DNA

16. The following drugs are indicated in management of Systemic Lupus Erythematosus with CNS involvement and diffuse proliferative renal disease EXCEPT:

- a. Methylprednisolone
- b. Azathioprine
- c. Hydroxychloroquine
- d. Cyclophosphamide
- e. Mycophenolate mofetil

17. Which of the following statement is INCORRECT with regard to the management of connective tissue disorders?

- a. Methotrexate is very effective therapy for skin involvement
- b. D-Penicillamine is an α -amino acid metabolite of penicillin with antibiotic properties
- c. β -blockers should be avoided in patients with systemic sclerosis

- d. Treatment of most of these diseases requires anti-inflammatory or immunosuppressive drugs
- e. Drug doses may need to be modified frequently

18. The following drug is indicated in the prevention of Acute uric acid nephropathy:

- a. Vasopressin
- b. Cyclophosphamide
- c. Allopurinol
- d. Amiloride
- e. Sodium Chloride

19. The following drug can precipitate a gouty attack:

- a. Acetazolamide
- b. Allopurinol
- c. Triamterene
- d. Spironolactone
- e. Furosemide

20. Adverse effects of furosemide include the following EXCEPT:

- a. Hyperglycemia
- b. Tinnitus
- c. Fluid and electrolyte imbalance
- d. Hypotension
- e. Metabolic acidosis

21. An opioid antagonist:

- a. Fentanyl
- b. Naloxone
- c. Nabumetone
- d. Naproxen
- e. Nalbuphine

22. The following constitute pharmacological effects of opioids EXCEPT:

- a. Sedation
- b. Miosis
- c. Cough Suppression
- d. Emesis
- e. Diarrhoea

23. A drug used for relieving pain in patients with spinal metastatic disease:

- a. Fentanyl
- b. Dexamethasone
- c. Amitriptyline
- d. Doxepine
- e. Phenobarbital

24. A drug that reverses opioids withdrawal symptoms:

- a. Clonidine
- b. Prazosin
- c. Phenoxybenzamine
- d. Guanethidine

e. Guanfacine

25. The following statements are correct about Tramadol EXCEPT:

- a. Is a weak opioid analgesic
- b. Prolonged used leads to development of tolerance
- c. It acts by blocking reuptake of norepinephrine and serotonin
- d. Is indicated for management of severe pain
- e. It has low incidence of respiratory depression and GI dysmotility

26. Which of the following is NOT a short acting β_2 agonist?

- a. Isoproterenol
- b. Formoterol
- c. Terbutaline
- d. Metaproterenol
- e. Pirbuterol

27. The following drug is NOT a bronchodilator:

- a. Ipratropium bromide
- b. Theophylline
- c. Formoterol
- d. Sodium cromoglycate
- e. Salmeterol

28. Inhaled salbutamol produces the following effect:

- a. Inhibits antigen-antibody reaction in the lungs
- b. Causes bronchodilatation

- c. Reduces bronchial hyperactivity
- d. Inhibits phosphodiesterase enzyme
- e. Inhibits phospholipase

29. Zileuton:

- a. Increases arachidonic acid breakdown
- b. Increases phospholipase activity
- c. Decreases arachidonic acid plasma levels
- d. Decreases leukotriene plasma levels
- e. Increases leukotriene receptor blockade

30. The mechanism of action of albuterol in management of Chronic Obstructive Airway Disease:

- a. Stimulate adenylyl cyclase
- b. Inhibit cyclic adenosine monophosphate
- c. Inhibit cyclic guanosine monophosphate
- d. Stimulate phosphodiesterase
- e. Activate adenosine receptors

31. An appropriate antibiotic regimen for an ICU patient with pneumonia where methicillin resistant *Staphylococcus aureus* (MRSA) is a consideration is:

- a. Cefotaxime plus azithromycin plus linezolid
- b. Ceftriaxone plus clarithromycin plus amikacin
- c. Ampicillin plus aztreonam plus amoxicillin
- d. Cefotaxim plus clarithromycin plus aztreonam
- e. Ceftriaxone plus levofloxacin plus tobramycin

32. Concerning antibiotic:

- a. Linezolid has greater activity against gram negative bacteria
- b. Combination of vancomycin and ampicillin-sulbactam is effective in *Pseudomonas aeruginosa* infection
- c. Aztreonam is a recommended alternative in patients allergic to penicillins
- d. Cefamandol and cefoxitin are third generation agents with broad coverage
- e. Cilastatin has intrinsic antibacterial effect when combined with imipenem

33. Which of the following drug classes would be indicated for the management of chronic stable angina in an asthmatic patient?

- a. Calcium channel blockers
- b. Angiotensin converting enzyme inhibitors
- c. Angiotensin receptor blockers
- d. β blockers
- e. Alpha blockers

34. In the management of dyslipidaemia, which drug class has the greatest potential to increase high density lipoprotein levels (HDL)?

- a. Fibrates
- b. Bile acid sequestrants
- c. Nicotinic acid derivatives
- d. Statins
- e. Fish oils

35. Which of the following is not a likely adverse effect of niacin when used in the management of hyperlipidaemia?

- a. Flushing

- b. Hypoglycaemia
 - c. Hyperuricaemia
 - d. Hepatotoxicity
 - e. Gastrointestinal distress
36. For which conditions would angiotensin receptor blockers be contraindicated?
- a. Anuric renal failure
 - b. Hypokalemia
 - c. Hypertension
 - d. Diabetes mellitus
 - e. Stage A heart failure
37. The following are conditions in which beta adrenergic blockers are contraindicated EXCEPT:
- a. Recent myocardial infarction
 - b. Symptomatic bradycardia
 - c. Reactive airway disease
 - d. Second degree heart block
 - e. Severe hypotension
38. The following drugs may be used in the management of bulimia nervosa EXCEPT:
- a. Orlistat
 - b. Amitriptylline
 - c. Trazodone
 - d. Lithium

d. Reduction in both morbidity and mortality

e. Improvement in symptoms

52. Concerning viral hepatitis, which of the following pairs of viruses DO NOT commonly cause chronic disease?

a. HAV and HEV

b. HBV and HCV

c. HBV and HDV

d. HCV and HEV

e. HAV and HCV

53. A 3-year-old boy was evaluated and found to have attention deficit hyperactivity disorder. What would be the first line pharmacologic agent to be used?

a. Risperidone

b. Paroxetine

c. Methylphenidate

d. Phenytoin

e. Sibutramine

54. In the management of type IIb combined hyperlipidaemia:

a. Niacin is less effective and should be avoided

b. Statins are the most effective and should be used alone

c. Fibrates are not effective and hence their use is contraindicated

d. Dietary intervention is the treatment of choice

e. The use of bile acid sequestrants alone should be avoided

55. Which of the following statements is true concerning the use of pramlintide?

- a. It is an effective oral antidiabetic drug
- b. When used in the management of type 1 diabetes mellitus, it is given as an injection mixed together with insulin
- c. When added to insulin therapy, the dosage of insulin should be reduced by up to 50%
- d. It is normally administered at bed time to prevent morning hyperglycaemia
- e. It can be used instead of insulin in diabetic patients

56. Which of the following statements is CORRECT concerning the use of pramlintide?

- a. It is an effective oral antidiabetic drug
- b. When used in the management of type 1 diabetes mellitus, it is given as an injection mixed together with insulin
- c. When added to insulin therapy, the dosage of insulin should be reduced by up to 50%
- d. It is normally administered at bed time to prevent morning hyperglycaemia
- e. It can be used instead of insulin in diabetic patients

57. When designing empiric insulin regimen:

- a. The basal insulin should constitute two thirds of the total daily insulin requirement
- b. The prandial insulin should constitute three quarters of the total insulin requirement and the rest be given as basal insulin
- c. The basal insulin should make up half of the total daily insulin requirement
- d. More of the prandial insulin should be administered before the evening meal
- e. The ratio of basal:prandial insulin doses should be 2:1 daily

INSTRUCTIONS:-

- i. THERE ARE TWO (2) QUESTIONS IN THIS SECTION.
- ii. ANSWER ALL OF THE MEQs.
- iii. START EACH QUESTION ON A FRESH SHEET OF PAPER OF THE BOOKLET PROVIDED.

MEQ 1

B.N is a 67 year old known diabetic male patient admitted at JOOTRH with generalized body pain and oedema of the lower limbs. He was referred from Bondo District Hospital, where he was diagnosed with uncontrolled hypertension. Laboratory tests done indicate elevations in the serum potassium, BUN, creatinine, phosphorous, sodium and the pH (acidosis). B.N has a medication history of Diclofenac/Ibuprofen tablets, which he takes on and off whenever he is in pain and Enalapril for his hypertension since the year 2005.

- a) Design an optimal therapeutic plan for managing B.N, indicating the goal of therapy, THREE (3) drug types and TWO (2) therapeutic outcome monitoring. (6 marks)
- b) Name FOUR (4) risk factors that B.N exhibits that are associated with the limb oedema. (2 marks)
- c) What nondrug therapies would you recommend in B.N's therapeutic plan? (2 marks)

MEQ 2

S.M., a 25-year-old, single, unemployed man, who was brought into the emergency department by the neighbours after he was found running barefoot in the estate where he stays. He was extremely frightened, agitated and threatened the health care staff when asked questions regarding his illness. He had no history of a previous psychiatric illness or hospitalization.

Mental Status Examination revealed a thin, disheveled-looking man with very poor hygiene, who appeared much older than his stated age, extremely suspicious of the interviewer and his

surroundings, very anxious and worried. His remote memory seemed to be intact, although his immediate and short-term memories were difficult to assess because he was uncooperative. He was oriented to person, place, time, and situation. A provisional diagnosis of Schizophrenia, Paranoid Type was made.

- a) State any TWO (2) initial goals of therapy in this patient (2 Marks)
- b) Explain the initial management of this patient, stating any TWO (2) drugs that can be used and their appropriate routes of administration and conditions for their use. (2 Marks)
- c) Give the pharmacotherapeutic options available for the management of this patient's condition, by explaining any FOUR(4) drug classes that can be used and their roles in this patient or in the course of the patient's management (4 marks)
- d) List any FOUR (4) non pharmacologic management that can be used to manage this patient (2 Marks)



MASENO UNIVERSITY

UNIVERSITY EXAMINATIONS 2017/2018

**FOURTH YEAR END YEAR EXAMINATION FOR THE DEGREE OF
BACHELOR OF MEDICINE AND BACHELOR OF SURGERY WITH
INFORMATION TECHNOLOGY**

CITY CAMPUS

SUPPLEMENTARY EXAMINATION

MCS 405: JUNIOR CLERKSHIP IN PSYCHIATRICS

Date: 23rd May, 2018

Time: 9.00 - 12.00pm

INSTRUCTIONS:

- As per the attached.



INSTRUCTIONS

(i) EACH OF THE QUESTIONS IN THIS SECTION HAS A STEM STATEMENT FOLLOWED BY FIVE COMPLETIONS NUMBERED (a) TO (e). A GIVEN COMPLETION WHEN ADDED TO THE STEM STATEMENT MAY PRODUCE A COMPLETE STATEMENT THAT IS TRUE (T) OR FALSE (F)

(ii) INDICATE AGAINST THE APPROPRIATE NUMBERS IN THE ANSWER SHEET PROVIDED WHETHER THE COMPLETE STATEMENT IS TRUE (T) OR FALSE BY PRINTING AN 'X' IN THE BOX IN THE COLUMN HEADED 'T' FOR TRUE OR 'F' FOR FALSE.

(iii) IF YOU DO NOT KNOW WHETHER THE COMPLETE STATEMENT IS TRUE OR FALSE, PRINT THE 'X' IN THE COLUMN CLEARLY LABELED (FOR DO NOT KNOW).

(vi) FOR EACH ITEM CORRECTLY COMPLETED ONE MARK WILL BE AWARDED. FOR AN ITEM INCORRECTLY COMPLETED NO MARK WILL BE LOST. FOR EACH ITEM COMPLETED AND LABELLED 'D' NO MARK WILL BE GAINED OR LOST. ANY UNANSWERED OPTION COUNTS AS WRONG

(v) THERE IS NO RESTRICTIONS ON THE NUMBER OF TRUE OR FALSE ITEMS IN A QUESTION. IT IS POSSIBLE FOR ALL THE ITEMS IN A QUESTION TO BE TRUE OR FOR ALL TO BE FALSE

SECTION 1: MULTIPLE CHOICE QUESTIONS (MCQs) (60 minutes)

1. Uncomplicated bereavement is **NOT** characterized by:

- a- Anger
- b- Depressive mood for one month
- c- Feelings of worthlessness
- d- Sensations of seeing or hearing the deceased
- e- Transient guilt

2. **The Mental Health Act, 1989**, provides for the following modes of admissions:

- a- Voluntary admission
- b- Military admission
- c- Emergency admission
- d- Doctor's recommendation
- e- Involuntary admission

3. **Exemption from criminal responsibility** may be considered under the following circumstances:

- a- A mistake has occurred
- b- An accident
- c- Duress
- d- Necessity
- e- Insanity

4. The following are **common causes of mental handicap**:

- a- Toxins
- b- CPD
- c- Prematurity
- d- Diarrhea and Vomiting
- e- Chromosomal abnormalities

5. **Affective Disorder**:

- a- Are disorders of mood
- b- Occur in psychotic forms only
- c- Are commonly seen in outpatient clinics
- d- Comprise depression, mania and Bipolar disorders
- e- Can be explained by the monoamine theory

6. Maladaptive Behaviours

- a- Represent an exaggeration of normal behaviour
- b- Increases morbidity and mortality
- c- Are a response to anxiety
- d- A sense of subjective compulsion is experienced
- e- Alcohol dependence is one example

7. Which of the following suitably describes "Mental Retardation"?

- a- Axis I
- b- Axis II
- c- Axis III
- d- Axis IV
- e- Axis V

8. Agoraphobia is characterized by the following:

- a- it is fear of open space
- b- It is caused by trauma
- c- It is treated by Beta-Blockers
- d- It occurs only in open space
- e- It may cause a House-Bound Syndrome

9. The following is/are a bad prognostic feature of Schizophrenia?

- a. late onset of the illness
- b. acute onset of symptoms
- c. high prevalence of family history of mood disorders
- d. presence of negative symptoms
- e. Obvious precipitating factors

10. Electroencephalography may be used in the diagnosis of:

- a. Schizophrenia
- b. Epilepsy
- c. Conversion disorder
- d. Major depressive disorder
- e. Somatoform pain disorder

11. A patient presents with involuntary frowning, blinking grimacing, and choreoathetoid movements of the upper extremities after several years of antipsychotic medication treatment. Which one of the following is the **MOST likely diagnosis?**

- a. Akinesia
- b. Akathisia
- c. Acute dystonia
- d. Parkinsonism
- e. Tardive dyskinesia

12. The feature(s) of mental retardation include:
- a- decreased motivation and response
 - b- slow movement
 - c- slow interaction
 - d- missed developmental milestones
 - e- all of the above is
13. One of the following is a Tricyclic Antidepressant:
- a- Amitriptyline
 - b- Sertraline
 - c- Phenezine
 - d- Mirtazapine
 - e- Fluoxetine
14. If a pt cries for one minute and laughs for another in response to non-existing stimuli. This is called:
- a- expansive mood
 - b- hallucination
 - c- labile affect
 - d- inappropriate affect
 - e- incongruous mood
15. Long-term treatment of Bipolar I patients include:
- a- mood stabilizers and prophylaxis
 - b- education of a patient and his/her family
 - c- increasing support and decreasing stressors
 - d- electroconvulsive therapy
 - e- all of the above
16. Regarding the CAGE questionnaire, which of the following statements is **TRUE**?
- a. It is a screening questionnaire
 - b. It asks whether a person feels like cutting back on his or her drinking
 - c. There is a question on whether a person feels guilty about his or her drinking
 - d. It asks about having a drink early in the morning as an 'eye-opener'
 - e. All of the above
17. The following is a pathognomonic feature of Schizophrenia (if found, the Diagnosis is sure)
- a- deterioration of functions
 - b- 3rd voice talking to a patient
 - c- delusion of control
 - d- all of the above
 - e- None of the above

18. The following have been associated with neuroleptic malignant syndrome EXCEPT:

- a- Nortriptyline
- b- Olanzapine
- c- Organic brain disease
- d- Haloperidol
- e- Fluphenazine

19. Regarding conversion disorders which is FALSE

- a- They are defined as a loss or change of function suggesting a physical disorder but caused by a psychological conflict
- b- 90% of the time present as neurological symptoms/disorders
- c- Often the patient seems relatively unconcerned about the symptom
- d- The symptom production is a conscious one
- e- All above are true

20. The following are/is a characteristic feature of a manic episode?

- a- Elevated, expansive, irritable mood
- b- Increased psychomotor activity
- c- Thought echo, thought insertion or thought withdrawal
- d- Flight of ideas
- e- may incur "debts" all over

21. The following condition is classified under personality disorders in ICD-10 classification?

- a- Paranoid
- b- Schizoid
- c- Schizotypal
- d- Dissocial
- e- Borderline

22. Exemption from criminal responsibility may be considered under the Following Circumstances:

- a- A mistake has occurred
- b- An accident
- c- Duress
- d- Necessity
- e- Insanity

23. The following are common causes of mental handicap:

- a-. Toxins
- b-. CPD
- c-. Prematurity
- d-. Diarrhea and Vomiting
- e-. Chromosomal abnormalities

24. These are **First rank symptoms** of Schizophrenia:

- 60. Audible Thoughts
- b. Thought Broadcasting
- c. Voice arguing or discussing or both
- d. Perplexity
- e. Somatic passivity experiences

25. **Atypical antipsychotic** drug(s) include

- a. Phenothiazines
- b. Butyrophenones
- c. Olanzapine
- d. Fluphenazine
- e. Largactil

26. In Teenagers, Schizophrenia can present with the following complaints:

- a. Withdrawal from friends and family
- b. A drop in performance at school
- c. Trouble sleeping
- d. Irritability or depressed mood
- e. Lack of motivation

27. The underlying principles of a therapeutic community **include**:

- a. Democracy
- b. Reality confrontation
- c. Permissiveness
- d. Community
- e. All of the above

28. The following are **NOT** features of puerperal psychosis:

- a. Onset is gradual, in the first few months after childbirth
- b. It is associated with tearfulness
- c. The child's life may be in danger
- d. It is more common in primiparae
- e. There is an increased occurrence in poor spousal support

29.. . A **delusion** is:

- a. A perception in the absence of a stimulus
- b. The misinterpretation of a stimulus
- c. A false but firmly held belief despite evidence to the contrary
- d. An inaccurate thought or belief based on interpretation of a stimulus
- e. A common feature in hypochondriasis

30. The following imaging techniques can be used to measure glucose metabolism in different parts of the brain?

- a. CT scan
- b. MR scan
- c. PET scan
- d. SPECT scan
- e. None of the above

SECTION II - SHORT ANSWER QUESTIONS (SAQ) (60 minutes)

INSTRUCTIONS

- (i) **THERE ARE 7 TOPICS IN THIS SECTION**
- (ii) **WRITE SHORT NOTES ON THE TOPICS OF YOUR CHOICE**
- (iii) **RESPOND TO ONLY FOUR TOPICS**
- (iv) **USE A SEPARATE BOOKLET FOR EACH TOPIC**
- (v) **EACH TOPIC ANSWERED CARRIES 15 MARKS TO BE ANSWERED IN 60 MINUTES.**

- 1. Neuroleptic Malignant Syndrome (NMS)
- 2. Tardive Dyskinesia
- 3. 'Cheese reaction'
- 4. Catastrophic Reaction
- 5. Normal Grief
- 6. "Cold turkey"
- 7. Multiaxial Diagnostic Formulation
- 8. Avolition Disorder

SECTION III - LONG ESSAY QUESTIONS (LEQ) (40 minutes)

INSTRUCTIONS:

- (i) **THERE ARE TWO QUESTIONS IN THIS SECTION**
- (ii) **ANSWER ONLY ONE QUESTION**
- (iii) **THE QUESTION TO BE ANSWERED IN 40 MINUTES**

- LEQ 1 **Discuss the (a) Aetiological factors for mental disorders**
- LEQ 2 **Discuss Mental Status Assessment in Psychiatry**



MASENO UNIVERSITY

UNIVERSITY EXAMINATIONS 2017/2018

**FIFTH YEAR END YEAR EXAMINATION FOR THE DEGREE OF
BACHELOR OF MEDICINE AND BACHELOR OF SURGERY WITH
INFORMATION TECHNOLOGY**

CITY CAMPUS

SUPPLEMENTARY EXAMINATION

MCS 406: JUNIOR CLERKSHIP IN SURGERY

Date: 23rd May, 2018

Time: 9.00 - 12.00pm

INSTRUCTIONS:

- This paper contains One hundred (100) multiple choice questions each with options A – E.
- Choose ONE single best answer by circling the correct option
- Answer ALL Questions
- Write your admission number on ALL pages



