Critical Care Guidelines FOR CRITICAL CARE USE ONLY



6. PosiMitt Protocol

На		
1.	RASS	YES
2.	CAM-ICU	YES
3.	Delirium Tool	YES
4.	Management of Agitation	YES 🔲

vialing efficit of Agriculon				
	Υ	N	Comments	
Do we need to re consider staff to patient ratio?				
Is patient a danger to themselves despite appropriate sedation and management?				
Do risks of over sedation outweigh risks of mechanical restraint What is the main risk for the patient? (please specify) Have you considered all other possible aids to avoid that risk (see assessment)? Is PosiMitt appropriate?				
is Posiviiti appropriate?				
Has capacity been assessed? Complete and date Adult Incapacity certificate (Scotland Act,2000, section 47) If restraint beyond mittens is required, document "appropriate restraint to facilitate treatment".				
Is the appropriate relative or carer aware of need for mechanical restraint and provided with relevant information?				
Decision made by medical staff and nurse in charge				
Signed senior medical staff				
Signed senior nursing staff				
Date				

REMEMBER

Evaluate effectiveness after 1 hour and every 2 hours thereafter until restraints assessed to be no longer necessary

Title: 6. PossiMitt protocol			
Authors: Michelle Jack, Alasdair Hay	Approved by: QIT editorial group		
Status Final:	Written: June 2015		
Reviewed on: May 2019	Next Review: May 2021		