


## Critical Care Guidelines for the emergency management of acute upper GI bleeding and Oesophago-Gastro-Duodenoscopy (OGD)

### 1. Urgent/emergency endoscopy for Acute Upper GI Bleeding



**i) Patients most likely to benefit from IR for haemorrhage control**

- Non-variceal haemorrhage (NB. TIPSS may be indicated post banding of OV)
- Failed endoscopic management
- Known lesion, potentially amenable to IR intervention
- Patient is stable enough to transfer to IR (May require GA and anaesthesia assistance)
- Risk of need for laparotomy is low (see below)
- Significant co-morbidities, which may preclude further OGD/surgery

**ii) Patients at high risk of need for laparotomy (OGD must be performed in theatre)**

- Uncontrolled non-variceal haemorrhage, patient too unstable for transfer to IR
- Failed endoscopic management and IR contraindicated/bleeding point not amenable to IR intervention
- Suspected perforation (de novo or iatrogenic), surgical abdomen, bleeding GIST
- GI consultant considers risk to be high

**iii) Urgent OGD in Critical Care**

OGD for oesophageal varices (or other acute GI haemorrhage, where the risk of need for laparotomy, is low), **may** be performed in 118 within dayshift hours, in selected patients, e.g. if the patient is already a critical care inpatient, intubated and staffing levels/experience levels allow. **This is at the discretion of the duty Critical Care Consultant, who must be involved in discussions.**

**iv) Admission to critical care, prior to OGD**

Patients who require urgent, but not emergency endoscopy, may benefit from admission to Critical Care pre-endoscopy, for resuscitation and transfusion. There is evidence that, this may improve outcome (2,3).

**Further management of acute upper GI haemorrhage**

- There is no role for Tranexamic acid -see HALT-It study (4)
- Adopt a restrictive post-emergency resuscitation blood transfusion strategy, that aims for a target haemoglobin between 7g/dL and 9g/dL. A higher target haemoglobin, should be considered in patients with significant co-morbidity e.g. ischaemic cardiovascular disease (1,2,6)
- PPI infusions should be administered for 72 hours post endoscopy (unless directed otherwise, by senior GI physicians) as per the Hong Kong protocol (5)
- Consider correction of coagulopathy and reversal of antiplatelets -may require specialist input from haematology (1,2,6)

## 2. Non-urgent endoscopy for Critical Care inpatients

Non-urgent OGD can be performed in a level 3 Critical Care area (pods 118A, 118B and 116D) for patients already in ICU, requiring procedural interventions e.g. NJ tube insertion, providing the following conditions are met:

- The ICU base duty Critical Care consultant and Charge nurse, are in agreement.
- Medical and nursing staffing levels/experience are adequate for the procedure to be performed safely.
- The procedure is performed in dayshift hours.

OGD should not be performed in a level 2 Critical Care area, i.e. 116C

## References

1. NICE guideline: “Acute upper gastrointestinal bleeding in over 16s: management” Clinical guideline [CG141] Published: 13 June 2012 Last updated: 25 August 2016. <https://www.nice.org.uk/guidance/cg141>
2. “British Society of Gastroenterology (BSG)-led multi-society consensus care bundle for the early clinical management of acute upper gastrointestinal bleeding”. 2019 <https://www.bsg.org.uk/clinical-resource/bsge-acute-upper-gi-bleed-care-bundle/>
3. Guo CLT, Wong SH, Lau LHS, *et al.* “Timing of endoscopy for acute upper gastrointestinal bleeding: a territory-wide cohort study”. *Gut* 2022;**71**:1544-1550. <https://gut.bmj.com/content/gutjnl/71/8/1544.full.pdf>
4. HALT-It trial. “Effects of a high-dose 24-h infusion of tranexamic acid on death and thromboembolic events in patients with acute gastrointestinal bleeding (HALT-IT): an international randomised, double-blind, placebo-controlled trial”. 2020 [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(20\)30848-5/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)30848-5/fulltext)
5. Hong Kong protocol [https://www.cag-acg.org/images/publications/Intl\\_consensus\\_recommendations\\_on\\_managmen\\_%20of\\_UGIB\\_2010.pdf](https://www.cag-acg.org/images/publications/Intl_consensus_recommendations_on_managmen_%20of_UGIB_2010.pdf)
6. “Diagnosis and management of nonvariceal upper gastrointestinal haemorrhage: European Society of Gastrointestinal Endoscopy (ESGE) Guideline”; 2015 <https://pubmed.ncbi.nlm.nih.gov/26417980/>

<b>Title:</b> Critical Care Guidelines for Oesophago-Gastro-Duodenoscopy (OGD)	
<b>ID</b>	<b>Authors:</b> M Beatty, N Church
<b>Category</b>	<b>Document Version:</b> 4
<b>Status:</b> Final	<b>Review Date:</b> October 2025
<b>Authoriser:</b> RIE QIT team	<b>Date Authorisation:</b> September 22nd 2022
<b>Date added to intranet</b>	
<b>Key words:</b> Endoscopy, OGD, Acute upper GI haemorrhage	