

FOR ICU USE ONLY

ADRENALINE

PRESENTATION:	Ampoules containing 1mg/ml of adrenaline.
INDICATION:	Inotropic support with variable degree of vasoconstriction and vasodilation. Bronchospasm and anaphylaxis.

DOSE AND ADMINISTRATION: ICU STANDARD INTRAVENOUS INFUSION

The infusion prepared depends on the dose and infusion rate. See table below.

Strength	Adrenaline (mg)	Glucose 5% (ml)	Comments
Single Strength Final concentration 80 micrograms/ml	8mg 20mg 40mg	100ml 250ml 500ml	Remove volume of glucose 5% from bag equivalent to volume of adrenaline which is to be added
Double strength Final concentration 160 micrograms/ml	40mg 80mg	250ml 500ml	
Quadruple strength Final concentration 320 micrograms/ml	80mg 160mg	250ml 500ml	

Initially 1-7mls/hour (0.01-0.1 micrograms/kg/min) of single strength titrated to required level.

Infuse through a central venous catheter.

Single Strength may be infused via a large peripheral vein until central venous access is established. Closely monitor the insertion site for the development of phlebitis.

CONCENTRATION:	80micrograms/ml 160micrograms/ml 320micrograms/ml
-----------------------	---

STABILITY:	Physically and chemically stable for 24 hours at room temperature. Protect from light.
-------------------	--

ADDITIONAL NOTES:	Also stable for 24 hours in 0.9% sodium chloride but more stable in glucose 5%. Do not use if discoloured or has a precipitate.
--------------------------	---

References

1. Trissel's IV Compatibility- Adrenaline. Accessed 06/03/19 via Micromedex 2.0, www.micromedexsolutions.com
2. Peddicord TE, Olsen KM, Zumbunnen TL, Warner DJ, Webb L. Stability of high-concentration dopamine hydrochloride, norepinephrine hydrochloride, and nitroglycerin in 5% dextrose injection. Am J Health-Syst Pharm. 1997; 54:1417-9.
3. Brustugun J, Kristensen S, Tonnesen HH. Photosensitivity of Sympathomimetic Agents in Commonly Used Infusion Media in the Absence and Presence of Bisulfite. PDA Journal of Pharmaceutical Science and Technology. 2004;58 (6):296-308.
4. Minimum Infusion Volumes for fluid restricted critically ill patients. Critical Care Group, UKCPA V 4.4, Dec 2012
5. BNF- adrenaline/epinephrine. Accessed 05/08/19 via www.bnf.nice.org.uk
6. Medusa IV guide –adrenaline/epinephrine. Accessed 07/08/19 via <https://medusa.wales.nhs.uk/IVGuideDisplay.asp>

Title: ADRENALINE	
ID:	Authors: Anne Neally
Category:	Document Version: 2.0
Status Draft/Final:Final	Review Date: August 2019
Authoriser: Lothian Critical Care QIT editorial group	Authorisation Date: August 2021
Date added to Intranet:18/10/19	

FOR ICU USE ONLY

FOR ICU USE ONLY

Key Words/comments	
---------------------------	--

FOR ICU USE ONLY