
Inhaled Nitric Oxide (iNO) for severe hypoxaemia

- For unselected patients with ARDS iNO is associated with transient improvements in oxygenation but does not carry a mortality benefit and increases renal dysfunction.
- The use of iNO for hypoxaemic patients with ARDS should be severely limited to
 those in whom other rescue therapies are not possible but are believed to have a
 recoverable disease process eg severe hypoxaemia while awaiting ECMO retrieval –
 in consultation with ECMO referral centre with consideration as to whether in-house
 ECMO is necessary/possible if prolonged retrieval anticipated. The use of iNO
 should ALWAYS be a consultant led decision.
- Patients in this category would typically have a Pa02 of < 8 KPa with FIO2 > 0.9 and PEEP > 10 following consideration of prone ventilation, neuromuscular blocking drugs and ECMO.

Suggested titration of iNO for severe hypoxaemia:



- Rebound pulmonary hypertension can occur on withdrawal of iNO therapy with hypoxaemia or haemodynamic compromise.
- iNO may also be used under specialist advice in cases of right heart failure and pulmonary hypertension and falls out-with this guidance.
- Methaemoglobin levels should be monitored during therapy with iNO at 0 hours, 1 hour, 6 hours then daily, and iNO is contra-indicated in patients with congenital or acquired methaemoglobin reductase deficiency.
- Relative contra-indications to iNO therapy include severe coagulopathy, intracranial haemorrhage and severe left ventricular failure.

| Title: inhaled nitric oxide for hypoxaemia | |
|--|--------------------------------------|
| ID: iNO2020 | Authors: N Young – adapted from L |
| | Ferguson and I DeBeaux 2013 guidance |
| Category: Respiratory | Document Version: 2.0 |
| Status Draft/Final: Draft | Review Date: April 2022 |
| Authoriser: G McNeill | Date Authorised: April 2020 |
| Date added to Intranet: March 2020 | |
| Key Words: iNO; respiratory failure; ARDS | |
| Comments: | |