

Management of Acute Type B Aortic Dissections Guideline

Haemodynamic targets (initial)

- Systolic BP 100-120 mmHg
- MAP <80 mmHg
- Targets should be changed **ONLY** after consultation with Vascular team

If patient develops leg weakness, the Vascular surgeon and Vascular anaesthetist must be contacted immediately. **Potential** interventions for spinal cord ischaemia

- Increasing target BP to avoid potential spinal cord infarction
- Emergency CSF drain
- Repeat CT or MRI imaging

Early medical management: Aggressive BP control, analgesia and anti-emetics

Analgesia

- **Morphine** (1-10mg) IV titrated to effect

Then

- **Morphine PCA** 1mg bolus 5 min lockout

*If the patient has **renal impairment**, morphine can be replaced with **fentanyl** 10 microgram bolus 5 min lockout*

- Regular **Paracetamol** (unless contra-indications)

Anti-emetics

- **Ondansetron** 4mg IV every 8 hours
- Supplemental cyclizine 50mg IV every 8 hours and metoclopramide 10mg IV every 8 hours may be used

BP control

Intravenous therapy

1. **Labetalol** (first choice)
 - a. Administer IV bolus injections for initial control of blood pressure (10mg slow IV bolus injections at 2 minute intervals to a maximum of 200mg per course of boluses).
 - b. AND ALSO start an IV infusion to maintain blood pressure control.
 - i. Concentration 5mg/ml for CVC use OR 1mg/ml for PVC use
 - ii. Dose – Start at 15mg/hr and titrate to clinical effect, but often 10-60mg/hour.
2. **Nicardipine** (second line in addition to labetalol, or first line if contra-indications to beta-blocker)
 - a. IV infusion (change IV infusion site every 12h if peripherally administered)
 - i. Concentration 25mg made up to 250ml (5% glucose) = 100micrograms/ml
 - ii. Dose – titrated to clinical effect
 - iii. Start at 50ml/hour (5mg/hour). The rate may be increased every 10 mins by 25ml/hour to a maximum of 150ml/hour (15mg/hour).
 - iv. Once target BP is achieved reduce dose gradually, usual maintenance dose 2-4mg/hour

3. **Hydralazine** (third line) **NB OFF LICENCE USE – Consultant directed use only**
PATIENT MUST BE RATE CONTROLLED BEFORE COMMENCING HYDRALAZINE
 - a. IV bolus – 5mg slow IV injection bolus at 20 minute intervals to a usual maximum of 20mg
 - b. IV infusion
 - i. Concentration 60mg made up to 60ml (0.9% sodium chloride) = 1mg/ml
 - ii. Dose – titrated to clinical effect
 - iii. Start at 3ml/hr (50micrograms/min). The rate may be increased every 10 mins by 3ml/hour to a maximum of 18ml/hour (300micrograms/min).

Oral therapy – Start as soon as possible (Day 1 unless contra-indicated)

Titrate first line drug to maximum tolerated dose before introducing next line drugs

1. **Bisoprolol** (first choice)
 - a. 2.5-20mg once daily
2. **Amlodipine** (second line in addition to bisoprolol, or first line if contra-indications to beta-blocker)
 - a. 5-10mg once daily
3. **Doxazosin** (third line in addition to bisoprolol and amlodipine)
 - a. 1-16mg once daily

NB ACE Inhibitors and diuretics should be avoided initially while the kidneys are at risk.

References

1. Curran MP, Robinson DM, Keating GM. Intravenous nicardipine: its use in the short-term treatment of hypertension and various other indications. *Drugs*. 2006;66(13):1755–82.
2. Dade J et al. UKCPA Critical Care Group. Minimum infusion volumes for fluid restricted critically ill patients. 4th Ed. V4 2012. Accessed 4.2.15 via <http://www.ukcpa.net/groups/critical-care/documents/?category=71>
3. Summary of Product Characteristics. Nicardipine 10mg/10ml solution for Injection. Concordia International. www.medicines.org.uk. Accessed 12/12/17.
4. Summary of Product Characteristics. Hydralazine 20mg Powder for Concentrate for Solution for Injection/Infusion. Concordia International. www.medicines.org.uk. Accessed 12/12/17.
5. Summary of Product Characteristics. Labetalol Hydrochloride 5mg/ml solution for injection. Concordia International. www.medicines.org.uk. Accessed 02/02/2018.
6. Hydralazine Hydrochloride. The UK Injectable Medicines Guide. Accessed 09/04/18.

Title: Management of Acute Type B Aortic Dissection Guideline	
ID: MATBADGv5.20190702	Authors: O Falah C Moores E McGregor M Dunn C Hannah
Category:	Document Version: 5
Status Draft/Final: Final	Review Date: July 2022
Authoriser: QIT, Vascular Surgery & Anaesthesia	Date Authorisation: July 2019
Date added to Intranet: Oct 2015	
Key words: Aortic Dissection	

The use of this guideline is subject to professional judgement and accountability. It should not be interpreted as setting a standard of care. This guideline has been prepared carefully and in good faith for use within the Directorate of Critical Care at NHS Lothian. No liability can be accepted by NHS Lothian for any errors, costs or losses arising from the use of this guideline or the information contained herein

Think DISSECTION

The Management of Acute Type B Aortic Dissections

Maaz Syed, ST Vascular Surgery
Orwa Falah, Consultant Vascular Surgery
Mark Dunn, Consultant Critical Care
Euan McGregor, Consultant Anaesthetics
Alistair Nimmo, Consultant Anaesthetics



1 RESUSCITATION


ABCDE

Early anticipation of critical care involvement

4 ANALGESIA AND ANTI-EMETICS

Morphine 1 – 10mg IV titrated to effect;
Morphine PCA 1mg bolus; 5min lock out;
Fentanyl 10µg IV bolus 5min lock out [renal failure]
Paracetamol 1g QDS (unless contraindicated)
Ondansetron 4mg IV every 8 hours OR
Cyclizine 50mg IV PRN 8 hourly OR
Metoclopramide 10mg IV PRN 8 hourly


2 PROMPT DIAGNOSIS



Hypotension may be due to myocardial ischaemia, tamponade, aortic root incompetence, or an aortic bleed. These warrant urgent investigations.



3 SYSTEMIC MONITORING



TYPE A

Urgent Cardiothoracic Opinion [bleep: 1682]

Urgent Vascular Surgery [via switchboard] and **Critical Care** [bleep: 2306] **Opinion**

TYPE B

6 LONG-TERM MANAGEMENT

CONVERT TO ORAL BLOOD PRESSURE CONTROL AS EARLY AS TOLERATED

Target BP 120/80 mmHg
 Repeat CT before discharge (usually at 48 hours)
 Follow-up CT at 1, 6 and 12 months.
 Outpatient follow-up at 8 weeks post discharge, unless indicated sooner.

- Bisoprolol** 2.5 – 20mg once daily
- Amlodipine** 5 – 10mg once daily
- Doxazosin** 1 – 16mg once daily

HIGH RISK FEATURES OF TYPE B DISSECTIONS !

Visceral / limb ischaemia
 Entry tear $\geq 10\text{mm}$
 Inner curve entry tear
 Aortic diameter $\geq 4\text{cm}$
 On-going Pain or HTN

Retrograde dissection
 False lumen (FL) $\geq 22\text{mm}$
 Partial FL thrombosis
 Fusiform index ≥ 0.64
 Grow $\geq 1\text{cm/yr}$ or $\geq 5.5\text{cm}$