

# Nasogastric Enteral Feeding Protocol

When NG tube position is confirmed (Refer to Critical Care Guideline)  
Commence **Nutrison 1.0 kcal/ml at 25mls/hr**

- Check aspirate after 4 hours
- Is aspirate > 250mls?

YES

NO

- Return 100mls
- Feed rate 25 mls/hr

- Return 100mls
- Increase to 50mls/hr

- Check aspirate after 4 hours
- Is aspirate > 250mls?

- Check aspirate after 4 hours
- Is aspirate > 250mls?

NO

YES

YES

NO

- Return 100mls
- Reduce rate to 10mls/hr

- Return 100mls
- Increase rate to 75mls/hr x 24 hours
- Aspirate 4 hourly until review by dietitian

Commence prokinetic(s)  
Metoclopramide  
and/or Erythromycin

- Once established on enteral feeding, consider changing to fine bore NG tube (Medicina 10Fr – 30 days)

If NG feed not tolerated within 48 hrs of commencing prokinetic consider post-pyloric feeding (NJ) and/or parenteral nutrition

<b>Title: NASOGASTRIC ENTERAL FEEDING PROTOCOL</b>	
<b>Status Draft/Final: FINAL</b>	<b>Authors: J MERRIWEATHER</b>
	<b>Approved by: QIT editorial group</b>
	<b>Written: AUG 2015</b>
<b>Reviewed on:</b>	<b>Next review : AUG 2018</b>