

Anidulafungin

ONLY TO BE USED WITH AGREEMENT OF ICU CONSULTANT AND CONSULTANT MICROBIOLOGIST

PRESENTATION: Vials containing 100mg of powder for reconstitution.

INDICATION: Antifungal.

DOSE AND ADMINISTRATION: 200mg dose on first day, then 100mg once daily.

ICU STANDARD INTRAVENOUS INFUSION

Reconstitute 100mg vial with 30 ml water for injections. Reconstitution may take up to 5 minutes. Further dilute the reconstituted solution with glucose 5% as follows:

Dose	Total volume of reconstituted solution	Infusion volume	Total infusion volume
100mg	30ml	100ml	130ml
200mg	60ml	200ml	260ml

Infuse 100mg over 90 minutes.
Infuse 200mg over 180 minutes.

CONCENTRATION: Must be diluted to a concentration of 0.77mg/ml.

STABILITY: Physically and chemically stable for 24 hours at room temperature.

ADDITIONAL INFORMATION: Anidulafungin is stable in sodium chloride 0.9%

References

1. ECALTA 100mg powder for concentrate for solution for infusion. Pfizer Ltd. Summary of Product Characteristics.2017. <https://www.medicines.org.uk/emc/product/454> (Accessed 15/08/2018)
(Revised 10/18)

Title: ANIDULAFUNGIN	
ID:	Authors: Hannah Prescott
Category:	Document Version: 2.0
Status Draft/Final: FINAL	Review Date: December 2021
Authoriser:	Authorisation Date: March 2019
Date added to Intranet	
Key Words	
Comments	