


Upper GI Bleeding – Urgent Endoscopy & Patient Monitoring

Aim: rapid, safe and appropriate access to Critical Care for urgent/emergency endoscopy and continued monitoring and care.

Referral to Critical Care by GI Registrar / Consultant



Indications for endoscopy on CEPD list in theatre 17 (TEL 23241)

- Urgent UGIE and no readily available bed in Critical Care
- Planned/expected need for GA (NB varices should go to 118)
- High risk for laparotomy (e.g. recurrent bleed in patient with PUD)
- Difficult airway

If conflict over priority for theatre arises, the Consultant Anaesthetist on call for emergencies should be the final arbiter. This may involve opening a second theatre.

Patient disposal from theatre- high-risk re-bleeders go to critical care (most appropriate ward) and non high-risk to ward 107 or 205. It is clearly not always possible to identify a specific bed/ward before endoscopy and this should not delay patients being brought to theatre.

UGIE under sedation in Ward 116 must *always* be discussed with the HDU Consultant from 08:00 to 18:00 hrs and Anaesthetic SpR Page 2200 from 18:00 to 08:00hrs (if unavailable call ICU Registrar page 2306) **prior to commencing procedure.**

Title: Upper GI Bleeding – Urgent GI Endoscopy & Patient Monitoring	
ID: CCUGIB010712v2	Authors: Dr M. Beatty, Dr N. Church & Dr B. Cook
Category: 1	Document Version: 2
Status Draft/Final: Final	Review Date: 01.07.14
Authoriser: QIT Editorial Board	Date Authorisation: 01.07.12
Date added to intranet	31.07.12
Key Words: Critical care, upper GI bleeding, endoscopy	