

Rocuronium

| | |
|---------------------------------|--|
| PRESENTATION: | Ampoules containing 100mg/10ml and 50mg/5ml. |
| INDICATION: | <p>Second line treatment for muscle relaxation in ICU patients when atracurium supply not available or not suitable.</p> <p>Rocuronium is a rapid onset, intermediate non-depolarising neuromuscular blocking drug.</p> |
| DOSE AND ADMINISTRATION: | <p>ICU STANDARD INTRAVENOUS INFUSION</p> <p>IV loading dose: 0.6 mg/kg by rapid injection.</p> <p>IV infusion: Administered undiluted e.g. 500mg in a 50ml syringe.</p> <p>Usual maintenance doses are between 0.3 - 0.6 mg/kg/hr. For a 70kg patient, this would equate to 2.1-4.2ml/h.</p> |
| CONCENTRATION: | 10mg/1ml. |
| MONITORING: | Monitoring of neuromuscular function using train-of-four peripheral nerve stimulation, is recommended during the use of rocuronium in order to individualise dosage requirements. |
| STABILITY: | Physically and chemically stable for 24 hours at 25° C. |
| OTHER INFORMATION: | Rocuronium has a low pH and may cause venous irritation and tissue damage in cases of extravasation. If a CVC is unavailable, administer via a large peripheral vein and monitor closely. |

References

1. Rocuronium bromide 100mg/10ml injection, 2013. Bowmed Ibisqus Limited. Summary of Product Characteristics. Available from: <https://www.medicines.org.uk/emc/product/8758/smpc> (Revision of text 28/12/23)
2. United Kingdom Clinical Pharmacy Association, 2012. Minimum Infusion Volumes. Fourth Edition.
3. Rocuronium Bromide, Injectable Medicines Guide, Medusa. Updated: April 2024. Accessed: May 2024

| | |
|---|--|
| Title: Rocuronium | |
| ID: | Authors: Mairi Cromarty, Fiona Clarke, Euan Simcox |
| Category: | Document Version: 2.0 |
| Status Draft/Final: Final | Review Date: May 2026 |
| Authoriser: Lothian Critical Care QIT Editorial Group | Authorisation Date: March 2022 |
| Date added to Intranet: March 2022 | Updated: May 2024 |
| Key Words | |
| Comments | |