NHS Lothian Critical Care CoVid Intubation Checklist

R Baruah, M Blacstock (Adapted from DAS/ICS Intubation Checklist 2018)
Review Date: March 2020



- Covid RSI box and trolley
- Intubation drugs
 Allergies, 个K+?
- Pre-oxygenate using C circuit
 Check O₂ supply
- Patient Optimisation

 i.v. access checked

 Optimise head position
- Monitoring visible
 ECG, SpO₂, BP
 ETCO₂ turned ON &
 waveform visible
- Covid PPE
- Communication
 Site mobile phone

2 Equipment

- Yankauer suction -on
 - Cut tube
 Check cuff & lubricate
 Tie or tape
 10 ml syringe
 Stylet (if requested)

ETT \times **2** (size requested)

- 2 x laryngoscope (McGrath and backup Mac))
- McGrath handle check turns on + Mac 4
- OPA (Sizes 3, 4, 5)
- Bougie
- FoNA kit in trolley
- Closed suction catheter mount
- Stethoscope

3 Team Plan

- Verbalise role allocation
- 1 Intubator
- 2 Cricoid pressure3 Intubator's assistant
- 4 Drugs administrator
- 4 Drugs administrator
- 5 Runner (outside room)
- Induction Drugs Plan
- Emergency Drugs incl. vasopressors
- On-going sedation
- Is further help required?
- Who will contact and how?ICU Cons / Theatre

Anaesthetist via site mobile phone

- Plan for Difficulty
- Can we wake this patient up if intubation fails?
- Vocalise "Airway plan is.."
- Drugs and laryngoscopy
- Plan B/C

Plan A

- Supraglottic airway
- Fibreoptic intubation via supraglottic airway
- Plan D
 - FoNA
 - Scalpel-bougie-tube
- Double check
 Understand roles?
- RAISE CONCERNS NOW
- PROCEED
- Confirm ETT tip in trachea 3cm - max 5cm from carina