

KETAMINE

For resistant status epilepticus only. For acute pain see "[guideline for ketamine subcutaneous infusion for acute pain](#)"

PRESENTATION:	Vial containing 500mg in 10ml, 50mg/ml of ketamine.																																																										
INDICATION:	Resistant status epilepticus (unlicensed use).																																																										
DOSE AND ADMINISTRATION:	<p>The evidence on the use of ketamine is mostly based on isolated case reports. From the information available:</p> <p>Intravenous loading dose of 3mg/kg. Use ideal body weight in obese patients.</p> <p>Maintenance intravenous infusion of 1-5mg/kg/hr. Use ideal body weight in obese patients.</p> <p>ICU STANDARD INFUSION</p> <p>Loading dose: 3mg/kg intravenously, over at least one minute, then,</p> <p>Continuous infusion: 2500mg in 50ml undiluted.</p> <table><tr><th rowspan="2">IBW</th><th colspan="5">Infusion rate (ml/hr)</th></tr><tr><th>1mg/kg/hr</th><th>2mg/kg/hr</th><th>3mg/kg/hr</th><th>4mg/kg/hr</th><th>5mg/kg/hr</th></tr><tr><td>40kg</td><td>0.8ml/hr</td><td>1.6ml/hr</td><td>2.4ml/hr</td><td>3.2ml/hr</td><td>4.0ml/hr</td></tr><tr><td>50kg</td><td>1.0ml/hr</td><td>2.0ml/hr</td><td>3.0ml/hr</td><td>4.0ml/hr</td><td>5.0ml/hr</td></tr><tr><td>60kg</td><td>1.2ml/hr</td><td>2.4ml/hr</td><td>3.6ml/hr</td><td>4.8ml/hr</td><td>6.0ml/hr</td></tr><tr><td>70kg</td><td>1.4ml/hr</td><td>2.8ml/hr</td><td>4.2ml/hr</td><td>5.6ml/hr</td><td>7.0ml/hr</td></tr><tr><td>80kg</td><td>1.6ml/hr</td><td>3.2ml/hr</td><td>4.8ml/hr</td><td>6.4ml/hr</td><td>8.0ml/hr</td></tr><tr><td>90kg</td><td>1.8ml/hr</td><td>3.6ml/hr</td><td>5.4ml/hr</td><td>7.2ml/hr</td><td>9.0ml/hr</td></tr><tr><td>100kg</td><td>2.0ml/hr</td><td>4.0ml/hr</td><td>6.0ml/hr</td><td>8.0ml/hr</td><td>10.0ml/hr</td></tr></table> <p>Administer intravenously via a central access device. Commence intravenous infusion at 1mg/kg/hr and titrate according to response (i.e achievement of burst suppression on the EEG).</p> <p>Ketamine has a low pH and may cause venous irritation and tissue damage in cases of extravasation.</p> <p>Ketamine does not need to be administered in a locked syringe in intensive care.</p>						IBW	Infusion rate (ml/hr)					1mg/kg/hr	2mg/kg/hr	3mg/kg/hr	4mg/kg/hr	5mg/kg/hr	40kg	0.8ml/hr	1.6ml/hr	2.4ml/hr	3.2ml/hr	4.0ml/hr	50kg	1.0ml/hr	2.0ml/hr	3.0ml/hr	4.0ml/hr	5.0ml/hr	60kg	1.2ml/hr	2.4ml/hr	3.6ml/hr	4.8ml/hr	6.0ml/hr	70kg	1.4ml/hr	2.8ml/hr	4.2ml/hr	5.6ml/hr	7.0ml/hr	80kg	1.6ml/hr	3.2ml/hr	4.8ml/hr	6.4ml/hr	8.0ml/hr	90kg	1.8ml/hr	3.6ml/hr	5.4ml/hr	7.2ml/hr	9.0ml/hr	100kg	2.0ml/hr	4.0ml/hr	6.0ml/hr	8.0ml/hr	10.0ml/hr
IBW	Infusion rate (ml/hr)																																																										
	1mg/kg/hr	2mg/kg/hr	3mg/kg/hr	4mg/kg/hr	5mg/kg/hr																																																						
40kg	0.8ml/hr	1.6ml/hr	2.4ml/hr	3.2ml/hr	4.0ml/hr																																																						
50kg	1.0ml/hr	2.0ml/hr	3.0ml/hr	4.0ml/hr	5.0ml/hr																																																						
60kg	1.2ml/hr	2.4ml/hr	3.6ml/hr	4.8ml/hr	6.0ml/hr																																																						
70kg	1.4ml/hr	2.8ml/hr	4.2ml/hr	5.6ml/hr	7.0ml/hr																																																						
80kg	1.6ml/hr	3.2ml/hr	4.8ml/hr	6.4ml/hr	8.0ml/hr																																																						
90kg	1.8ml/hr	3.6ml/hr	5.4ml/hr	7.2ml/hr	9.0ml/hr																																																						
100kg	2.0ml/hr	4.0ml/hr	6.0ml/hr	8.0ml/hr	10.0ml/hr																																																						
CONCENTRATION:	50mg/ml																																																										
STABILITY:	Physically and chemically stable for 24 hours at room temperature. Protect from light.																																																										

Critical Care Guidelines FOR CRITICAL CARE USE ONLY

References:

1. Kramer A.H. Early ketamine to treat refractory status epilepticus. *Neurocrit Care* (2012) 16: 299-305.
2. Shorvon S, Ferlisi M. The treatment of super-refractory status epilepticus. *Brain*. 2011; 134(10):2802-2818.
3. Chen JWY, Wasterlain CG. Status epilepticus: pathophysiology and management in adults. *Lancet Neurol* 2006; 5:246-56.
4. Sheth RD, Gidal BE. Refractory status epilepticus: response to ketamine. *Neurology*. Dec 1998; 51:1765.
5. Synowiec AS, Singh DS, Yenugadhati V, Valeriano JP, Schramke CJ et al. Ketamine use in the treatment of refractory status epilepticus. *Epilepsy Research* 2013 in press.
6. UKCPA Minimum Infusion Volumes. Critical Care Group 2012.
7. Injectable Medicines Guide online accessed on 25/09/2023.
8. Yao Fang, Xuefeng Wang: Ketamine for the treatment of refractory status epilepticus. *Seizure*. 30 (2015) 14-20.
9. Summary of Product Characteristics. Ketamine 50mg/ml Injection. Hameln pharmaceuticals. Last updated 26/06/20.
10. Summary of Product Characteristics. Ketamine 50mg/ml Solution for Injection. Panpharma UK Ltd. Last updated 27/09/21.
11. The Walton Centre NHS Foundation Trust: Status Epilepticus Guideline. Last updated May 2023. Accessed 25/09/23.
12. Erstad B, Barletta J. Drug dosing in the critically ill obese patient-a focus on sedation, analgesia and delirium. *Critical Care* 24. Article Number:315. 2020.

Title: KETAMINE	
ID:	Authors: Claire Hannah, Dr M Blackstock, Dr R Baruah, G Smyth.
Category:	Document Version: 3.0
Status Draft/Final: Final	Review Date: November 2025
Authoriser: Lothian Critical Care QIT Editorial Group	Authorisation Date: November 2023
Date added to Intranet: November 2023.	
Key Words	
Comments	