Critical Care Guidelines FOR CRITICAL CARE USE ONLY

DECEMBER OF THE PROPERTY OF TH			
PRODUCT	CLEANING PRODCEDURE	INSERTION PRODCEDURE	REMOVAL PROCEDURE
Clinell® wipes 2% chlorhexidine/ 70% alcohol	SmartSite® disinfection Urinary catheter port to obtain a CSU Blood culture bottle top disinfection Arterial line port disinfection Port disinfection prior to intrathecal drug administration Sampling from established EVD		
Clinell® wipes 2% chlorhexidine/ 70% alcohol	Skin disinfection for dressing changes. PVC, Arterial line Central line Central line = CVC, Quinton, AVA, PICC, Swan introducers, Mahurkar, PA Cath, ECMO	Skin disinfection before PVC insertion Skin disinfection before taking blood cultures	Skin disinfection before central line removal, and before epidural catheter removal
3 ml applicator ChloraPrep® with tint 2% chlorhexidine/ 70% alcohol		Inserting a butterfly into an established VAD Skin disinfection before perc trache insertion, arterial line insertion, central line insertion ie CVC, Quinton, AVA, PICC, Swan introducers, Mahurkar, PA Cath	
10.5 ml applicator ChloraPrep® with tint 2% chlorhexidine/ 70% alcohol	Skin disinifection before lumbar puncture	Skin disinifection before chest drain insertion Skin disinfection before intra Cranial Pressure bolt insertion	
26 ml applicator ChloraPrep® with tint 2% chlorhexidine/ 70% alcohol		Skin disinfection for ECMO line insertion	
Sterile saline 0.9% NaCl + sterile dressing pack	Chest drains dressing change		Chest drain removal Lumbar drain removal post TAAA

It is important to clean the area and allow drying time

Title: Chlorhexidine products		
ID: ChlohexProducts V1.0 20200228	Authors: Critical Care Infection Group	
Category:	Document Version: 1.0	
Status Draft/Final: Final	Review Date: 19/11/2021	
Authoriser: QIT	Date Authorisation: 19/11/2019	
Date added to Intranet:		
Key words: Chlorhexidine, ChloraPrep, Clinell, insertion, cleaning, removal		