Critical Care Guidelines FOR ICU USE ONLY



ALFENTANIL

PRESENTATION: Ampoules containing 5mg in 10ml or 25mg in

50ml (500micrograms/ml) of alfentanil.

INDICATION: Analgesia and/or endotracheal tube toleration.

DOSE AND ADMINISTRATION: ICU STANDARD INTRAVENOUS INFUSION

Alfentanil is administered undiluted e.g.15mg in

30ml or 25mg in 50ml according to infusion

rate.

Initially 1-4mls/hour (500micrograms-2mg/hour)

titrated to the required level.

In assisted ventilation, bolus doses of

500micrograms to 1 mg may be given.

CONCENTRATION: 500micrograms/ml

STABILITY: Physically and chemically stable for 24 hours at

room temperature.

References

- 1. Alfentanil 500micrograms/ml solution for injection. Summary of Product Characteristics. Hameln Pharmaceuticals Ltd. www.emc.medicines.org.uk. Updated on 22/04/20. Accessed 08/11/2022.
- 2. Critical Illness. Medicines Complete. <u>MedicinesComplete CONTENT > Critical Illness > Drug: Alfentanil</u>. Acessed 08/11/2022.
- 3. Medusa, NHS Injectable Medicines Guide. Alfentanil. Date published 13/07/2020. www.medusa.wales.nhs.uk. Accessed 08/11/2022.

Title: Alfentanil	
ID:	Authors: Claire Hannah
Category:	Document Version: 4.0
Status Draft/Final: FINAL	Review Date: Nov 24
Authoriser: Lothian Critical Care QIT	
Editorial Board.	Authorisation Date: Nov 2022
Date added to Intranet:	
Key Words	
Comments	

FOR ICU USE ONLY Critical Care Guidelines