

Lothian Intra-Aortic Balloon Pump Guidelines

PATIENT LABEL

Critical Care Guidelines
FOR CRITICAL CARE USE ONLY

Indication:

On Arrival

ECG	Position on CXR	Initial limb check		
Power on	He tank full	Tubing check	Pressure bag	

Day 1 Date:

Platelets	Urea	Creatinine	ECG	CXR	
				_	

	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
Power on																								
He tank level																								
Tubing check																								
Pressure bag																								
Insertion site																								
MAP																								
Aug. diastolic																								
Frequency																								
Trigger																								
♠inotropes																								
Rt pedal pulse																								
Lt pedal pulse																								
Lt radial pulse																								

Title: Lothian Critical Care Intra-Aortic Balloon Pump	Guideline 1
	Authors: A Abu-Arafeh G McNeill
Status Draft/Final: Final	Approved by: Directorate QIT Sept 2018
	Written: August 2018
Reviewed on:11.09.18	Next review : August 2020



Lothian Intra-Aortic Balloon Pump Guidelines

PATIENT LABEL

Critical Care Guidelines
FOR CRITICAL CARE USE ONLY

Day 2 Date:

Plate	elets		Ur	ea		Creat	inine			EC	CG		CXR			
															·	
	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	

	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
Power on																								
He tank level																								
Tubing check																								
Pressure bag																								
Insertion site																								
MAP																								
Aug. diastolic																								
Frequency																								
Trigger																								
♠inotropes																								
Rt pedal pulse																								
Lt pedal pulse		_	_		_	_	_	_	_					_	_	_	_				_			
Lt radial pulse																								

If circulatory support is anticipated to be required beyond 48 hours:

- -ICU consultant must liaise with cardiology consultant then document a plan for ongoing circulatory support
- -Anticoagulation should be considered with IV Heparin, if there are no contraindications
- -Consideration should be made to discussion with GJNH re: advanced circulatory support

Title: Lothian Critical Care Intra-Aortic Balloon Pump	Guideline 2
	Authors: A Abu-Arafeh G McNeill
Status Draft/Final: Final	Approved by: Directorate QIT Sept 2018
	Written: August 2018
Reviewed on:11.09.18	Next review : August 2020



Lothian Intra-Aortic Balloon Pump Guidelines

PATIENT LABEL

Critical Care Guidelines FOR CRITICAL CARE USE ONLY

Day 3							Date:	
	Platelets		Urea	Creatinine	ECG	CXR		
Outcom	e of discussion	on with	n GJNH					
Ongoing	circulatory s	suppor	t plan					
Anticoa	gulation plan							

	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23
Power on																								
He tank level																								
Tubing check																								
Pressure bag																								
Insertion site																								
MAP																								
Aug. Diastolic																								
Frequency																								
Trigger																								
↑ inotropes																								
Rt pedal pulse																								
Lt pedal pulse																								
Lt radial pulse																								

Title: Lothian Critical Care Intra-Aortic Balloon Pump	Guideline 3
	Authors: A Abu-Arafeh G McNeill
Status Draft/Final: Final	Approved by: Directorate QIT Sept 2018
	Written: August 2018
Reviewed on:11.09.18	Next review : August 2020