## Critical Care Guidelines FOR CRITICAL CARE USE ONLY

## Guideline for intra-hospital transfer by Advanced Critical Care Practitioners within NHS Lothian



Intra-hospital transfer of a level 3 critical care patient should only be undertaken by an Advanced Critical Care Practitioner (ACCP) within NHS Lothian when the following criteria are met:

- The decision for transfer is made by a **senior doctor** (Consultant, advanced trainee in Intensive Care Medicine, post-fellowship Anaesthetic trainee).
- There is an agreement from a senior doctor that the patient is suitable for transfer by an ACCP.
- There is an agreement from the ACCP that the patient is suitable for them to transfer.
- The ACCP has completed in-house training including supervised practice and has been signed off as having met basic competencies in intra-hospital transfer, and airway management.
- The patient should be physiologically stable.
- The transfer should take place 'in hours' 0800-2030.

Intra-hospital transfer by an ACCP is **not** appropriate in the following circumstances:

- The presence of severe cardiovascular instability (eg > 20 ml/hour single strength noradrenaline).
- High oxygen requirement (eg F<sub>i</sub>O<sub>2</sub> > 0.5)
- Known or anticipated difficult intubation.
- Patients with significant intracranial hypertension.
- Any other anticipated significant difficulty.

Title: Guideline for intra-hospital transfer by Advanced Critical Care Practitioners within NHS Lothian	
	Authors: G Nimmo, N Young,
Status: Final	Approved by: ACCP Clinical Leads, ACCP
	Managerial Leads
Reviewed on: G McNeill, G Renwick June 2020	Next review: June 2023