



Management of Subarachnoid Haemorrhage (SAH) Patients


Learning Outcomes



- Awareness of incidence and aetiology of SAH
- Appreciation of goals for care management
- Understand the components of secondary injury
- Know common complications to look out for
- Know what nursing and medical interventions are employed to prevent elevated ICP and other secondary injuries/complications.

Monro- Kellie Hypothesis





Putting SAH in Perspective



- SAH is a form of “stroke” and comprises around 7% of all strokes
- Incidence is around 9 per 100,000 per annum
- Up to 50% of all SAH are fatal, with 10 to 15% of patients dying before they reach hospital and a further 35-40% dying in hospital
- Approx 200-400 admissions to ICU per year in NHS Lothian

SUBARACHNOID HAEMORRHAGE




- Bleeding into the subarachnoid space (e.g. aneurysm or trauma)
- Risk of aneurysmal SAH increases with middle age and the elderly
- Smoking/excessive alcohol/ untreated hypertension
- Risk is 25% higher in women (? Hormonal changes)
- Genetics (3 to 5 fold ↑ risk in 1st degree relatives)

SUBARACHNOID HAEMORRHAGE



- Bleeding into the subarachnoid space (e.g. aneurysm or trauma)
- Risk of aneurysmal SAH increases with middle age and the elderly
- Smoking/excessive alcohol/ untreated hypertension
- Risk is 25% higher in women (? Hormonal changes)
- Genetics (3 to 5 fold ↑ risk in 1st degree relatives)

Common Sites



Symptoms/ Presentation



- ‘Thunderclap Headache’
- Nausea Vomiting
- Some patients experience ‘sentinel symptoms’ indicating leak before collapse
- collapse

Grading SAH



Hunt and Hess Classification of subarachnoid hemorrhage

Grade 1: Asymptomatic, mild headache, slight nuchal rigidity

Grade 2: Moderate to severe headache, nuchal rigidity, no neurologic deficit other than cranial nerve palsy

Grade 3: Drowsiness / confusion, mild focal neurologic deficit

Grade 4: Stupor, moderate-severe hemiparesis

Grade 5: Coma, decerebrate posturing

Tufts Comprehensive Stroke Center at
NEMC



(c) 2005, David E. Thaler, M.D., Ph.D.

Sedation hold within first 12 hours


Treatment



Clipping Treatment for Cerebral Aneurysm



Sub Arachnoid Haemorrhage



Basilar bifurcation aneurysm
before coil embolisation

after coil embolisation
showing complete obliteration.

Nursing Care



- As per protocol
- Head up
- Tape rather than tape
- $PO_2 > 12$
- PCO_2 5-6
- SBP 120-160 uncoiled
- SBP > 140
- Careful Fluid and Electrolyte balance
- Nimodipine
- Temperature control
- Scan if drop in GCS
- Phenytoin if seizure at the time of th aneurysm

Secondary Brain Injury/ Complications



- Decreased Cerebral Perfusion /blood flow (Hypotension, Intracranial Hypertension)
- Decreased Oxygen Supply (Hypoxia, Anaemia)
- Seizure
- Rebleed
- Delayed cerebra ischemia – caused by cerebral vasospasm.
- Hydrocephalus
- Neurogenic Pulmonary Oedema
- Myocardial

Cerebral Vasospasm



- *Cerebral vasospasm* is the prolonged, intense vasoconstriction of the larger conducting arteries in the subarachnoid space which is initially surrounded by a clot. Significant narrowing develops gradually over the first few days after the aneurysmal rupture
- Nimodipine has shown to be effective at prevention 60mg 4hrly
- Hypertension- if aneurysm secured (otherwise SBP 120-160)
- Worse 48hrs-10 days
- Causes significant secondary complications

SECONDARY INJURIES; THE CULPRITS!






The Eight H'S;

1. Hypoxia
2. Hypertension (Intracranial)
3. Hypotension
4. Hypo/ Hypercapnia
5. Hyponatraemia
6. Hypoglycaemia
7. Hyperpyrexia
8. Haematoma

Hydrocephalus



External Ventricular Drain




Therapeutic Goals



- $\text{SpO}_2 > 95\%$
- $\text{BP/MAP} > 90\text{mmHg}$
- Unsecured SAH- $\text{SBP } 120\text{-}160\text{mmHg}$ Secured $140\text{-}200\text{mmHg}$ (vasospasm)
- $\text{PaO}_2, > 12\text{kPa}$
- $\text{PaCO}_2 \text{ } 4.5\text{-}5\text{kPa}$
- Normothermia
- $\text{Hgb} > 90\text{g/l}$
- $\text{Glucose } 4.5\text{-}10\text{mmol/l}$

Abnormal ECGs




MEASURES TO REDUCE ICP



Consistently high ICP

RESCAN EARLY

- To exclude surgically remedial causes
- Decompressive hemi-craniectomy or bifrontal craniectomy (Stage 3 therapy)



Complications of SAH

- Cerebral Oedema/Swelling
- Hydrocephalus
- Neurogenic Pulmonary Oedema
- Disorders of Salt/Water Balance
- Seizures
- Vasospasm (ischemia)
- Rebleeding (aneurysm)
- ECG Changes
- VAP





- **DVT** (/Pneumatic Compression. Heparin later-once coiled and confirmed with neurosurgeons)
- **Stress Ulceration prophylaxis**
- **Bowels**

BRAIN STEM HERNIATION



- Continued ICP rise
- Cushings' reflex (widening of pulse pressure)
- CPP falls
- ICP so high that brain herniates
- BP falls
- Pupils fixed and dilated
- Brain-stem death tests

CARE OF THE FAMILY



- Empathetic
- Flexible regarding visiting
- Involve in care

PROGNOSTIC INDICATORS



- CAT SCAN
- GCS
- AGE
- PUPIL RESPONSE
- HAEMODYNAMICS

References



- The Guidelines for the Management of Severe Traumatic Brain Injury (Brain Trauma Foundation, 3rd Edition 2007)
- Scottish Intercollegiate Guidelines Network May 2009
- <http://www.sign.ac.uk/guidelines/fulltext/110/index.html>