Critical Care response to high risk haemorrhagic patients Department of Critical Care, RIE, WGH, SJH



ACTION CARD - NURSE IN CHARGE (NIC) CRITICAL CARE FOR VIRAL HAEMORRHAGIC FEVER (VHF)

On receipt of phone call requiring nurses to provide critical care support for patients in ED/RIDU/ARU isolation room:

- 1. Confirm information on patient risk assessment and discuss with Critical Care Consultant.
- 2. Allocate 2 nurses to be deployed to ED/RIDU/ARU, if staffing allows. (Ensure they have been PPE fitted and trained)
- 3. Locate all equipment, ready for transport to ED/RIDU/ARU if needed
 - Ventilator- Oxylog 3000 (ED will provide Oxylog 2000)
 - Mobile monitor
 - Ebola boxes
 - FFP3 masks (WGH) or Jupiter Hoods (RIE/SJH) for allocated staff
 - Ensure appropriate PPE available in ED/RIDU/ARU

4. Escalate:

- In hours:
 - CNM for Critical Care
 - Chief Nurse for RIE/WGH/SJH as appropriate
- Out of hours :
 - o Bleep 1590 (RIE), 8100 (WGH)
 - Contact Consultant on-call for Critical Care and ensure an appropriate member of medical staff is deployed to ED/RIDU/ARU to assess and potentially deliver critical care to the patient.
- 5. Liase with ED/RIDU/ARU NIC regularly
- 6. Arrange staffing provision of PPE trained nursing staff for subsequent shifts

Author: M Dunn, M Blackstock, S Wright v3.20141017