

PHYSICS 341, Assignment # 2
Due: Friday, September 29, 2017

(1) Suppose the force acting on a particle is given as $\mathbf{F} = f_0(y\hat{\mathbf{i}} + x\hat{\mathbf{j}})$. Two particles begin at the origin (0,0) and follow two different paths to the same point in the x - y plane. The first particle travels along a trajectory $y = \sqrt{ax}$ and the second particle travels along a trajectory $y = bx$.

- (a) At what point (x_m, y_m) do the two particles meet?
- (b) Compute the work done in moving the the first particle from the origin to (x_m, y_m) .
- (b) Compute the work done in moving the the second particle from the origin to (x_m, y_m) .
- (d) Is the work performed on both particles the same or different? Explain why.

(2) Two three-dimensional vectors are given as:

$$\mathbf{A} = 3\hat{\mathbf{e}}_1 - 2\hat{\mathbf{e}}_2 + \hat{\mathbf{e}}_3, \quad \mathbf{B} = -2\hat{\mathbf{e}}_1 + \hat{\mathbf{e}}_2 + 3\hat{\mathbf{e}}_3$$

find the following:

- (a) $\mathbf{A} - \mathbf{B}$ and $|\mathbf{A} - \mathbf{B}|$
- (b) the component of \mathbf{B} along \mathbf{A}
- (c) the angle between \mathbf{A} and \mathbf{B}
- (d) $\mathbf{A} \times \mathbf{B}$
- (e) $(\mathbf{A} - \mathbf{B}) \times (\mathbf{A} + \mathbf{B})$

(3) Using the cross-product relationships among the unit vectors $\hat{\mathbf{i}}$, $\hat{\mathbf{j}}$ and $\hat{\mathbf{k}}$ prove that the following triple vector product relation holds:

$$\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) = \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B}).$$

This is known as the *bac minus cab* rule and can be used to show that the cross product is not associative. (See Example 1.7.3 in the text.)

(4) (a) Find the transformation matrix that rotates the y -axis of a cartesian coordinate system 30° toward z around the x -axis.

(b) If the same rotation is performed twice what are the new coordinate directions $(\bar{x}, \bar{y}, \bar{z})$ in terms of the original directions (x, y, z) ?

(c) Is the rotation positive or negative and why?

(5) Find the values of σ needed to make the following transformation matrix describe an orthogonal transformation.

$$\mathbf{T} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \sigma & -\sigma \\ 0 & \sigma & \sigma \end{pmatrix}$$