

# Applied Probability

## 1 Continuous time Markov Chains

### 1.1 Poisson process

- Poisson process with intensity  $\lambda$

(i)  $N(0) = 0, N(s) \leq N(t)$  for  $s < t$

(ii)  $\mathbb{P}(N(t+h) = n+m | N(t) = n) = \begin{cases} \lambda h + o(h) & \text{if } m = 1 \\ o(h) & \text{if } m > 1 \\ 1 - \lambda h + o(h) & \text{if } m = 0 \end{cases}$

(iii)  $N(t) - N(s)$  independent of  $(N(k))_{k \leq s}$

**Theorem 1.1.**  $N(t) \sim Poi(\lambda t)$

*Proof.* derive differential equation, then generating function □

- $p_j(t) = \mathbb{P}(N(t) = j)$
- Generating function  $G(s, t) = \sum p_j(t) s^j$
- Arrival time  $T_n$
- interarrival time  $U_n$

**Theorem 1.2.**

(i)  $U_i \sim Exp(\lambda)$

(ii)  $U_i$  independent

*Proof.* use  $N(t)$  Poisson □

**Fact.**  $N(t) \geq j \iff T_j \leq t$

- order statistics

**Theorem 1.3.**  $T_1, \dots, T_n$  conditional on  $\{N(t) = n\}$  same as joint distribution of order statistics of  $n$  i.i.d. Uniform $[0, t]$

*Proof.*  $U$  to  $T$ , then calculate density □

## 1.2 Birth process

- birth process with birth rates  $\lambda_0, \lambda_1, \dots$
- (i)  $N(0) = 0, N(s) \leq N(t)$  for  $s < t$
- (ii)  $\mathbb{P}(N(t+h) = n+m | N(t) = n) = \begin{cases} \lambda_n h + o(h) & \text{if } m = 1 \\ o(h) & \text{if } m > 1 \\ 1 - \lambda_n h + o(h) & \text{if } m = 0 \end{cases}$
- (iii)  $N(t) - N(s)$  independent of  $(N(k))_{k \leq s}$

**Example.**

- (i) *Poisson process*:  $\lambda_n = \lambda$
- (ii) *Simple birth*:  $\lambda_n = n\lambda$
- (iii) *Simple birth with immigration*:  $\lambda_n = n\lambda + \nu$
- $T_\infty = \lim T_n = \sum_{i=0}^{\infty} U_i$
- non-explosive / honest —  $\mathbb{P}(T_\infty = \infty) = 1$

**Theorem 1.4.** birth process  $N$ ,  $\lambda_n > 0$ , then non-explosive  $\iff \sum_n \frac{1}{\lambda_n} = \infty$

**Lemma 1.5.**  $U_n \sim \text{Exp}(\lambda_n)$ , independent, then  $\mathbb{P}(T_\infty < \infty) = \begin{cases} 0 & \text{if } \sum_n \frac{1}{\lambda_n} = \infty \\ 1 & \text{if } \sum_n \frac{1}{\lambda_n} < \infty \end{cases}$

- forward system of equations:  $p'_{ij}(t) = \lambda_{j-1}p_{i,j-1}(t) - \lambda_j p_{ij}(t)$
- backward system of equations:  $p'_{ij}(t) = \lambda_i p_{i+1,j}(t) - \lambda_i p_{ij}(t)$

**Theorem 1.6.**

- (i) forward system has unique solution  $\{p_{ij}(t)\}$
- (ii)  $\{p_{ij}(t)\}$  satisfy backward system

**Theorem 1.7.**  $\{p_{ij}(t)\}$  unique solution of forward equations,  $\{\pi_{ij}(t)\}$  any solution of backward equations, then  $p_{ij}(t) \leq \pi_{ij}(t)$

**Fact.**  $\sum_j p_{ij}(t) = 1 \iff \mathbb{P}(T_\infty > t) = 1$

- weak Markov property
- stopping time

- strong Markov property
- right continuity
- stationary independent increments
  - (i)  $N(t) - N(s)$  only depends on  $t - s$
  - (ii)  $\{N(t_i) - N(s_i)\}$  independent where  $s_1 \leq t_1 \leq \dots \leq s_n \leq t_n$

### 1.3 Continuous Markov Chain

**Setting 1.**  $(X(t))$  takes values in countable  $S$

- Markov property —
  - $\mathbb{P}(X(t_n) = j | X(t_{n-1}) = i_{n-1}, \dots, X(t_1) = i_1) = \mathbb{P}(X(t_n) = j | X(t_{n-1}) = i_{n-1})$
- continuous-time Markov chain — right-continuous, Markov property
- transition probability  $p_{ij}(s, t) = \mathbb{P}(X(t) = j | X(s) = i)$
- homogeneous —  $p_{ij}(s, t) = p_{ij}(0, t - s)$
- transition semigroup  $(P_t)_{ij} = p_{ij}(t)$
- stochastic semigroup
  - (i)  $P_0 = I$
  - (ii)  $P_t$  stochastic — non-negative entries, row sum 1
  - (iii) (Chapman-Kolmogorov)  $P_{s+t} = P_s P_t$

**Setting 2.**  $(X(t))$  homogeneous Markov chain

**Theorem 1.8.**  $P_t$  stochastic semigroup

- $\mathbb{P}_i$  — probability measure conditional on  $X(0) = i$
- $\mathbb{E}_i$
- $t$ -historical — events given by  $\{X(s) : s < t\}$
- $t$ -future — events given by  $\{X(s) : s > t\}$
- stopping time  $T$  —  $\{T \leq t\}$  given by  $\{X(s) : s \leq t\}$

**Theorem 1.9** (Extended Markov property).  $H$   $t$ -historical,  $F$   $t$ -future, then  $\mathbb{P}(F | X(t) = j, H) = \mathbb{P}(F | X(t) = j)$

**Theorem 1.10** (Strong Markov property).  $T$  stopping time, conditional on  $\{T \leq T_\infty\} \cap \{X(T) = i\}$ , then

- (i)  $(X_{T+u})_u$  continuous Markov chain start at state  $i$
- (ii) same transition prob
- (iii) independent to  $\{X(s) : s < T\}$

**Setting 3.**  $X(0) = i$

–  $U_0 = \inf \{t : X(t) \neq i\}$

**Fact.** right continuous  $\Rightarrow U_0 > 0$

**Theorem 1.11.**

- (i)  $U_0 \sim \text{Exp}(g_i)$
- (ii)  $U_0$  stopping time

*Proof.* Extended Markov and homogeneity to deduce memoryless □

- transition matrix  $\mathbf{Y} = (y_{ij})$  —  $y_{ij} = \begin{cases} \delta_{ij} & \text{if } g_i = 0 \\ \mathbb{P}_i(X(U_0) = j) & \text{if } g_i > 0 \end{cases}$
- generator  $\mathbf{G} = (g_{ij})$  —  $g_{ij} = \begin{cases} g_i y_{ij} & \text{if } j \neq i \\ -g_i & \text{if } j = i \end{cases}$

**Fact.**  $\mathbb{P}(X(t+h) = j | X(t) = i) = g_{ij}h + o(h)$

**Fact.**  $g_{ij} = g_i(y_{ij} - \delta_{ij})$

**Theorem 1.12.**  $X(0) = i$ , then

- (i)  $X(U_0)$  independent of  $U_0$
- (ii) conditional on  $X(U_0) = j$ ,  $X^*(s) = X(U_0 + s)$  continuous-time Markov chain, same transition prob, initial state  $j$ , independent to the past

- $T_m$
- holding time  $U_m = T_{m+1} - T_m$
- jump chain  $Y = \{Y_n\}$
- $T_\infty = \lim T_n$
- minimal process
- explode from state  $i$  —  $\mathbb{P}_i(T_\infty < \infty) > 0$

**Proposition 1.13.**  $X$  minimal process, then  $P_{s+t} = P_s$

*Proof.* may go to  $\{\infty\}$  □

**Theorem 1.14.**  $i \in S$ , non-explosive from  $i$  if any of the following holds:

- (i)  $S$  finite
- (ii)  $\sup_j g_j < \infty$
- (iii)  $i$  recurrent in jump chain  $Y$

*Proof.* be dominated by Poisson process which is non-explosive □

- irreducible —  $\forall i, j, \exists t > 0, p_{ij}(t) > 0$

**Theorem 1.15.**

- (i) (Levy dichotomy)  $X$  irreducible, then  $\forall t > 0, p_{ij}(t) > 0$
- (ii)  $X$  irreducible  $\iff Y$  irreducible

*Proof.* look at jump chain,  $g_{i_0} \cdots g_{i_n} > 0, p_{i_k, i_{k+1}}(t) > 0$  □

**Fact.** birth process not irreducible

- recurrent —  $\mathbb{P}(\{t : X(t) = i\} \text{ unbounded}) = 1$
- transient —  $\mathbb{P}(\{t : X(t) = i\} \text{ unbounded}) = 0$
- $R_i = \inf \{t > U_0 : X(t) = i\}$
- mean return time  $m_i = \mathbb{E}(R_i)$
- positive recurrent / non-null recurrent —  $m_i < \infty$

**Theorem 1.16.** continuous-time chain  $X$ , jump chain  $Y$

- (i)  $g_i = 0$ , then  $i$  recurrent for  $X$
- (ii)  $g_i > 0$ , then  $i$  recurrent for  $X \iff$  recurrent for  $Y$
- (iii)  $i$  recurrent  $\iff \int p_{ii}(t)dt = \infty$
- (iv)  $i$  transient  $\iff \int p_{ii}(t)dt < \infty$
- (v)  $X$  irreducible, then every state recurrent or every state transient

*Proof.* main point is no explosion. Interchange summation, then old result.  $\square$

- **Forward equation:**  $P'_t = P_t G$  with boundary condition  $P_0 = 1$
- **Backward equation:**  $P'_t = G P_t$  with boundary condition  $P_0 = 1$

**Fact.** If states  $S$  finite, then  $P_t = e^{tG}$

- minimal solution —  $p_{ij}(t) \leq \pi_{ij}(t)$
- sub-stochastic —  $\sum_j p_{ij}(t) < 1$

**Theorem 1.17.**  $S$  countable,  $X$  minimal Markov chain with generator  $G$ , then

- (i)  $P_t$  minimal non-negative solution of backward equation  $P'_t = G P_t$  with boundary condition  $P_0 = 1$
- (ii)  $P_t$  minimal non-negative solution of forward equation  $P'_t = P_t G$

*Proof. Solution:* condition on  $T_1 > t$  or  $T_1 \leq t$ .

**Minimal:** reverse argument and induction.  $\square$

**Fact.** any solution to both equations sub-stochastic

**Fact.** non-explosive  $\Rightarrow P_t$  unique solution to both equations

- measure
- stationary measure —  $\pi = \pi P_t$
- stationary distribution
- unique measure — unique up to scalar multiplication
- first return time  $R_i$
- $m_i = \mathbb{E}_i(R_i)$

**Theorem 1.18.**  $X$  irreducible,  $|S| \geq 2$

- (i) some state  $k$  positive recurrent, then
  - (a)  $\exists$  unique stationary distribution  $\pi$
  - (b) unique distribution st  $\pi G = 0$
  - (c) all states positive recurrent
- (ii)  $X$  non-explosive,  $\exists \pi$  st  $\pi G = 0$ , then
  - (a) all states positive recurrent
  - (b)  $\pi$  stationary
  - (c)  $\pi_k = \frac{1}{m_k g_k}$

*Proof.* (i) use 1.19(iv)  $\pi = \mu(k)/m_k$ , then uniqueness of measure  $\Rightarrow$  all state non-null

(ii)  $\nu' = \frac{\pi_i g_i}{\pi_k g_k}$ , then  $\rho(k) \leq \nu'$  from discrete MC

□

–  $\nu_i = x_i g_i$

–  $\mu(k) = (\mu_j(k))_j \longrightarrow \mu_j(k) = \mathbb{E}_k \left( \int_0^{R_k} \mathbb{1} \{X(s) = j\} ds \right)$

–  $\rho(k) = (\rho_j(k)) \longrightarrow$  mean visit to  $j$  starting from  $k$  in jump chain  $Y$

**Lemma 1.19.**  $X$  irreducible Markov chain,  $|S| \geq 2$

(i) measure  $x$ , then  $xG = 0 \iff \nu Y = \nu$

(ii)  $X$  recurrent,  $xG = 0$  unique measure

(iii)  $x$  measure,  $xG = 0$ , then  $x_j > 0$

(iv)  $X$  recurrent,  $k \in S$ , then  $\mu(k)G = 0$  and stationary

*Proof.* (i) expand

(ii)  $\nu Y = \nu$ , then uniqueness from discrete MC

(iii)  $\mu_j(k) = \frac{1}{g_j} \rho_j(k)$ , then  $\rho(k)Y = \rho(k)$  from discrete MC

(iv) strong Markov to shift time  $t$

□

**Fact.**  $X$  non-explosive, then  $R_k = \sum_j \int_0^{R_k} \mathbb{1} \{X(s) = j\} ds$

**Fact.**  $X$  irreducible,  $\exists$  more than one stationary distribution, then  $X$  explosive

**Theorem 1.20** (Markov chain limit theorem).  $X$  irreducible, non-explosive

(i) if  $\exists$  stationary distribution  $\pi$ , then

(a)  $\pi$  unique

(b)  $p_{ij}(t) \rightarrow \pi_j$

(ii) if no stationary distribution, then  $p_{ij}(t) \rightarrow 0$

*Proof.* skeleton  $Z_n = X(nh)$

□

**Lemma 1.21.**  $X$  minimal, then  $|p_{ij}(t+u) - p_{ij}(t)| \leq 1 - e^{-g_i u}$

## 1.4 Birth-death process and imbedding

- birth rate  $\lambda_0, \lambda_1, \dots$
- death rate  $\mu_1, \mu_2, \dots$
- birth-death process

**Theorem 1.22.**  $X$  birth-death process, generator  $G$

- (i) measure  $x_n = \frac{\lambda_0 \lambda_1 \dots \lambda_{n-1}}{\mu_1 \mu_2 \dots \mu_n} x_0$  satisfies  $\mathbf{x}G = 0$
- (ii)  $\exists$  distribution  $\pi$  satisfies  $\pi G = 0 \iff \sum \frac{\lambda_0 \lambda_1 \dots \lambda_{n-1}}{\mu_1 \mu_2 \dots \mu_n} < \infty$
- (iii) if  $\sum \frac{\lambda_0 \lambda_1 \dots \lambda_{n-1}}{\mu_1 \mu_2 \dots \mu_n} (\lambda_n + \mu_n) < \infty$ , then  $\pi$  stationary

*Proof.* (i) solve  $\mathbf{x}G = 0$

(ii) trivial

(iii) condition for jump chain  $Y$  recurrent, then non-explosive

□

**Example.**

- **Pure birth**  $\mu_n = 0$
- **Simple death with immigration**  $\lambda_n = \lambda, \mu_n = n\mu$

**Theorem 1.23.**  $X(t)$  asymptotically  $\text{Poi}(\rho) = \text{Poi}\left(\frac{\lambda}{\mu}\right)$

- **Simple birth-death**  $\lambda_n = n\lambda, \mu_n = n\mu, X(0) = I$

**Fact.** state 0 absorbing

**Theorem 1.24.**  $G(s, t) = \mathbb{E}(s^{X(t)}) = \begin{cases} \left( \frac{\lambda t(1-s) + s}{\lambda t(1-s) + 1} \right)^I & \text{if } \mu = \lambda \\ \left( \frac{\mu(1-s) - (\mu - \lambda s)e^{-t(\lambda - \mu)}}{\lambda(1-s) - (\mu - \lambda s)e^{-t(\lambda - \mu)}} \right)^I & \text{if } \mu \neq \lambda \end{cases}$

*Proof.* Forward equation

□

**Fact.** non-explosive as  $\sum p_j(t) = G(1, t) = 1$

**Fact.**  $\mathbb{E}_I(X(t)) \rightarrow \begin{cases} 0 & \text{if } \rho < 1 \\ \infty & \text{if } \rho > 1 \end{cases}$

- extinction probability  $\eta(t) = \mathbb{P}_I(X(t) = 0)$



**Corollary 1.25.**  $\eta(t) \rightarrow \begin{cases} 1 & \text{if } \rho \leq 1 \\ \rho^{-I} & \text{if } \rho > 1 \end{cases}$

- imbedded random walk — jump chain  $Y$  with parameter  $\frac{\lambda}{\lambda+\mu}$ , absorbing at 0
- imbedded branching process — lives  $Exp(\lambda+\mu)$ , then born  $n$  individuals where  $\begin{cases} p_0 = \mathbb{P}(n=0) = \frac{\mu}{\lambda+\mu} \\ p_2 = \mathbb{P}(n=2) = \frac{\lambda}{\lambda+\mu} \end{cases}$
- age-dependent branching process
- age density function  $f_T(u) = (\lambda + \mu)e^{-(\lambda+\mu)u}$
- family-size generating function  $G(s) = \frac{\mu+\lambda s^2}{\mu+\lambda} = p_0 + p_2 s^2$