

# Numerical Solution to the Optimal Portfolio Selection Problem for Power Utility Function

by

Kenrick So

## Chapter 1

# Numerical Solution to the Optimal Portfolio Selection Problem for Power Utility Function

### 1.1 Problem Setup

Consider a financial market containing two financial assets that are traded continuously on a finite horizon  $[0, T]$ . The first asset is a risk-free bond  $P$  that evolves according to the following ordinary differential equation

$$dP_t = rP_t dt, \quad t \in [0, T], \quad (1.1)$$

where  $r$  is a risk-free rate. The second asset is a risky stock that evolves according to the following stochastic differential equation

$$dS_t = \mu S_t dt + \sigma S_t dW_t, \quad t \in [0, T] \quad (1.2)$$

where  $\mu \in \mathbb{R}$  is a constant drift term,  $\sigma \geq 0$  is a constant volatility term, and  $W_t$  is a standard Brownian motion.

The investor is interested in determining an efficient strategy for her final wealth. Define the wealth process  $X_t$ , at any time  $t < T$ , the investor needs to decide what proportion  $u_t$  of her wealth to invest in the risky asset  $S_t$  and invest the remaining proportion  $1 - u_t$  to the risk-free bond. The wealth process must evolve according to the following stochastic differential equation

$$\begin{aligned} dX_t &= u_t X_t \frac{dS_t}{S_t} + (1 - u_t) X_t r dt \\ &= u_t X_t (\mu dt + \sigma dW_t) + (1 - u_t) X_t r dt \\ &= u_t X_t \mu dt + u_t X_t \sigma dW_t + X_t r dt - u_t X_t r dt \\ &= [r + u_t(\mu - r)] X_t dt + X_t u_t \sigma dW_t. \end{aligned} \quad (1.3)$$

Assume that the initial wealth is positive i.e.  $X_0 = x_0 > 0$ . The investor is interested in finding an optimal investment strategy  $u_t^*$  such that the expected utility of the terminal

wealth  $X_T$  is maximized. The objective function is given by,

$$\max_{u(\cdot) \in \mathcal{A}} \mathbb{E}[U(X_T^u)]. \quad (1.4)$$

Here,  $\mathcal{A}$  is the set of admissible controls if the investor had an initial endowment of  $x_0$ , and  $U(x)$  is a utility function that is strictly increasing and concave up. We apply the power utility function such that the objective function becomes,

$$\max_{u(\cdot) \in \mathcal{A}} \mathbb{E} \left[ \frac{(X_T^u)^\gamma}{\gamma} \right]. \quad \gamma \in (0, 1) \quad (1.5)$$

By the martingale optimality principle and the verification theorem for Hamilton-Jacobi-Bellman (HJB) equation for the Merton problem, the portfolio problem (1.4) can be evaluated by solving the following HJB-equation

$$\begin{cases} \frac{\partial V}{\partial t} + \max_{u_t \in \mathcal{A}} \left\{ [u_t(\mu - r) + r]X_t \frac{\partial V}{\partial x} + \frac{1}{2}X_t^2 \sigma^2 u_t^2 \frac{\partial^2 V}{(\partial x)^2} \right\} = 0 \\ V(T, x) = \frac{x^\gamma}{\gamma}. \end{cases} \quad (1.6)$$

## 1.2 Numerical Scheme for the HJB Equation

The PDE (1.6) can be written as,

$$\begin{cases} \frac{\partial V}{\partial t} + \max_{u_t \in \mathcal{A}} \{\mathcal{L}^u V\} = 0 \\ V(T, x) = \frac{x^\gamma}{\gamma}. \end{cases} \quad (1.7)$$

Here,

$$\begin{cases} \mathcal{L}^u V = a(X_t, u_t) \frac{\partial^2 V}{(\partial x)^2} + b(X_t, u_t) \frac{\partial V}{\partial x} \\ a(X_t, u_t) = \frac{1}{2}X_t^2 \sigma^2 u_t^2 \\ b(X_t, u) = [r + u_t(\mu - r)]X_t. \end{cases} \quad (1.8)$$

$\mathcal{L}$  is the infinitesimal generator of the process. The domain of the PDE is  $\Omega = \{(t, X_t) \in [0, T] \times \mathbb{R}\}$ .

## 1.3 Boundary Conditions

In order to evaluate the HJB PDE numerically, boundary conditions need to be established. At the boundary  $X_t = 0$ , the PDE reduces to

$$\frac{\partial V}{\partial t} = 0 \quad (1.9)$$

Furthermore, for large  $X_t$ , we make the assumption that the optimal control is  $u = 0$ . The intuition for this assumption is that for a sufficiently large wealth, the marginal gain an investor generates from investing in the risky asset is small compared to investing in the risk-free asset. The PDE (1.6) reduces to

$$\frac{\partial V}{\partial t} + rX_t \frac{\partial V}{\partial x} = 0. \quad (1.10)$$

Define  $\tau = T - t$ , the solution (See Appendix A) to the PDE above is given by,

$$V(T - \tau, x) = \delta^2(\tau)x^2 + \delta(\tau)x, \quad (1.11)$$

where,

$$\delta(\tau) = \exp(r\tau) \quad (1.12)$$

## 1.4 Finite Difference

The first step is to localize the PDE to a finite interval  $[0, X_{max}]$ . Suppose that the price process can be discretized into  $M$  nodes, and the time process can be discretized into  $N$  nodes. Define  $\Delta x = \frac{X_{max}}{M-1}$  and  $\Delta \tau = \frac{T}{N-1}$ ,

$$\begin{aligned} x_i &= (i-1)\Delta x, \quad i = 1, 2, \dots, M \\ \tau_n &= (n-1)\Delta \tau, \quad n = 1, 2, \dots, N. \end{aligned} \quad (1.13)$$

The values at each grid point is denoted as  $V_i^n = V(\tau_n, x_i)$ . The infinitesimal generator  $\mathcal{L}^u V$  can be approximated as,

$$(\mathcal{L}_{\Delta x}^u V)_i = \alpha_i V_{i-1} + \beta_i V_{i+1} - (\alpha_i + \beta_i) V_i. \quad (1.14)$$

The PDE (1.6) can be approximated using an explicit timestepping as,

$$\begin{aligned} \frac{V_i^{n+1} - V_i^n}{\Delta t} + \max_{u_t \in \mathbb{R}} [(\mathcal{L}_{\Delta x}^u V^n)_i] &= 0 \\ \frac{V_i^{n+1} - V_i^n}{\Delta t} + \alpha_i^n V_{i-1}^n + \beta_i^n V_{i+1}^n - (\alpha_i^n + \beta_i^n) V_i^n &= 0 \end{aligned} \quad (1.15)$$

From (1.8), note that  $a(X_t, u)$  is always positive, while  $b(X_t, u)$  is not always positive. When either  $\alpha_i$  or  $\beta_i$  is negative, oscillations may appear in the numerical solution. The numerical scheme presented involves the maximal use of central differences to improve the convergence rate and accuracy. In cases that either  $\alpha_i$  or  $\beta_i$  is negative, we adopt forward or backward differences to ensure that the new coefficient is positive. The central

difference is given by,

$$\begin{aligned}\alpha_{i,central}^n &= \frac{a(x_i, u^n)}{\Delta x^2} - \frac{b(x_i, u^n)}{2\Delta x} \\ \beta_{i,central}^n &= \frac{a(x_i, u^n)}{\Delta x^2} + \frac{b(x_i, u^n)}{2\Delta x}\end{aligned}\tag{1.16}$$

If  $\alpha_{i,central}^n < 0$ , we apply a forward difference to  $\frac{\partial V}{\partial x}$ ,

$$\begin{aligned}\alpha_{i,forward}^n &= \frac{a(x_i, u^n)}{\Delta x^2} \\ \beta_{i,forward}^n &= \frac{a(x_i, u^n)}{\Delta x^2} + \frac{b(x_i, u^n)}{\Delta x}.\end{aligned}\tag{1.17}$$

If  $\beta_{i,central}^n < 0$ , we apply a backward difference to  $\frac{\partial V}{\partial x}$ ,

$$\begin{aligned}\alpha_{i,backward}^n &= \frac{a(x_i, u^n)}{\Delta x^2} - \frac{b(x_i, u^n)}{\Delta x} \\ \beta_{i,backward}^n &= \frac{a(x_i, u^n)}{\Delta x^2}\end{aligned}\tag{1.18}$$

More detail is provided in Appendix B.

## 1.5 Numerical Algorithm

Define the matrix,

$$A^n(u^n) = \begin{pmatrix} -(\alpha_1^n + \beta_1^n) & \beta_1^n & & & \\ \alpha_2^n & -(\alpha_2^n + \beta_2^n) & \beta_2^n & & \\ & \ddots & \ddots & \ddots & \\ & \alpha_{M-2}^n & -(\alpha_{M-2}^n + \beta_{M-2}^n) & \beta_{M-2}^n & \\ & & \alpha_{M-1}^n & -(\alpha_{M-1}^n + \beta_{M-1}^n) & \\ 0 & 0 & \dots & 0 & \end{pmatrix}.$$

Then,

$$\begin{aligned}(\mathcal{L}_{\Delta x}^u V)_i &= \alpha_i V_{i-1} + \beta_i V_{i+1} - (\alpha_i + \beta_i) V_i \\ &= [A^n(u^n) V^n]_i.\end{aligned}\tag{1.19}$$

Define the boundary condition vector  $G^n = [0, 0, \dots, 0, G_M^n]$ , where the entries are populated using the equation for the boundary condition for large  $X_t$ ,

$$\begin{aligned}V(T - \tau, x_{max}) &= \delta(\tau) x_{max}^2 + \delta(\tau) x_{max} \\ \rightarrow V(t, x_{max}) &= \delta(T - t) x_{max}^2 + \delta(T - t) x_{max}\end{aligned}\tag{1.20}$$

such that

$$\begin{aligned} G_u^M &:= V(n\Delta t, x_{max}) \\ &= \delta(T - n\Delta t)x_{max}^2 + \delta(T - n\Delta t)x_{max} \end{aligned} \quad (1.21)$$

Then the discretized PDE (1.15) can be written in terms of the control vector  $u^n$

$$\begin{aligned} \frac{V_i^{n+1} - V_i^n}{\Delta t} + \max_{u_t \in \mathbb{R}} [\mathcal{L}^u V^n]_i &= 0 \\ \frac{V_i^{n+1} - V_i^n}{\Delta t} + [A^n(u^n) V^n]_i &= 0 \\ V_i^{n+1} - V_i^n + A^n(u^n) V^n \Delta t &= 0 \\ [I - A^n(u^n) \Delta t] V^n &= V^{n+1} + (G^{n+1} - G^n) \end{aligned} \quad (1.22)$$


Here,  $(G^{n+1} - G^n)$  enforces the boundary condition at  $x_{max}$ .

## 1.6 Simulation

The scheme described above is implemented in Python with the following parameter values:

- $r = 0.03$
- $\mu = 0.02$
- $\sigma = 0.15$
- $T = 20.0$
- $\gamma = 0.5$
- $x_{max} = 5.0$
- $M = 100$
- $N = 1600$
- $U_{max} = 1$

A plot of the value function  $V(X_t, t)$  versus the wealth process  $X_t$  and a plot of the optimal control  $u$  against wealth  $X_t$  derived from the numerical scheme above is presented below:



## 1.7 Solution to the PDE (Appendix A)

Assume that the solution to the PDE is given by,

$$V(T - \tau, X_t) = \delta^2(\tau)X_t^2 + \delta(\tau)X_t \quad (1.23)$$

where

$$\delta(\tau) = \exp(r\tau). \quad (1.24)$$

It follows that

$$\begin{aligned} \frac{\partial V}{\partial t} &= \frac{\partial V}{\partial \tau} \frac{\partial \tau}{\partial t} \\ &= -\frac{\partial V}{\partial \tau} \\ &= -2r \exp(2r\tau)X_t^2 - r \exp(r\tau)X_t \end{aligned} \quad (1.25)$$

Also,

$$\begin{aligned} rX_t \frac{\partial V}{\partial X_t} &= rX_t(2X_t\delta(\tau) + \delta(\tau)) \\ &= rX_t(2X_t \exp(2r\tau) + \exp(r\tau)) \\ &= 2r \exp(2r\tau)X_t^2 + r \exp(r\tau)X_t. \end{aligned} \quad (1.26)$$

Hence,

$$\begin{aligned} \frac{\partial V}{\partial t} + rX_t \frac{\partial V}{\partial X_t} &= -2r \exp(2r\tau)X_t^2 - r \exp(r\tau)X_t + 2r \exp(2r\tau)X_t^2 + r \exp(r\tau)X_t \\ &= 0. \end{aligned} \quad (1.27)$$

Therefore,  $V(t, X_t) = \delta(\tau)X_t^2 + \delta(\tau)X_t$  is a solution to the PDE.



## 1.8 Discretization Scheme (Appendix B)

We discretize the operator  $\mathcal{L}^u V$  and derive the coefficients  $\alpha_i^{n+1}$  and  $\beta_i^{n+1}$ ,

$$\mathcal{L}^u V = a(X_t, u_t) \frac{\partial^2 V}{(\partial x)^2} + b(X_t, u_t) \frac{\partial V}{\partial x} a(X_t, u_t) = \frac{1}{2} X_t^2 \sigma^2 u_t^2. \quad (1.28)$$

Suppose we apply central differencing to  $\frac{\partial V}{\partial x}$  and  $\frac{\partial^2 V}{(\partial x)^2}$ , then

$$\begin{aligned} \frac{\partial V}{\partial x} &\approx \frac{V_{i-1} + V_{i+1}}{\Delta x} \\ \frac{\partial^2 V}{(\partial x)^2} &\approx \frac{V_{i-1} + V_{i+1} - 2V_i}{(\Delta x)^2} \end{aligned} \quad (1.29)$$

Then, (1.28) results in,

$$\mathcal{L}^u V \approx a(X_t, u_t) \frac{V_{i-1} + V_{i+1} - 2V_i}{(\Delta x)^2} + b(X_t, u_t) \frac{V_{i-1} + V_{i+1}}{\Delta x} \quad (1.30)$$

Matching the coefficient with (1.14) results in:

$$\alpha_i V_{i-1} + \beta_i V_{i+1} - (\alpha_i + \beta_i) V_i = a(X_t, u_t) \frac{V_{i-1} + V_{i+1} - 2V_i}{(\Delta x)^2} + b(X_t, u_t) \frac{V_{i-1} + V_{i+1}}{\Delta x} \quad (1.31)$$

For the above equation to hold,

$$\begin{aligned} \alpha_{i,central}^n &= \frac{a(x_i, u^n)}{\Delta x^2} - \frac{b(x_i, u^n)}{2\Delta x} \\ \beta_{i,central}^n &= \frac{a(x_i, u^n)}{\Delta x^2} + \frac{b(x_i, u^n)}{2\Delta x} \end{aligned} \quad (1.32)$$

Suppose instead that  $\alpha_{i,central}^n < 0$  and we apply forward difference to  $\frac{\partial V}{\partial x}$ , and central difference to  $\frac{\partial^2 V}{(\partial x)^2}$ , then

$$\begin{aligned} \frac{\partial V}{\partial x} &\approx \frac{V_{i+1} - V_i}{\Delta x} \\ \frac{\partial^2 V}{(\partial x)^2} &\approx \frac{V_{i+1} + V_{i-1} - 2V_i}{(\Delta x)^2} \end{aligned} \quad (1.33)$$

Matching the coefficient with (1.14) results in:

$$\alpha_i V_{i-1} + \beta_i V_{i+1} - (\alpha_i + \beta_i) V_i = a(X_t, u_t) \frac{V_{i+1} + V_{i-1} - 2V_i}{(\Delta x)^2} + b(X_t, u_t) \frac{\partial V}{\partial x} \frac{V_{i+1} - V_i}{\Delta x} \quad (1.34)$$

For the above equation to hold,

$$\begin{aligned}\alpha_{i,forward}^n &= \frac{a(x_i, u^n)}{\Delta x^2} \\ \beta_{i,forward}^n &= \frac{a(x_i, u^n)}{\Delta x^2} + \frac{b(x_i, u^n)}{\Delta x}.\end{aligned}\tag{1.35}$$

Suppose instead that  $\beta_{i,central}^n < 0$  and we apply backward difference to  $\frac{\partial V}{\partial x}$ , and central difference to  $\frac{\partial^2 V}{(\partial x)^2}$ , then

$$\begin{aligned}\frac{\partial V}{\partial x} &\approx \frac{V_i - V_{i-1}}{\Delta x} \\ \frac{\partial^2 V}{(\partial x)^2} &\approx \frac{V_{i+1} + V_{i-1} - 2V_i}{(\Delta x)^2}\end{aligned}\tag{1.36}$$

Matching the coefficient with (1.14) results in:

$$\alpha_i V_{i-1} + \beta_i V_{i+1} - (\alpha_i + \beta_i) V_i = a(X_t, u_t) \frac{V_{i+1} + V_{i-1} - 2V_i}{(\Delta x)^2} + b(X_t, u_t) \frac{V_i - V_{i-1}}{\Delta x}\tag{1.37}$$

For the above equation to hold,

$$\begin{aligned}\alpha_{i,backward}^n &= \frac{a(x_i, u^n)}{\Delta x^2} - \frac{b(x_i, u^n)}{\Delta x} \\ \beta_{i,backward}^n &= \frac{a(x_i, u^n)}{\Delta x^2}\end{aligned}\tag{1.38}$$