

Energetics at mesoscopic scale

Ken Sekimoto (PCT-ESPCI, MSC-P7)
ken.sekimoto@espci.fr

June 8, 2009 CEA

Introduction

A few key papers of theory of fluctuations ...

1905 – *Einstein's relation* (Fluctuation and dissipation)

1906 – Smoluchowski (diffusion process)

1908 – Langevin's equation (equation of *stochastic* motion)

1915 – Fokker (diffusion process)

1917 – Planck (diffusion process)

...

1940 – Kramers' equation (Brownian motion and chemical reaction)

1951 – Itô; "*On stochastic differential equations*" (mathematization!)

1960 – Zwanzig's *projection method* (from Liouville eq. to *diffusion*)

1963 – Feynman's ratchet wheel and pawl (mesoscopic heat engine)

1965 – *Mori's formula* (from Hamiltonian to linear Langevin eq.)

1973 – *Kawasaki's identity* (non-linear extension of *Mori's formula*)

...

1990's – Jarzynski's equality, Fluctuation Theorems, *StEng*

Introduction

A few key papers of mesoscopic experiments ...

1827 – Robert *Brown*

1913 – Perrin; *Les Atomes* (experimental proofs)

1960 – Perutz *et al.* (X-ray structure of protein)

1960's – Shimomura (Green Fluorescent Protein; GFP)

1970 – Ashkin (optical tweezer)

1980's – Binnig *et al.* (STM, AFM)

1984 – Yanagida, *et al.* (dynamics of single filament)

1993 – Raymer (X-ray structure of *myosin*)

1990's – (single protein probe)

Introduction

A few key papers of mesoscopic experiments ...

1827 – Robert *Brown*

1913 – Perrin; *Les Atomes* (experimental proofs)*

1960 – Perutz *et al.* (X-ray structure of protein)

1960's – Shimomura (Green Fluorescent Protein)*

1970 – Ashkin (optical tweezer)

1980's – Bennig *et al.* (STM, AFM)

1984 – Yanagida, *et al.* (dynamics of single filament)

1993 – Rayment (X-ray structure of *myosin*)

1990's – (single protein probe)



Perrin 1913

Introduction

A few key papers of mesoscopic experiments ...

1827 – Robert *Brown*

1913 – Perrin; *Les Atomes* (experimental proofs)

1960 – Perutz *et al.* (X-ray structure of protein)

1960's – Shimomura (Green Fluorescent Protein)*


1970 – Ashkin (optical tweezer)

1980's – Bennig *et al.* (STM, AFM)

1984 – Yanagida, *et al.* (dynamics of single filament)*

1993 – Rayment (X-ray structure of *myosin*)

1990's – (single protein probe)



(GFP)

Introduction

A few key papers of mesoscopic experiments ...

1827 – Robert *Brown*

1913 – Perrin; *Les Atomes* (experimental proofs)

1960 – Perutz *et al.* (X-ray structure of protein)

1960's – Shimomura (Green Fluorescent Protein)

1970 – Ashkin (optical tweezer)

1980's – Bennig *et al.* (STM, AFM)

1984 – Yanagida, *et al.* (dynamics of single filament)*

1993 – Rayment (X-ray structure of *myosin*)*

1990's – (single protein probe)



Yanagida *et al.* 1984

Introduction

A few key papers of mesoscopic experiments ...

1827 – Robert *Brown*

1913 – Perrin; *Les Atomes* (experimental proofs)

1960 – Perutz *et al.* (X-ray structure of protein)

1960's – Shimomura (Green Fluorescent Protein)


1970 – Ashkin (optical tweezer)

1980's – Bennig *et al.* (STM, AFM)

1984 – Yanagida, *et al.* (dynamics of single filament)

1993 – Rayment (X-ray structure of *myosin*)*

1990's – (single protein probe)*



Rayment et al. 1993

Introduction

A few key papers of mesoscopic experiments ...

1827 – Robert *Brown*

1913 – Perrin; *Les Atomes* (experimental proofs)

1960 – Perutz *et al.* (X-ray structure of protein)

1960's – Shimomura (Green Fluorescent Protein)


1970 – Ashkin (optical tweezer)

1980's – Bennig *et al.* (STM, AFM)

1984 – Yanagida, *et al.* (dynamics of single filament)

1993 – Rayment (X-ray structure of *myosin*)

1990's – (single protein probe)*




Kitamura *et al.* 1999

Introduction

Short summary

More and more *mesoscopic* phenomena
— described by Langevin equations



Reminder : Langevin equation

"Newton's 1st law" → *Free Brownian motion without external field*

$$0 = -\underbrace{\gamma \frac{dx}{dt}}_{\text{forces from environment}} + \sqrt{2\gamma k_B T} \theta(t), \quad \langle \theta(t) \rangle = 0, \quad \langle \theta(t) \theta(t') \rangle = \delta(t - t')$$

$\theta(t)$: white Gaussian process
($\int_0^t \theta(s) ds = B_t$: Wiener process)

"Newton's 2nd law" → *Brownian motion under external field*

$$0 = -\frac{\partial U(x, a)}{\partial x} - \gamma \frac{dx}{dt} + \sqrt{2\gamma k_B T} \theta(t)$$

a : control parameter (by an external system)


+ many generalizations e.g. D -dim, inertia, memory, manifold,...

Introduction

What have been *less* discussed ?

1. Missing link – energetics absent (cf. QM, hydro, Newton, gen-rel)

cf. Known : steady-state of Langevin eq \leftrightarrow canonical ensemble




Introduction

What have been *less* discussed ?

1. Missing link (cf. QM,hydro,Newton,GR) – energetics absent

e.g. *Feynman's ratchet wheel and pawl*




Modelling by Langevin equation ? Efficiency ? (ks 1997)

Introduction

What have been *less* discussed ?

1. Missing link (cf. QM,hydro,Newton,GR) – energetics absent

e.g. *Protein motors*



cf.
ATPase free energy
~70% : *entropic*
~30% : *energetic*

Introduction

What have been *less* discussed ?

1. Missing link (cf. QM, hydro, Newton, GR) – energetics absent
2. How do the fluctuations look *at* mesoscale ?


a) Analyses along a single realisations (cf. ensemble properties)

$$0 = -\frac{\partial U(x, a)}{\partial x} - \gamma \frac{dx}{dt} + \sqrt{2\gamma k_B T} \theta(t)$$


b) "Newton's 3rd law" → [force from environment]
= -[force onto environment]

Energetics at mesoscopic scales

Thermodynamics



Langevin equation (single trajectory)



Energetics at mesoscopic scales

$$0 = -\frac{\partial U(x, a)}{\partial x} - \gamma \frac{dx}{dt} + \sqrt{2\gamma k_B T} \theta(t)$$


Definition of *mesoscopic heat*

heat, $d'Q$: work done by the heat bath
through a (true) displacement, dx

$$d'Q \equiv \left(-\gamma \frac{\partial x}{\partial t} + \sqrt{2\gamma k_B T} \theta(t) \right) \circ dx(t)$$

$$A(t) \circ dx(t) \equiv \frac{A(t+dt) + A(t)}{2} [x(t+dt) - x(t)]$$

STRATONOVICH type product



e.g. Free Brownian motion; $d'Q = 0$ (for overdamped case).

Energetics at mesoscopic scales

1st law of stochastic energetics

$$\text{heat : } d'Q \equiv \left(-\gamma \frac{\partial x}{\partial t} + \sqrt{2\gamma k_B T} \theta(t) \right) \circ dx(t) = \frac{\partial U}{\partial x} \circ dx(t)$$


$$0 = -\frac{\partial U(x, a)}{\partial x} - \gamma \frac{dx}{dt} + \sqrt{2\gamma k_B T} \theta(t)$$

$$\text{energy change : } dE = dU(x, a)$$

$$\text{work by external system : } d'W \equiv \frac{\partial U}{\partial a} \circ da(t)$$

$$\rightarrow dE = d'Q + d'W$$

Energy balance (along a single trajectory)



Energetics at mesoscopic scales

Experimental demonstrations of *energy balance*


PRL **96**, 070603 (2006)

PHYSICAL REVIEW LETTERS

week ending
24 FEBRUARY 2006

Thermodynamics of a Colloidal Particle in a Time-Dependent Nonharmonic Potential

V. Blickle,¹ T. Speck,² L. Helden,¹ U. Seifert,² and C. Bechinger¹



$z(t)$: Single-Particle
Tracking (SPT)

a : laser force \uparrow

$U(z,a)$: equilibrium
distribution data

Energetics at mesoscopic scales


Experimental demonstrations of *energy balance*

Journal of Statistical Mechanics: Theory and Experiment
An IOP and SISSA journal

stacks.iop.org/JSTAT/2007/P09018

Fluctuation theorems for harmonic oscillators

S Joubaud, N B Garnier and S Ciliberto



$\theta(t)$: tilt angle

$$I_{\text{eff}} \frac{d^2\theta}{dt^2} + \nu \frac{d\theta}{dt} + C\theta = M + \sqrt{2k_B T \nu} \eta,$$

$$Q_\tau = \Delta U_\tau - W_\tau$$

$$= -\frac{1}{k_B T} \int_{t_i}^{t_i+\tau} \nu \left[\frac{d\theta}{dt}(t') \right]^2 dt' + \frac{\sqrt{2k_B T \nu}}{k_B T} \int_{t_i}^{t_i+\tau} \eta(t') \frac{d\theta}{dt}(t') dt'.$$

Energetics at mesoscopic scales


Experimental demonstrations of *energy balance*

PHYSICAL REVIEW E **75**, 011122 (2007)

Experimental test of a new equality: Measuring heat dissipation in an optically driven colloidal system

Shoichi Toyabe,^{1,*} Hong-Ren Jiang,¹ Takenobu Nakamura,² Yoshihiro Murayama,¹ and Masaki Sano^{1,†}

Random Poisson switching



Harada-Sasa* identity : (*PRL '05)

$$\langle J \rangle_0 = \gamma \int_{-\infty}^{\infty} [\tilde{C}(\omega) - 2k_B T \tilde{R}'(\omega)] \frac{d\omega}{2\pi},$$

deviation from equilibrium F-D relation

f^p : probe force

$$\gamma \dot{x}(t) = F(x(t), t) + \varepsilon f^p(t) + \hat{\xi}(t),$$

heat : $\langle J \rangle = \langle F(x(t), t) \circ v(t) \rangle$,

correlation : $C(t) \equiv \langle \dot{x}(t) \dot{x}(0) \rangle_0$.

response :

$$\langle \dot{x}(t) \rangle_\varepsilon - v_s = \varepsilon \int_{-\infty}^t R(t-s) f^p(s) ds + o(\varepsilon^2),$$

Energetics at mesoscopic scales

Experimental demonstrations of *energy balance*


PRL **104**, 198103 (2010)

PHYSICAL REVIEW LETTERS

week ending
14 MAY 2010

Nonequilibrium Energetics of a Single F_1 -ATPase Molecule

Shoichi Toyabe,¹ Tetsuaki Okamoto,¹ Takahiro Watanabe-Nakayama,² Hiroshi Taketani,¹
Seishi Kudo,³ and Eiro Muneyuki^{1,*}





$$\text{heat} : J \equiv \langle [\Gamma v(t) - \xi(t)] \circ v(t) \rangle = \left\langle \frac{d'Q}{dt} \right\rangle$$

Harada-Sasa equality :

$$J = \Gamma v_s^2 + \Gamma \int_{-\infty}^{\infty} df [\tilde{C}(f) - 2T\tilde{R}'(f)],$$

Torque by electric field :



Energetics at mesoscopic scales

2nd law of stochastic energetics

$$\begin{array}{lcl} a(t_0) & = & a_i \\ \downarrow & & \\ a(t_1) & = & a_f \end{array}$$

Irreversible work : $W_{\text{irr}} \equiv W - \Delta F$

$$= \int_{a_i}^{a_f} \left\{ \frac{\partial U(x(t), a(t))}{\partial a} - \left\langle \frac{\partial U(x, a(t))}{\partial a} \right\rangle_{\text{can}; T} \right\} da(t).$$


$$F : e^{-F/k_B T} = \int e^{-U(x, a)/k_B T} dx$$

Quasi-static limit : $W_{\text{irr}} \rightarrow 0, \quad (\text{prob.1})$
 i.e. $W \rightarrow \Delta F$ *without ensemble averages*

Jarzynski equality (Jarzynski, PRL97, Crooks, JSP98)

$$1 = \langle e^{-\beta W_{\text{irr}}} \rangle_{\text{eq. at } t=0} \Rightarrow \langle W_{\text{irr}} \rangle_{\text{eq. at } t=0} \geq 0$$


Complementarity: $\langle W_{\text{irr}} \rangle \Delta t \geq S(a_i, a_f) \quad (\Delta t \rightarrow \infty)$
 (correction : initial condition sensitive)



Energetics at mesoscopic scales

REMARK : energetics of *open system* can also be formulated.



Base : *energy of the open system*



Individual particles do *not* carry the chemical potential μ .

Energetics at mesoscopic scales

RESUMÉ : “thermodynamic structure”



(cf. steady-state thermodyn., Q' m entangling,...)

Introduction

Stochastic
Energetics

Misconceptions

Misconceptions about the fluctuations

*Are Langevin equations **always** irreversible ?*

“the mesoscopic equations are therefore irreversible”

(van Kampen’s book “Stochastic Processes in Physics and Chemistry”)

Yes & No :

Yes — microscopic viewpoint

The heat bath does not accumulate the memory*.

(**temperature* [energy] & *mean-flow* [momentum])

No — **meso**scopic viewpoint

The trajectories can **describe quasi-equilibrium process**.

cf. thermodynamic adiabatic process of chaotic Hamiltonian system
(macro) (micro)


Misconceptions about the fluctuations

Numerical convergence of Langevin eq.

$$0 = -\frac{\partial U(x, a)}{\partial x} - \gamma \frac{dx}{dt} + \sqrt{2\gamma k_B T} \theta(t) \quad \text{under a given } \theta(t); \quad B_t = \int_0^t \theta(s) ds$$

1st order Euler scheme for Langevin equation ($t_0 < t < t_1$):

$$0 = -\frac{\partial U(x_t, a_t)}{\partial x} - \gamma \frac{x_{t+\Delta t} - x_t}{\Delta t} + \sqrt{2\gamma k_B T} (B_{t+\Delta t} - B_t)$$

convergence at $\mathcal{O}(\Delta t^{\frac{1}{2}})$  solution $x(t)$

But **error of $\mathcal{O}(1)$ for energy balance.**

Implication :

Apparent equilibrium state can dissipate heat.

Order $\mathcal{O}(\Delta t)$ convergence required for correct energy balance.

Misconceptions about the fluctuations


Can Carnot cycle be reversible ?

Yes & No :

Yes — macroscopic [thermodynamic] limit

No — mesoscopic case (with some exceptions)

Detachment from heat baths



* **divergence of relaxation time** → Impossibility of *quasi-static* operations

associated **irreversible work** : *non-extensive* & *arbitrarily small*

Misconceptions about the fluctuations


Can Carnot cycle be reversible ?

Yes & No :

Yes — macroscopic [thermodynamic] limit

No — **meso**scopic case (with some exceptions)

Adiabatic process



* relaxation from **non-gibbs distribution** → Irreversible information loss

associated **irreversible work** : non retrievable otherwise

Misconceptions about the fluctuations


Can Feynman's ratchet wheel and pawl attain Carnot efficiency?

"... our device cannot extract more work than this $[(T_1 - T_2)/T_1]$, operating reversibly."
— Feynman's lecture 46-3 (1963)

No :


wiggling between the "jumps"

Feynman's ratchet



\approx

mechanical *heat* conductor




"stall state" (jump frequency=0) \neq Carnot efficiency
= **null** efficiency

(Parrondo 1996, KS 1997)

Misconceptions about the fluctuations

REMARK : mechanical *heat* conductor

→ “random motion” can **absorb** heat (cf. Joule’s law)



cf. “Maxwell’s demon”

$T'=0$ case :

$$\gamma \dot{x}(t) = -K \left[x(t) - \frac{K}{\gamma'} \int_0^\infty e^{-\frac{K}{\gamma'} s} \underline{x(t-s)} ds \right] + \sqrt{2\gamma k_B T} \theta(t)$$

feed-back as 0-K bath

Introduction

Stochastic
Energetics

Misconceptions

Misconceptions about the fluctuations

Is the over-damped limit $m \rightarrow 0$ *regular* in energetics ?

under-damped : $\frac{dp}{dt} = -\frac{\partial U(x, a)}{\partial x} - \gamma \frac{dx}{dt} + \sqrt{2\gamma k_B T} \theta(t), \quad \frac{dp}{dt} = \frac{p}{m}$

time coarse graining \downarrow $\gamma[x(t + \Delta t) - x(t)] = \sqrt{2\gamma k_B T(x(t))}[B_{t+\Delta t} - B_t] - \frac{\partial U(x(t), a(t))}{\partial x} \Delta t + k_B T'(x(t))\{[B_{t+\Delta t} - B_t]^2 - \Delta t\} + o(\Delta t),$

over-dumped : $0 = -\frac{\partial U(x, a)}{\partial x} - \gamma \frac{dx}{dt} + \sqrt{2\gamma k_B T} \theta(t)$



under-damped

over-damped

Introduction


Stochastic
Energetics

Misconceptions

Misconceptions about the fluctuations

Is the over-damped limit $m \rightarrow 0$ *regular* in energetics ?

No, sometimes :




Model : $m = 0$

"Carnot's limit efficiency" $\frac{T' - T}{T'}$
at stalled state ($v=0$).

Model : $m \rightarrow 0$

divergence of heat leak $\propto \frac{1}{\sqrt{m}}$
(\Rightarrow zero efficiency)



Introduction

Stochastic
Energetics

Misconceptions

Misconceptions about the fluctuations

QUESTION :

Relation between mesoscopic heat AND “calorimetric heat” ?

Introduction


Stochastic
Energetics

Misconceptions

What is the heat ?


What is the heat ?

Example



$$U(x,a) = U_{\text{chain}}(x) + U_{\text{optic}}(x,a)$$

$$0 = -\frac{\partial U(x,a)}{\partial x} - \gamma \frac{dx}{dt} + \sqrt{2\gamma k_B T} \theta(t)$$



Introduction


Stochastic
Energetics

Misconceptions

What is the heat ?


What is the heat ?

Example



$$U(x, a) = U_{\text{chain}}(x) + U_{\text{optic}}(x, a)$$

$$0 = -\frac{\partial U(x, a)}{\partial x} - \gamma \frac{dx}{dt} + \sqrt{2\gamma k_B T} \theta(t)$$



Introduction

Stochastic
Energetics

Misconceptions

What is the heat ?


What is the heat ?

Is **mesoscopic heat** = “calorimetric heat” $d'Q_m$?

$$d'Q \equiv \left(-\gamma \frac{\partial x}{\partial t} + \sqrt{2\gamma k_B T} \theta(t) \right) \circ dx(t)$$

$$0 = -\frac{\partial U(x, a)}{\partial x} - \gamma \frac{dx}{dt} + \sqrt{2\gamma k_B T} \theta(t)$$

No in general



$U(x, a)$: free energy — entropy renormalized

$d'W$: scale independent

energy balance for calorimetric heat

➡ relation between **mesoscopic heat** AND “calorimetric heat”

Introduction

Stochastic
Energetics

Misconceptions

What is the heat ?

What is the heat ?

Is **mesoscopic heat** = “calorimetric heat” $d'Q_m$?

$$d'Q \equiv \left(-\gamma \frac{\partial x}{\partial t} + \sqrt{2\gamma k_B T} \theta(t) \right) \circ dx(t)$$

$$0 = -\frac{\partial U(x, a)}{\partial x} - \gamma \frac{dx}{dt} + \sqrt{2\gamma k_B T} \theta(t)$$

No in general

$$dE - T \frac{\partial U}{\partial T} \mapsto dE_m$$

$$dQ - T \frac{\partial U}{\partial T} \mapsto dQ_m$$

mesoscopic energy balance

$$dE = d'Q + d'W$$



calorimetric energy balance

$$dE_m = d'Q_m + d'W$$

Introduction


Stochastic
Energetics

Misconceptions


What is the heat ?

What is the heat ?

IMPLICATION : local energy transfer *in the environment*



No calorimetric heat



Introduction


Stochastic
Energetics

Misconceptions

What is the heat ?

What is the heat ?

IMPLICATION : scale cascade of heat



(controversy ~2000)

does the **filament** generate a *work* or a *heat* ?

Introduction


Stochastic
Energetics

Misconceptions

What is the heat ?

What is the heat ?

IMPLICATION : scale cascade of heat



(controversy ~2000)

does the **filament** generate a *work* or a *heat* ?

Introduction

Stochastic
Energetics

Misconceptions

Different heats

Project

Project

RÉSUMÉ :

stochastic energetics

=> new perspectives of fluctuation phenomena

What can we do with these perspectives ?

> exploration of **structure-function relation** of proteins

> **mesoscopic** mechanism of *motor proteins*

> modelling common to ***myosin*** & ***kinesin***

— similar / different

Introduction


Stochastic
Energetics

Misconceptions

Different heats

Project

Project



(Vale, J. Cell Biol. '96)

Introduction


Stochastic
Energetics

Misconceptions

Different heats

Project

Project




(Vale, J. Cell Biol. '96)

Identical QM reaction

J. Phys. Chem. B, Vol. 108, No. 10, 2004

(modified from) Li and Cui



Pi = H_2PO_4

(cf. protein kinase; γ -Pi is given to Se, which loses a proton)

Introduction


Stochastic
Energetics

Misconceptions

Different heats


Project

Project



(Vale, J. Cell Biol. '96)

Very similar 3D atomic structure



F. J. Kull, R. D.Vale and R.J.Fletcher
J. Muscle Research and Cell Motility 19, 877-886 (1998)

Introduction


Stochastic
Energetics

Misconceptions

Different heats

Project

Project



(Vale, J. Cell Biol. '96)

Very similar AA sequences

Nucleotide Binding Motifs				
	$\alpha, \beta\text{-PO}_4$	$\gamma\text{-PO}_4$	$\gamma\text{-PO}_4$	Purine
	<u>N-1</u>	<u>N-2</u>	<u>N-3</u>	<u>N-4</u>
G Proteins	GxxGxGKS/T	T	DxxG	NKxD
Kinesins	GQTxxGKS/T	NxxSSR	DxxGxE	RxRP
Myosins	GESxxGKS/T	NxxSSR	DxxGxE	NP

(Vale, J. Cell Biol. '96)

Introduction


Stochastic
Energetics

Misconceptions

Different heats

Project


Project



(Vale, J. Cell Biol. '96)

Similar folding modules

Conserved β -strands



Kull, Vale Fletterick
J.Muscle Res. Cell Motility 1998

Introduction


Stochastic
Energetics

Misconceptions

Different heats

Project


Project




(Vale, J. Cell Biol. '96)

Different chemo-mechanical cycles

kinesin - microtubule



myosin – F-actin



Introduction


Stochastic
Energetics

Misconceptions

Different heats

Project

Project



(Vale, J. Cell Biol. '96, modified)

Different filament binding

Introduction


Stochastic
Energetics

Misconceptions

Different heats

Project

Project



(Vale, J. Cell Biol. '96)

Designing principle :

— common *core mechanism* & different attachments

Introduction

Stochastic
Energetics



Misconceptions

Different heats

Project

Project

Working hypothesis :
— bidirectional control



(Vale, J. Cell Biol. '96)

2 “allosteric” couplings (sensor → gate)

Introduction

Stochastic
Energetics


Misconceptions

Different heats

Project

Project

Working hypothesis :
— bidirectional control



kinesin

myosin

(KS: 2005,2007,
work in progress)

Introduction

Stochastic
Energetics

Misconceptions

Different heats

Project

Summary

*Welcome
to
fluctuating world
and
stochastic energetics*

Sekimoto

LECTURE NOTES IN PHYSICS 799



Stochastic Energetics