

Direct measurement of the $\bar{K}N \rightarrow \pi\Sigma$ scattering
amplitude below the $\bar{K}N$ threshold employing the
 $d(K^-, N)''\pi\Sigma''$ reaction

Kentaro Inoue

February 1, 2025

Contents

1	Introduction	3
1.1	$\Lambda(1405)$ and $\bar{K}N$ interaction	3
1.2	$d(K^-, n)$ reaction	3
1.3	The J-PARC E31 experiment	3
2	Experimental setup	4
3	Analysis	5
4	Discussion	6
4.1	Decomposition of the $K^-d \rightarrow n\pi^+\pi^-n$ events	6
4.1.1	Backward $\pi^\mp\Sigma^\pm$ event selection	6
4.1.2	Template fitting	6
4.2	Conversion to the cross section	6
4.3	Spectra	6
4.3.1	Qualitative properties of obtained spectra	6
4.3.2	Comparison with theoretical calculations	6
4.3.3	$\bar{K}N$ Pole parameters assuming the 2-step reaction . .	6
4.3.4	Comparison with DCC model	6
5	Conclusion	7
A	$\pi^0\Sigma^0$ spectrum analysis	8
B	Detector resolution	9
B.1	CDC resolution	9
B.2	Detector resolution on the $d(K^-, N)$	9
C	$d(K^-, n)K^0n$ analysis	10

Chapter 1

Introduction

1.1 $\Lambda(1405)$ and $\bar{K}N$ interaction

1.2 $d(K^-, n)$ reaction

1.3 The J-PARC E31 experiment

Chapter 2

Experimental setup

Chapter 3

Analysis

Chapter 4

Discussion

4.1 Decomposition of the $K^-d \rightarrow n\pi^+\pi^-n$ events

4.1.1 Backward $\pi^\mp\Sigma^\pm$ event selection

4.1.2 Template fitting

4.2 Conversion to the cross section

4.3 Spectra

4.3.1 Qualitative properties of obtained spectra

4.3.2 Comparison with theoretical calculations

4.3.3 $\bar{K}N$ Pole parameters assuming the 2-step reaction

4.3.4 Comparison with DCC model

Chapter 5

Conclusion

Appendix A

$\pi^0\Sigma^0$ spectrum analysis

Appendix B

Detector resolution

B.1 CDC resolution

B.2 Detector resolution on the $d(K^-, N)$

Appendix C

$d(K^-, n)K^0n$ analysis

Bibliography

- [1] R.L. Workman et al. (Particle Data Group), Prog. Theor. Exp. Phys. 2022, 083C01 (2022)
"Review of Particle Physics"
- [2] R. H. Dalitz and A. Deloff, J. Phys. G17, 281 (1991).
"The Shape and Parameters of the $\Lambda(1405)$ Resonance"
- [3] M. Hassanvand et al., Phys. Rev. C **87**, 055202 (2013)
"Theoretical analysis of $\Lambda(1405) \rightarrow (\pi\Sigma)^0$ mass spectra produced in $p + p \rightarrow p + \Lambda(1405) + p$ reactions"
- [4] J. Esmaili, Y. Akaishi, and T. Yamazaki, Phys. Lett. B **686**, 23 (2010)
"Experimental confirmation of the $\Lambda(1405)$ ansatz from resonant formation of a K^-p quasi-bound state in K^- absorption by ^3He and ^4He "
- [5] R. H. Dalitz and S. F. Tuan, Phys. Rev. Lett. 2 (1959).
"Possible Resonant State in Pion-Hyperon Scattering"
- [6] M. H. Alston, L. W. Alvarez, P. Eberhard and M. L. Good, Phys. Rev. Lett. 6, 698 (1961).
"Study of Resonances of the Σ - π System"
- [7] R. J. Hemingway, Nucl Phys B **253**, 742 (1985).
"Production of $\Lambda(1405)$ in K^-p Reactions at 4.2GeV/ c "
- [8] B. Conforto et al., Nucl. Phys. B **34**, 41 (1971).
"New experimental results on the Reactions $K^-p \rightarrow \bar{K}N$ and $K^-p \rightarrow \Sigma\pi$ a partial-wave analysis between 430 and 800MeV/ c "
- [9] A. J. Van Horn, Nucl. Phys. B **87**, 145 (1975).
"Energy dependent partial-wave analysis of $K^-p \rightarrow \Lambda\pi^0$ between 1540 and 2215MeV"

- [10] R. J. Hemingway et al., Nucl. Phys. B **91**, 12 (1975).
"New data on $K^-p \rightarrow K^-p$ and K^0n and a partial-wave analysis between 1840 and 2234 MeV center of mass energy"
- [11] P. Baillon and P. J. Litchfield, Nucl. Phys. B **94**, 39 (1975).
"Energy-independent partial-wave analysis of $\bar{K}N \rightarrow \Lambda\pi$ between 1540 and 2150 MeV"
- [12] G. P. Gopal et al., Nucl. Phys. B **119**, 362 (1977).
"Partial-wave analyses of KN two-body reactions between 1480 and 2170 MeV"
- [13] W. E. Humphrey and R. R. Rose, Phys. Rev. **127**, 1305(1962).
"Low-Energy Interactions of K^- Mesons in Hydrogen"
- [14] J. K. Kim, Phys. Rev. Lett. **19**, 1074 (1967).
"Multichannel Phase-Shift Analysis of $\bar{K}N$ Interaction in the Region 0 to 550 MeV/c"
- [15] A. D. Martin, Nucl. Phys. B **179**, 33 (1981).
"Kaon-Nucleon Parameters"
- [16] J. D. Davies et al., Phys. Lett. **83B**, 55 (1979).
"Observation of kaonic hydrogen atom X-rays"
- [17] M. Izycki et al., Z. Phys. A **297**, 11 (1980).
"Results of the search for K -series X-rays from kaonic hydrogen"
- [18] P. M. Bird et al., Nucl. Phys. A **404**, 482 (1983).
"Kaonic Hydrogen atom X-rays"
- [19] M. Iwasaki et al., Phys. Rev. Lett. **78**, 3067 (1997).
"Observation of Kaonic Hydrogen K_α X Rays"
- [20] M. Bazzi et al., Phys. Lett. B **704**, 113 (2011).
"A New Measurement of Kaonic Hydrogen X-Rays"
- [21] J.C.Nacher et al., Phys. Lett. B **455**, 55 (1999).
"Photoproduction of the $\Lambda(1405)$ on the proton and nuclei"
- [22] M. Niiyama et al., Phys. Rev. C **78**, 035202 (2008).
"Photoproduction of $\Lambda(1405)$ and $\Sigma(1385)$ on the proton at $E_\gamma = 1.5$ -2.4 GeV/c"

- [23] K. Morita for the CLAS Collaboration,
Phys. Rev. C **87**, 035206 (2013).
"Measurement of the $\pi\Sigma$ photoproduction line shapes near the $\Lambda(1405)$ "
- [24] K. Morita for the CLAS Collaboration,
Phys. Rev. Lett. **112**, 082004 (2014).
"Spin and parity measurement of the $\Lambda(1405)$ baryon"
- [25] S. X. Nakamura, and D. Jido, Prog. Theor. Exp. Phys., 023D01 (2014)
"Lambda (1405) photoproduction based on the chiral unitary model"
- [26] D. Jido et al., Nucl. Phys. A **725**, 181 (2003).
"Chiral Dynamics of the Two $\Lambda(1405)$ States"
- [27] G. Agakishiev for the HADES Collaboration,
Phys. Rev C **87**, 025201 (2013).
"Baryonic Resonances to the $\bar{K}N$ threshold: The case of $\Lambda(1405)$ in pp collisions"
- [28] Y. Ikeda, T. Hyodo, and W. Weise, Nucl. Phys. A **881**, 98 (2012)
"Chiral SU(3) theory of antikaon–nucleon interactions with improved threshold constraints"
- [29] Z.-H. Guo and J. Oller, Phys. Rev. C **87**, 3, 035202 (2013)
"Meson-baryon reactions with strangeness - 1 within a chiral framework"
- [30] M. Mai and U.-G. Meißner, Eur. Phys. J. A **51**, 3, 30 (2015),
"Constraints on the chiral unitary amplitude from $\pi\Sigma K^+$ photoproduction data"
- [31] H. Zhang et al., Phys. Rev. C **88**, 035204 (2013).
"Partial-wave analysis of $\bar{K}N$ scattering reactions"
- [32] H. Zhang et al., Phys. Rev. C **88**, 035205 (2013).
"Multichannel parametrization of $\bar{K}N$ scattering amplitudes and extraction of resonance parameters"
- [33] H. Kamano et al., Phys. Rev. C **90**, 065202 (2014).
"Dynamical Coupled-Channels Model of K^-p Reactions: Determination of Partial Wave Amplitudes"
Phys. Rev. C **92**, 025205 (2015).
"Dynamical Coupled-Channels Model of K^-p Reactions."

Extraction of Λ^* and Σ^* Hyperon Resonances”
 Phys. ReV. C**95**, 044903(E) (2015).

- [34] O. Braun et al., Nucl. Phys. B **129**, 1 (1977).
 ”New Information About the Kaon-Nucleon-Hyperon Coupling Constants $g(KN\Sigma(1197))$, $g(KN\Sigma(1385))$ and $g(KN\Lambda(1405))$ ”
- [35] D. Jido, E. Oset and T. Sekihara, Eur. Phys. J. A **42**, 257 (2009).
 ”Kaonic Production of $\Lambda(1405)$ off deuteron target in chiral dynamics”
- [36] J. Yamagata-Sekihara, T. Sekihara, and D. Jido, Prog. Theor. Exp. Phys.**2013**, 043D02 (2013).
 ”Production of hyperon resonances induced by kaons on a deuteron target”
- [37] H. Noumi et al., Proposals for the 15th PAC meeting
 ”Spectroscopic study of hyperon resonances below $\bar{K}N$ threshold via the (K^-, n) reaction on Deuteron”
- [38] K. Miyagawa, J. Haidenbauer, and H. Kamada Phys. Rev. C **97**, 055209 (2018)
 ”Faddev approach to the reaction $K^-d \rightarrow \pi\Sigma n$ at $p_K = 1.0\text{GeV}/c$ ”
- [39] E. Oset, A. Ramos, and C. Bennhold, Phys. Lett. B **527**, 99 (2002); **530**, 260(E) (2002).
 ”Low lying $S = -1$ excited baryons and chiral symmetry”
- [40] A. Cieplý and J. Smejkal, Nucl. Phys. A **881**, 115 (2012).
 ”Chirally motivated $\bar{K}N$ amplitudes for in-medium applications”
- [41] S. Ohnishi et al, Phys. Rev. C **93**, 025207 (2016).
 ”Structure of the $\Lambda(1405)$ and the $K^-d \rightarrow \pi\Sigma n$ reaction”
- [42] H. Kamano et al., Phys. ReV. C**94**, 065205 (2016).
 ”Toward Establishing Low-Lying Λ and Σ Hyperon Resonances with the $\bar{K} + d \rightarrow \pi + Y + N$ Reaction”
- [43] H. Noumi for the E31 Collaboration, Phys. ReV. B**837**, 137637 (2023).
 ”Pole Position of $\Lambda(1405)$ measured in $d(K^-, n)\pi\Sigma$ reaction”
- [44] S. Kawasaki’s Doctor thesis will be submitted.
- [45] K. Agari et al, Prog. Theor. Exp. Phys., 02B009 (2012)

- [46] K. Agari et al, Prog. Theor. Exp. Phys., 02B011 (2012)
- [47] TRANSPORT <http://linac96.web.cern.ch/Linac96/Proceedings/Thursday/THP72/Paper.pdf>
- [48] T. K. Ohoka et al., Nuclear Science, IEEE Transactions on 33, 98 (1986).
- [49] M. Shiozawa and et al., A new TKO system manager board for a dead-time-free data acquisition system, in 1994 IEEE Nuclear Science Symposium-NSS'94, pages 632–635, (1994)
- [50] M. Iio et al., Nucl. Instrum. Methods Phys. Res., Sect. **A** 687, 1 (2012).
- [51] S. Agostinelli et al., Nucl. Instrum. Methods Phys. Res., Sect. **A** 506, 250 (2003)
J. Allison et al., IEEE Transactions on Phys. Sci. 53, 207 (2006)
J. Allison et al., Nucl. Instrum. Methods Phys. Res., Sect. **A** 835, 186 (2016)
- [52] K. Fuji, https://www-jlc.kek.jp/subg/offl/lib/docs/helix_manip/node3.html (1968).
- [53] Opera Electromagnetic FEA Solution Software
- [54] V. Flaminio et al., CERN-HARA-87-01, 121 (1983).
- [55] M. Jones et al, Nucl. Phys. B **90**, 349 (1975)
- [56] R. Barlow and C. Beeston, Comp. Phys. Comm. **77**, 219 (1993).
"Fitting using finite Monte Carlo samples"
- [57] A. Nappi, Comp. Phys. Comm. **180**, 269 (2009).
- [58] C.J.S. Damerell et. al., Nucl. Phys. B **129**, 397 (1977). "
 K^-n elastic scattering between 610 and 840 MeV/ c "
- [59] M. Jones, R. Levi, Setti, D. Merrill and R. D. Tripp, Phys. Rev. B **90**, 349 (1975). K^-p charge exchange and hyperon production cross section from 860 to 1000 MeV/ c
- [60] M. Bernheim and et. al., Nucl. Phys. A **365**, 349, (1981). Momentum distributions of nucleons in the deuteron from $d(e, e'p)n$ reaction
- [61] R. Machleidt, Phys. Rev. C **63**, 024001 (2001).
- [62] R. Barlow and C. Beeston, Comp. Phys. Comm. **77**, 219 (1993).

- [63] L. Lensniak, AIP Conf. Proc. 1030, 238–243 (2008). New formula for a resonant scattering near an inelastic threshold
- [64] S. Agostinelli et al., Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment **506**, 250 (2003).
J.Allison et el., Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment **835**, 186 (2016).