

## CHAPTER 2: ALGEBRAIC STRUCTURES PART 1

**Hamza Moufek**

Algebra 1

November 2024



# OUTLINES OF THIS TALK

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition

Examples

### Properties of binary operation

Commutativity

associativity

Distributivity

Identity and inverse  
elements

Identity or neutral  
elements

Inverse Element

### Groups

Definition and

Examples

Properties

### Subgroups

Definition

Characterization of  
subgroups

### Group Homo- morphism

Definition and

Examples

Types of  
homomorphisms

- Binary operations
- Groups
- Subgroups
- Group homomorphisms

# DEFINITION

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

Definition and  
Examples

Definition  
Examples

Properties of  
binary  
operation

Commutativity

Associativity

Distributivity

Identity and inverse  
elements

Identity or neutral  
elements

Inverse Element

Groups

Definition and

Examples

Properties

Subgroups

Definition

Characterization of  
subgroups

Group Homo-  
morphism

Definition and

Examples

Types of  
homomorphisms

## DEFINITION

A binary operation on a set  $S$  is a function

$$\begin{aligned} * &: S \times S \rightarrow S \\ (a, b) &\mapsto *(a, b) \end{aligned}$$

For convenience we write  $a * b$  instead of  $*(a, b)$ . The set  $S$  is said closed under the operation  $*$ . We call the data of a set  $S$  together with a binary operation  $*$ , written  $(S; *)$ , magma.

Binary operations are usually denoted by other symbols (for example,  $+$  or  $\times$  for numbers) but, for the moment, we use the generic notation  $a * b$ .

FIGURE: 1

# EXAMPLE 1

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition  
Examples

### Properties of binary operation

Commutativity  
associativity  
Distributivity  
Identity and inverse  
elements  
Identity or neutral  
elements  
Inverse Element

### Groups

Definition and  
Examples  
Properties

### Subgroups

Definition  
Characterization of  
subgroups

### Group Homo- morphism

Definition and  
Examples  
Types of  
homomorphisms

- Let  $S = \mathbb{R}$  and  $*$  be  $+$ . For  $a, b \in \mathbb{R}$ ,  $a * b = a + b$ , addition (note that  $a + b \in \mathbb{R}$ ). This is a binary operation.
- Let  $S = \mathbb{Z}$  and  $*$  be  $\times$ . For  $a, b \in \mathbb{Z}$ ,  $a * b = ab$ , multiplication (note that  $ab \in \mathbb{Z}$ ). This is a binary operation.
- Let  $S = \mathbb{Z}$  and  $a * b = \max \{a, b\}$ , the largest of  $a$  and  $b$ . This is a binary operation.
- Let  $S = \mathbb{Q}$  and define  $*$  by  $a * b = a$ . This is a binary operation.
- Let  $S = \mathbb{Z}$  and  $a * b = \frac{a}{b}$ . This is not a binary operation, as it's not defined when  $b = 0$ , and also  $\frac{a}{b}$  need not be in  $\mathbb{Z}$ .
- Let  $S = \{f : A \rightarrow A\}$ , with  $*$  composition of functions and  $A$  non empty set. This is a binary operation.
- Let  $X$  be a set. On  $\mathcal{P}(X)$ , the union of

$$\mathcal{P}(X)^2 \rightarrow \mathcal{P}(X)$$

$$(A, B) \mapsto A \cup B$$

is a binary operation. The same, for the intersection  $A \cap B$  and symmetric difference  $A \Delta B$ .

# EXAMPLE 2

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition

Examples

### Properties of binary operation

Commutativity

associativity

Distributivity

Identity and inverse  
elements

Identity or neutral  
elements

Inverse Element

### Groups

Definition and  
Examples

Properties

### Subgroups

Definition

Characterization of  
subgroups

### Group Homo- morphism

Definition and  
Examples

Types of  
homomorphisms

## EXAMPLE 2

A binary operation  $\triangle$  is defined on the set  $\mathbb{R}$  of real numbers by

$$a \triangle b = \frac{2a - b}{a^2 + 2}, \text{ where } a, b \in \mathbb{R}$$

Evaluate: (a)  $2 \triangle 3$ , (b)  $3 \triangle 2$ .

### Solution

$$(a) \quad 2 \triangle 3 = \frac{2 \cdot 2 - 3}{2^2 + 2} = \frac{1}{6}, \quad (b) \quad 3 \triangle 2 = \frac{2 \cdot 3 - 2}{3^2 + 2} = \frac{4}{11}.$$

# EXAMPLE 3

## EXAMPLE 3

The operation  $\bar{+}$  is defined on the set  $S = \{\bar{0}, \bar{1}, \bar{2}, \bar{3}, \bar{4}\}$  by Table below

$\bar{+}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$
$\bar{0}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$
$\bar{1}$	$\bar{1}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{0}$
$\bar{2}$	$\bar{2}$	$\bar{3}$	$\bar{4}$	$\bar{0}$	$\bar{1}$
$\bar{3}$	$\bar{3}$	$\bar{4}$	$\bar{0}$	$\bar{1}$	$\bar{2}$
$\bar{4}$	$\bar{4}$	$\bar{0}$	$\bar{1}$	$\bar{2}$	$\bar{3}$

**Remark** These table (Cayley table for the magma  $(\mathbb{Z}/5\mathbb{Z}, \bar{+})$ ) defines the operation  $\bar{+}$  on the set  $S$ . The set  $S$  is closed with respect to  $\bar{+}$  since when any two elements of  $S$  are combined, the result is always an element of  $S$ . That is,

$$\forall x, y \in S \Rightarrow x \bar{+} y \in S.$$

for example  $\bar{3} \bar{+} \bar{2} = \bar{0}$  and  $(\bar{3} \bar{+} \bar{1}) \bar{+} \bar{2} = \bar{1}$  an other example with operation table:

# COMMUTATIVE OPERATIONS

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition

Examples

### Properties of binary operation

#### Commutativity

associativity

Distributivity

Identity and inverse  
elements

Identity or neutral  
elements

Inverse Element

### Groups

Definition and  
Examples

Properties

### Subgroups

Definition

Characterization of  
subgroups

### Group Homo- morphism

Definition and  
Examples

Types of  
homomorphisms

## DEFINITION

A binary operation  $*$  defined on a set  $E$  is said to be commutative if  $a * b = b * a$ , for all  $a, b \in E$ .

**Remark** Addition  $(+)$  and multiplication  $(\times)$  are both commutative since for all  $a, b \in \mathbb{R}$ ,  $a + b = b + a$  and  $a \times b = b \times a$ .

The union  $(\cup)$  and the intersection  $(\cap)$  are both commutative since for any two sets  $A$  and  $B$ ,  $A \cup B = B \cup A$  and  $A \cap B = B \cap A$ .

### EXAMPLE 1

The operation  $*$  is defined on the set of real numbers  $\mathbb{R}$  by  $a * b = a + b + 2ab$ . Show that  $*$  is commutative.

$$a * b = a + b + ab \dots (1)$$

$$b * a = b + a + ba$$

$$= a + b + ab \dots (2) \text{ [since } a + b = b + a \text{ and } ab = ba \text{ for all } a, b \in \mathbb{R} \text{]}$$

Comparing (1) and (2) it follows that  $a * b = b * a$  for all  $a, b \in \mathbb{R}$ . Hence the operation  $*$  is commutative.

### EXAMPLE 2

The operation  $\nabla$  is defined on the set  $\mathbb{R}$  of real numbers by  $x \nabla y = x - y + 3xy$ .

$$x \nabla y = x - y + 3xy \dots (1)$$

$$y \nabla x = y - x + 3yx \dots (2)$$

Since subtraction is not commutative it follows that  $x - y \neq y - x$  and therefore  $x \nabla y \neq y \nabla x$ . Hence the operation  $\nabla$  is not commutative.



# ASSOCIATIVITY

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition  
Examples

### Properties of binary operation

Commutativity  
**associativity**  
Distributivity  
Identity and inverse  
elements  
Identity or neutral  
elements  
Inverse Element

### Groups

Definition and  
Examples  
Properties

### Subgroups

Definition  
Characterization of  
subgroups

### Group Homo- morphism

Definition and  
Examples  
Types of  
homomorphisms

## DEFINITION

A binary operation  $*$  defined on a closed set  $E$  is said to be associative if for every  $a, b, c, \in E, (a * b) * c = a * (b * c) = a * b * c$ .

## EXAMPLE 1

Show that the operation  $*$  defined over the set of real numbers  $\mathbb{R}$  is associative, where  $a * b = a + b + ab$ .

$$\begin{aligned}(a * b) * c &= (a + b + ab) * c \\ &= (a + b + ab) + c + (a + b + ab)c \\ &= a + b + c + ab + ac + bc + abc \dots \dots \dots (1)\end{aligned}$$

$$\begin{aligned}a * (b * c) &= a * (b + c + bc) \\ &= a + (b + c + bc) + a(b + c + bc) \\ &= a + b + c + bc + ab + ac + abc \dots \dots \dots (2)\end{aligned}$$

From (1) and (2),  $(a * b) * c = a * (b * c)$  for all  $a, b, c \in \mathbb{R}$ . Hence the operation  $*$  is associative

**exercise** The operation defined on the set  $\mathbb{R}$  of real numbers by  $a \triangle b = ab + a$ . Determine whether or not the operation  $\triangle$  is associative.

## DEFINITION

Let  $*$  and  $\triangle$  be two binary operations both defined on the closed set  $E$ . We say the operation  $*$  is distributive over  $\triangle$  if for all

$$a, b, c \in E, a * (b \triangle c) = (a * b) \triangle (a * c) \text{ and } (b \triangle c) * a = (b * a) \triangle (c * a)$$

## EXAMPLE

- For example, multiplication ( $\times$ ) is distributive over addition ( $+$ ) since for all  $a, b, c \in \mathbb{R}$ ,  $a \times (b + c) = (a \times b) + (a \times c)$  and  $(b + c) \times a = (b \times a) + (c \times a)$  written as  $a(b + c) = ab + ac$ .
- In the set  $\mathcal{P}(X)$  of parts of a set  $X$ , the union is distributive with respect to the intersection and the intersection is distributive with respect to the union

$$\forall A, B, C \in \mathcal{P}(X) \quad A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$\forall A, B, C \in \mathcal{P}(X) \quad A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

# IDENTITY OR NEUTRAL ELEMENTS

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition

Examples

### Properties of binary operation

Commutativity

associativity

Distributivity

Identity and inverse  
elements

**Identity or neutral  
elements**

Inverse Element

### Groups

Definition and

Examples

Properties

### Subgroups

Definition

Characterization of  
subgroups

### Group Homo- morphism

Definition and

Examples

Types of  
homomorphisms

## DEFINITION

Let  $*$  be a binary operation on a non empty set  $A$ . An element  $e$  is called an identity element with respect to  $*$  if

$$e * x = x = x * e$$

for all  $x \in A$ .

## Examples

- 1 is an identity element for multiplication on the integers.
- 0 is an identity element for addition on the integers.
- If  $*$  is defined on  $\mathbb{Z}$  by  $x * y = x + y + 1$  Then  $-1$  is the identity.
- The operation  $*$  defined on  $\mathbb{Z}$  by  $x * y = 1 + xy$  has no identity element.

# IDENTITY OR NEUTRAL ELEMENTS

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

Definition and  
Examples

Definition  
Examples

Properties of  
binary  
operation

Commutativity

associativity

Distributivity

Identity and inverse  
elements

**Identity or neutral  
elements**

Inverse Element

Groups

Definition and

Examples

Properties

Subgroups

Definition

Characterization of  
subgroups

Group Homo-  
morphism

Definition and

Examples

Types of  
homomorphisms

## EXERCISE 1

The operation  $*$  is defined over the set  $\mathbb{R}$  of real numbers by  $a * b = a + b + 2ab$ . Find the identity element under the operation  $*$ .

**Solution** Let  $e \in \mathbb{R}$  be the identity element under the operation  $*$ . It then follows that for every element  $a \in \mathbb{R}$ ,  $a * e = e * a = a$ . Solving for  $e$  we have

$$\begin{aligned} a * e = a &\Rightarrow a + e + 2ae = a \\ &\Rightarrow e(1 + 2a) = 0 \\ &\Rightarrow e = 0 \end{aligned}$$

Hence the identity element under  $*$  is 0.

# INVERSE ELEMENT

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition  
Examples

### Properties of binary operation

Commutativity  
associativity  
Distributivity  
Identity and inverse  
elements  
Identity or neutral  
elements

### Inverse Element

### Groups

Definition and  
Examples  
Properties

### Subgroups

Definition  
Characterization of  
subgroups

### Group Homo- morphism

Definition and  
Examples  
Types of  
homomorphisms

## DEFINITION

Let  $G$  be a set equipped with a binary operation  $*$  that admits an identity element  $e$ . We say that an element  $x \in G$  is invertible if there exists an element  $y \in G$  such that:

$$x * y = y * x = e$$

We say then that  $y$  is the inverse of  $x$ .

## Remark

- When the binary operation is denoted additively:  $+$  (resp. multiplicatively:  $\times$ ), the identity element will be denoted by  $0$  (resp.  $1$ )
- the inverse of  $x$  will be denoted by  $-x$  (resp.  $x^{-1}$ ).

## EXAMPLE

- Consider the operation of addition on the integers. For any integer  $a$ , the inverse of  $a$  with respect to addition is  $-a$ .
- Consider the operation of multiplication on  $\mathbb{Z}$ . The invertible elements are 1 and  $-1$ .
- In  $\mathbb{R}$ , each element  $a$  has an inverse for addition which is its opposite  $-a$ . But  $a$  has only an inverse for multiplication if it is non-zero; its inverse is then the inverse  $1/a$  of  $a$ .
- In  $\mathcal{F}(X, X)$ , an element  $f$  has an inverse (for the composition  $\circ$ ) if and only if  $f$  is bijective and then its inverse is the reciprocal map  $f^{-1}$ .

# INVERSE ELEMENT

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition

Examples

### Properties of binary operation

Commutativity

associativity

Distributivity

Identity and inverse  
elements

Identity or neutral  
elements

**Inverse Element**

### Groups

Definition and

Examples

Properties

### Subgroups

Definition

Characterization of  
subgroups

### Group Homo- morphism

Definition and

Examples

Types of  
homomorphisms

# INVERSE ELEMENT

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition  
Examples

### Properties of binary operation

Commutativity  
associativity  
Distributivity  
Identity and inverse  
elements  
Identity or neutral  
elements

### Inverse Element

### Groups

Definition and  
Examples  
Properties

### Subgroups

Definition  
Characterization of  
subgroups

### Group Homo- morphism

Definition and  
Examples  
Types of  
homomorphisms

## EXERCISE 2

The operation  $*$  is defined over the set  $\mathbb{R}$  of real numbers by

$$a * b = a + b + 2ab, \text{ for all } a, b \in \mathbb{R}.$$

Find the inverse under  $*$  of a general element  $a \in \mathbb{R}$  and state which element has no inverse. Determine the inverses of 2 and 3.

For  $a \in \mathbb{R}$ , let  $a^{-1} \in \mathbb{R}$  be the inverse. From Exercise 1, the identity element is  $e = 0$  so :

$$a * a^{-1} = a^{-1} * a = e = 0$$

$$a + a^{-1} + 2aa^{-1} = 0$$

$$a^{-1} + 2aa^{-1} = -a$$

$$a^{-1}(1 + 2a) = -a$$

$$a^{-1} = \frac{-a}{1 + 2a}$$

The inverse expression  $a^{-1}$  becomes undefined when the denominator is zero. so:

$$a^{-1} \text{ is not defined } \Leftrightarrow 1 + 2a = 0$$

Hence  $\frac{-1}{2}$  has no inverse. The inverse of 2 is  $\frac{-2}{5}$  and the inverse of 3 is  $\frac{-3}{7}$ .



## DEFINITION OF GROUP

Let  $G$  be a non-empty set and  $*$  a binary operation on  $G$ . We call  $(G, *)$  a group if the following hold:

- The operation  $*$  is associative. That is, for all  $a, b, c \in G$  we have that  $(a * b) * c = a * (b * c)$ .
- The operation  $*$  admits an identity element  $e \in G$ . That is, there is an  $e \in G$  so that for all  $a \in G$  we have that  $e * a = a * e = a$ .
- Every element of  $G$  is invertible. That is, for every  $a \in G$  there is an  $\tilde{a} \in G$  so that  $a * \tilde{a} = \tilde{a} * a = e$ .

Moreover, a group is called commutative or Abelian group iff  $*$  is commutative.

# EXAMPLES

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition  
Examples

### Properties of binary operation

Commutativity  
associativity  
Distributivity  
Identity and inverse  
elements  
Identity or neutral  
elements  
Inverse Element

### Groups

Definition and  
Examples  
Properties

### Subgroups

Definition  
Characterization of  
subgroups

### Group Homo- morphism

Definition and  
Examples  
Types of  
homomorphisms

- $(\mathbb{Z}, +), (\mathbb{Q}, +), (\mathbb{R}, +)$  and  $(\mathbb{C}, +)$  are commutative groups.
- $(\mathbb{N}, +)$  and  $(\mathbb{Z}, \times)$  are not groups.
- Let  $G$  be the set of bijective functions from  $A \rightarrow A$ , and let  $\circ$  be the operation of composition of functions. Then  $(G, \circ)$  is a non commutative group. With neutral element:  $id_A$ , inverse element of  $f$  is  $f^{-1}$
- Let  $\mathbb{Z}/5\mathbb{Z} = \{\bar{0}, \bar{1}, \bar{2}, \bar{3}, \bar{4}\}$  be the set of integers modulo 5. Set

$$\forall \bar{x}, \bar{y}, \overline{\bar{x} + \bar{y}} = \overline{\bar{x}} + \overline{\bar{y}}$$

This operation is well defined, and  $(G, +)$  is an abelian group.

- Let  $G$  be the set  $\{1, -1\}$ . Then it forms an abelian group under multiplication.
- The subset  $\{1, -1, i, -i\}$  of the complex numbers is a group under complex multiplication. Note that  $-1$  is its own inverse, where as the inverse of  $i$  is  $-i$ , and vice versa.

## THEOREME

Let  $(G, *)$  be a group. Then we have:

- The identity element is unique.
- For all  $a, b, x \in G$ , we have the cancellation laws

$$a * x = b * x \Rightarrow a = b,$$

and

$$x * a = x * b \Rightarrow a = b.$$

- For all  $x \in G$ , the inverse of  $x$  is unique.
- For all  $x \in G$ , the inverse of  $x^{-1}$  is  $x$ .
- For all  $x, y \in G$ ,  $(x * y)^{-1} = y^{-1} * x^{-1}$ .

# EXPONENTIATION

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition

Examples

### Properties of binary operation

Commutativity

associativity

Distributivity

Identity and inverse  
elements

Identity or neutral  
elements

Inverse Element

### Groups

Definition and

Examples

Properties

### Subgroups

Definition

Characterization of  
subgroups

### Group Homo- morphism

Definition and

Examples

Types of  
homomorphisms

## DEFINITION AND PROPERTIES

Let  $G$  be a group with binary operation  $*$ . For any  $a \in G$  we define *nonnegative integral exponents* by

$$a^0 = e, \quad a^1 = a, \quad a^{n+1} = a^n * a \quad n > 0.$$

Negative integral exponents are defined by

$$a^{-n} = (a^{-1})^n \quad n > 0.$$

With

$$a^n = \underbrace{a * a * a * \dots * a}_{n \text{ times}}$$

Laws of Exponents

$$1) x^n \cdot x^{-n} = e, \quad 2) x^m \cdot x^n = x^{m+n} \quad 3) (x^m)^n = x^{mn}$$

and If  $G$  is abelian then

$$4) (xy)^n = x^n y^n.$$

# DEFINITION

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition  
Examples

### Properties of binary operation

Commutativity  
associativity  
Distributivity  
Identity and inverse  
elements  
Identity or neutral  
elements  
Inverse Element

### Groups

Definition and  
Examples  
Properties

### Subgroups

Definition  
Characterization of  
subgroups

### Group Homo- morphism

Definition and  
Examples  
Types of  
homomorphisms

## DEFINITION 1

If a subset  $\mathbf{H}$  of a group  $\mathbf{G}$  is itself a group under the operation of  $\mathbf{G}$ , we say that  $\mathbf{H}$  is a subgroup of  $\mathbf{G}$

**Remark** If  $(G, *)$  is a group and  $H \subseteq G$  with  $(H, *)$  is also a group, then  $H$  is a subgroup of a group  $G$

## DEFINITION 2

Let  $(G, *)$  be a group. A subgroup of  $G$  is a subset  $H \subseteq G$  that satisfies the following:

- $e_G \in H$
- $\forall x, y \in H, x * y \in H.$
- $\forall x \in H, x^{-1} \in H.$

**Remark** A subgroup is a group under the induced binary operation, with the same identity element.

# CHARACTERIZATION OF SUBGROUPS AND EXAMPLE

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

Definition and  
Examples

Definition  
Examples

Properties of  
binary  
operation

Commutativity  
associativity  
Distributivity  
Identity and inverse  
elements  
Identity or neutral  
elements  
Inverse Element

Groups

Definition and  
Examples  
Properties

Subgroups

Definition  
Characterization of  
subgroups

Group Homo-  
morphism

Definition and  
Examples  
Types of  
homomorphisms

## THEOREME

A non void subset  $H$  of a group  $(G, *)$  is a subgroup of  $G$  iff :

$$\forall (a, b) \in H^2, a * b^{-1} \in H.$$

## Examples

■  $(\mathbb{Z}, +)$ ,  $(\mathbb{Q}, +)$  and  $(\mathbb{R}, +)$  are subgroups of  $(\mathbb{C}, +)$

■  $(]0, +\infty[, \times)$  is a subgroup of  $(\mathbb{R}^*, \times)$ .

## Exercise 1

Let  $G$  group. Let

$$Z(G) = \{x \in G \mid x * g = g, \text{ for all } g \in G\}$$

Show that  $Z(G)$  is a subgroup of  $G$ .

## Exercise 2

Show that intersection of two subgroups of a group  $G$  is a subgroup of  $G$ .

# UNION OF TWO SUBGROUPS

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

Definition and  
Examples

Definition  
Examples

Properties of  
binary  
operation

Commutativity

associativity

Distributivity  
Identity and inverse  
elements

Identity or neutral  
elements

Inverse Element

Groups

Definition and

Examples

Properties

Subgroups

Definition

Characterization of  
subgroups

Group Homo-  
morphism

Definition and

Examples

Types of  
homomorphisms

## Exercise

show that the union of two subgroups is not a subgroup

## proof

Let

$$H_2 = \{2n | n \in \mathbb{Z}\}$$

and

$$H_3 = \{3n | n \in \mathbb{Z}\}$$

Wher  $(\mathbb{Z}, +)$  is the group of integers .  $H_2$  and  $H_3$  will be subgroups of  $\mathbb{Z}$ . Indeed

$$2n - 2m = 2(n - m) \in H_2$$

and

$$3n - 3m = 3(n - m) \in H_3$$

Now  $H_2 \cup H_3$  is not a subgroup as  $2, 3 \in H_2 \cup H_3$  but

$$2 - 3 \notin H_2 \cup H_3$$

**THE union of two subgroups is a subgroup iff one of them is contained in the other**

# DEFINITION

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

Definition and  
Examples

Definition  
Examples

Properties of  
binary  
operation

Commutativity  
associativity

Distributivity

Identity and inverse  
elements

Identity or neutral  
elements

Inverse Element

Groups

Definition and  
Examples  
Properties

Subgroups

Definition

Characterization of  
subgroups

Group Homo-  
morphism

Definition and  
Examples

Types of  
homomorphisms

## DEFINITION

Let  $(G, *)$  and  $(G', \diamond)$  be two groups. A map  $f : G \rightarrow G'$  is said to be a group homomorphism if

$$\forall x, x' \in G \quad \phi(x * x') = \phi(x) \diamond \phi(x')$$

Then,  $\phi$  is group homomorphism .

## REMARK

Not every function from one group to another is a homomorphism! The condition  $\phi(a * b) = \phi(a) \diamond \phi(b)$  means that the map  $\phi$  preserves the structure of  $G$ .



# EXAMPLES

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition  
Examples

### Properties of binary operation

Commutativity  
associativity  
Distributivity  
Identity and inverse  
elements  
Identity or neutral  
elements  
Inverse Element

### Groups

Definition and  
Examples  
Properties

### Subgroups

Definition  
Characterization of  
subgroups

### Group Homo- morphism

Definition and  
Examples

Types of  
homomorphisms

## EXAMPLE 1

Let  $G$  be the group  $(\mathbb{R}, +)$  and  $G'$  be the group  $(\mathbb{R}_+^*, \times)$ . The map  $f : \mathbb{R} \rightarrow \mathbb{R}_+^*$  defined by  $f(x) = e^x$  is an homomorphism.

Indeed, we have

$$\forall x, y \in \mathbb{R}, f(x + y) = e^{x+y} = e^x \times e^y = f(x) \times f(y).$$

## EXAMPLE 2

Consider the function

$$\begin{aligned} \varphi : (\mathbb{Z}, +) &\rightarrow (\mathbb{Z}/5\mathbb{Z}, \overline{+}) \\ x &\mapsto \overline{x} \end{aligned}$$

for all  $(a, b) \in \mathbb{Z}$ , we have  $\varphi(a + b) = \overline{a + b} = \overline{a} + \overline{b} = \varphi(a) \overline{+} \varphi(b)$

# TYPES OF HOMOMORPHISMS

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition  
Examples

### Properties of binary operation

Commutativity  
associativity  
Distributivity  
Identity and inverse  
elements  
Identity or neutral  
elements  
Inverse Element

### Groups

Definition and  
Examples  
Properties

### Subgroups

Definition  
Characterization of  
subgroups

### Group Homo- morphism

Definition and  
Examples

### Types of homomorphisms

- A homomorphism that is both injective and surjective is an isomorphism.
- Two groups are isomorphic if there exists an isomorphism between them .
- A homomorphism from a group to itself is called an endomorphism.
- An automorphism is an isomorphism from a group to itself .

# TWO BASIC PROPERTIES OF HOMOMORPHISMS

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition  
Examples

### Properties of binary operation

Commutativity  
associativity  
Distributivity  
Identity and inverse  
elements  
Identity or neutral  
elements  
Inverse Element

### Groups

Definition and  
Examples  
Properties

### Subgroups

Definition  
Characterization of  
subgroups

### Group Homo- morphism

Definition  
Examples

Types of  
homomorphisms

## PROPOSITION

Let  $\phi : G \longrightarrow H$  be a homomorphism. Denote the identity of  $G$  by  $1_G$ , and the identity of  $H$  by  $1_H$ .

- $\phi(1_G) = 1_H$  “ $\phi$  sends the identity to the identity”
- $\phi(g^{-1}) = \phi(g)^{-1}$  “ $\phi$  sends inverses to inverses”

## EXAMPLE

Let  $G$  be the group  $(\mathbb{R}, +)$  and  $G'$  be the group  $(\mathbb{R}_+^*, \times)$ . The map  $f : \mathbb{R} \mapsto \mathbb{R}_+^*$  defined by  $f(x) = e^x$ . We have  $f(0) = 1$ . The identity element of  $(\mathbb{R}, +)$  has as its image the identity element of  $(\mathbb{R}_+^*, \times)$ . For  $x \in \mathbb{R}$  its inverse in  $(\mathbb{R}, +)$  is here its opposite  $-x$ , Then  $f(-x) = e^{-x} = \frac{1}{e^x} = \frac{1}{f(x)}$  is the inverse of  $f(x)$  in  $(\mathbb{R}_+^*, \times)$ .

## PROPOSITION

- Let  $\varphi : G \rightarrow G'$  and  $\phi : G' \rightarrow G''$  be two groups homomorphisms. The  $\phi \circ \varphi : G \rightarrow G''$  is a group homomorphism.
- If  $\varphi : G \rightarrow G'$  is a bijective homomorphism, then  $\varphi^{-1} : G' \rightarrow G$  is also a group homomorphism.

# GROUP HOMOMORPHISM

## Chapter 2: Algebraic structures part 1

Dr. Hamza  
Moufek

### Definition and Examples

Definition  
Examples

### Properties of binary operation

Commutativity  
associativity  
Distributivity  
Identity and inverse  
elements  
Identity or neutral  
elements  
Inverse Element

### Groups

Definition and  
Examples  
Properties

### Subgroups

Definition  
Characterization of  
subgroups

### Group Homo- morphism

Definition and  
Examples

Types of  
homomorphisms

## DEFINITION

Given  $\varphi : G \rightarrow G'$  a homomorphism of groups, we define the **kernel** of  $\varphi$  to be:

$$\text{Ker}(\varphi) = \{x \in G \mid \varphi(x) = e_{G'}\}$$

We define the image of  $\varphi$  to be:

$$\text{Im}(\varphi) = \{y \in G' \mid \exists x \in G \text{ such that } \varphi(x) = y\}$$

## PROPOSITION

Let  $\varphi : G \rightarrow G'$  be a group homomorphism. Then we have :

- The **kernel** of  $\varphi$ ,  $\text{ker}(f)$  is a subgroup of  $G$ .
- The **image** of  $\varphi$ ,  $\text{Im}(f)$  is a subgroup of  $G'$ .
- The homomorphism  $\varphi$  is injective if, and only if,  $\text{ker}(\varphi) = \{e_G\}$ .
- $\varphi$  is surjective if, and only if,  $\text{Im}(\varphi) = G'$ .