

1.3. Schedule of Activities (SoA)

The Schedule of Activities described below should be followed for all participants enrolled in Study GZGI. However, for those participants whose participation in this study is affected by exceptional circumstances, such as pandemics or natural disasters, please refer to Section 10.11 (Appendix 11) for additional guidance.

Study Period I Screening/Lead-in			Study Period II Treatment Period For early terminations that occur before the last visit in treatment period, see the activities listed for ET in this table. Shaded columns represent the dose escalation period Weeks 0-16.																	Study Period III Safety Follow-Up
	Screening	Lead -In																		
Visit Number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	ET	801
Week Relative to Randomization	-2	-1	0	1	2	4	6	8	10	12	13	16	20	24	26	28	32	36	-	2 Wks Post End of TXP
Allowable Interval Tolerance (days)			-	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	-	±3
Fasting Visit	X		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Administrative																				
Informed consent	X																			
Inclusion and exclusion criteria review	X		X																	
Demographics	X																			
Preexisting conditions and medical history	X		X																	
Concomitant medications	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Substance use (alcohol, caffeine, tobacco use)	X																			
Adverse events (AEs)	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Physical Evaluation																				
Height	X																			
Weight	X		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Waist circumference	X		X			X		X		X		X	X		X		X	X	X	X

Study Period I Screening/Lead-in			Study Period II Treatment Period For early terminations that occur before the last visit in treatment period, see the activities listed for ET in this table. Shaded columns represent the dose escalation period Weeks 0-16.																		Study Period III Safety Follow-Up
	Screening	Lead -In																			
Visit Number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	ET	801	
Week Relative to Randomization	-2	-1	0	1	2	4	6	8	10	12	13	16	20	24	26	28	32	36	-	2 Wks Post End of TXP	
Allowable Interval Tolerance (days)			-	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	-	±3	
Fasting Visit	X		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Vital signs (2 sitting BP and PR measurements)	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
	Vital signs include pulse rate, blood pressure, and temperature. Vital sign measurements should be collected after participant has been sitting for at least 5 minutes and before obtaining an ECG tracing and before collection of blood samples for laboratory testing (see Section 10.7).																				
Physical examination	X																				
Symptom-directed physical examination			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
12-Lead ECG	X		X			X		X		X			X		X			X	X		
	Single central ECGs will be performed at screening and early termination. Central ECGs should be collected in triplicate at all other visits. ECG should be collected prior to collection of blood samples, including PK samples. ECG measurements should be obtained per the instructions in Section 8.2.3.																				
ABPM training		X													X						
Initiate ABPM device		X													X						
Return ABPM device		X													X						
Record ABPM measurements		X													X						
	At the time participant visit data are uploaded by the site, the ABPM Data Validity Report will be generated and should be reviewed to ensure ≥70% of readings are valid. If <70% of readings are valid, the 24-hour monitoring session should be repeated within the visit window.																				
Participant Education																					
Explain diet and physical activity plan	X	X	X				X				X										
May be performed more frequently based on usual site practices and participant needs.																					

Study Period I Screening/Lead-in			Study Period II Treatment Period For early terminations that occur before the last visit in treatment period, see the activities listed for ET in this table. Shaded columns represent the dose escalation period Weeks 0-16.																		Study Period III Safety Follow-Up
	Screening	Lead -In																			
Visit Number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	ET	801	
Week Relative to Randomization	-2	-1	0	1	2	4	6	8	10	12	13	16	20	24	26	28	32	36	-	2 Wks Post End of TXP	
Allowable Interval Tolerance (days)			-	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	-	±3	
Fasting Visit	X		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Discuss diet and physical activity progress				X		X				X				X				X			
	May be performed based on participant needs.																				
Participant Administration Log																					
Dispense study drug administration log, instruct in use			X	X	X	X	X	X	X	X		X	X	X		X	X				
	The date and time of all dose administrations will be recorded in the log by the participant the day prior to a study visit, but may be recorded more frequently depending on site usual practice.																				
Study drug administration log return and review				X	X	X	X	X	X	X		X	X	X	X	X	X	X	X		
Patient-Reported Outcomes																					
SF-36v2 acute form			X									X			X			X	X		
IWQOL-Lite CT			X									X			X			X	X		
PGIS for Physical Activity			X									X			X			X	X		
Participant Survey															X				X		
	The Participant Survey should be administered at early termination if it occurs before 26 weeks.																				
Clinician-Administered Assessments (Paper)																					
Patient Health Questionnaire-9 (PHQ-9)	X		X									X			X			X	X	X	

Study Period I Screening/Lead-in			Study Period II Treatment Period For early terminations that occur before the last visit in treatment period, see the activities listed for ET in this table. Shaded columns represent the dose escalation period Weeks 0-16.																		Study Period III Safety Follow-Up
	Screening	Lead -In																			
Visit Number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	ET	801	
Week Relative to Randomization	-2	-1	0	1	2	4	6	8	10	12	13	16	20	24	26	28	32	36	-	2 Wks Post End of TXP	
Allowable Interval Tolerance (days)			-	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	-	±3	
Fasting Visit	X		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
C-SSRS screening/baseline	X																				
C-SSRS since last visit			X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Laboratory Tests and Sample Collections																					
Hematology	X		X							X					X			X	X		
Hemoglobin A1c (HbA1c)	X		X							X					X			X	X		
Clinical chemistry	X		X			X				X					X			X	X	X	
Lipid panel	X		X			X				X					X			X	X		
Urinalysis	X									X					X			X	X		
Serum pregnancy	X																				
Follicle-stimulating hormone (FSH)	X																				
	Collect serum estradiol, FSH, and LH in women whose menopausal status needs to be determined. For participants known to be either premenopausal or postmenopausal, these tests do not need to be collected.																				
Luteinizing hormone (LH)	X																				
Estradiol	X																				
Glucagon			X												X						
C-peptide			X												X						
Calcitonin	X		X							X					X			X	X		
Pancreatic amylase	X		X							X					X			X	X		

Study Period I Screening/Lead-in			Study Period II Treatment Period For early terminations that occur before the last visit in treatment period, see the activities listed for ET in this table. Shaded columns represent the dose escalation period Weeks 0-16.																		Study Period III Safety Follow-Up
	Screening	Lead -In																			
Visit Number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	ET	801	
Week Relative to Randomization	-2	-1	0	1	2	4	6	8	10	12	13	16	20	24	26	28	32	36	-	2 Wks Post End of TXP	
Allowable Interval Tolerance (days)			-	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	-	±3	
Fasting Visit	X		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Lipase	X		X							X					X			X	X		
HIV screening tests	X																				
Hepatitis C Virus (HCV) screening tests	X																				
Hepatitis B Virus (HBV) screening tests	X																				
Pro-C3			X												X			X			
High sensitivity CRP			X															X			
Apo B and Apo C3			X															X			
Leptin			X															X			
Cytokeratin 18			X															X			
eGFR (CKD-EPI)	X									X					X			X			
Urinary albumin/creatinine ratio (UACR)	X									X					X			X			
Pharmacokinetic (PK) sample	Predose		X					X		X			X					X			
	Postdose					X		X				X			X				X		
PK predose sample should be collected predose (~up to 1 hour predose). PK postdose sample time windows: 3-6 hours postdose (Week 4); 6-12 hours postdose (Week 8), 1-3 hours postdose (Week 16), 3-6 hours postdose (Week 26). Participants may need to return to clinical site for additional PK-specific visits to provide postdose PK samples.																					

Study Period I Screening/Lead-in			Study Period II Treatment Period For early terminations that occur before the last visit in treatment period, see the activities listed for ET in this table. Shaded columns represent the dose escalation period Weeks 0-16.																		Study Period III Safety Follow-Up
	Screening	Lead -In																			
Visit Number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	ET	801	
Week Relative to Randomization	-2	-1	0	1	2	4	6	8	10	12	13	16	20	24	26	28	32	36	-	2 Wks Post End of TXP	
Allowable Interval Tolerance (days)			-	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	-	±3	
Fasting Visit	X		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Stored Samples																					
Exploratory biomarker samples			X			X						X			X			X			
Randomization and Dosing																					
IWRS	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Randomization			X																		
Dispense study intervention capsules			X	X	X	X	X	X	X	X		X	X	X		X	X				
			One bottle of capsules is to be dispensed at Visit 3 and at Visit 4. Two bottles of capsules are to be dispensed at Visits 5-9 (during the rest of the dose escalation period). The site pharmacy should clearly label which bottle to take first. At Visits 10-17, 4 bottles of capsules should be dispensed.																		
Participant returns unused study drug				X	X	X	X	X	X	X		X	X	X		X	X	X			
Assess drug compliance				X	X	X	X	X	X	X		X	X	X		X	X	X	X		

Abbreviations: ABPM = ambulatory blood pressure monitoring; AEs = adverse events; BP = blood pressure; C-SSRS = Columbia Suicide Severity Rating Scale; CKD-EPI = chronic kidney disease epidemiology; ECG = electrocardiogram; eGFR = estimated glomerular filtration rate; ET = early termination; FSH = follicle-stimulating hormone; HbA1c = hemoglobin A1c; HBC = hepatitis B virus; HCV = hepatitis C virus; IWQOL-Lite- CT = Impact of Weight on Quality of Life-Lite Clinical Trials Version; IWRS = interactive web-response systems; PGIS for Physical Activity = Patient Global Impression of Status for Physical Activity; PHQ-9 = Patient Health Questionnaire-9; PK = pharmacokinetic; SF-36v2 acute form = Short Form 36 Version 2 health survey; PR = pulse rate; TXP = treatment; UACR = urinary albumin creatinine ratio; Wks = weeks