

1.3 SCHEDULE OF ACTIVITIES (SOA)

Evaluation	Screening (up to 28 days before Day 1)		Treatment Phase ^a			End of Treatment (EOT)	Post Treatment Follow-up Phase			Notes
										^a A cycle is 21 days
	Cycle 1			Cycle 2 and Beyond				Safety follow-up Period		Survival follow-up
D- 28 to D- 15	D-14 to D-1	D1 (±1)	D8 (± 1)	D15 (±1)	D1 (± 2)	30 (±7) days after last IMPs admin	At 60 (±7) days after last IMPs admin	At 90 (±7) days after last IMPs admin	Every 90 days (±7) after last safety follow-up	
Informed consent/ Inclusion and exclusion criteria	X									Informed Consent: Informed consent may be signed prior to D -28.
Demography, Medical/Surgical and Disease History	X									
Physical examination		X (^{<} 7days prior to first dose)		X ^C	X ^C	X	X	X		Section 8.2.1
Height (at baseline only) /Weight/ ECOG (HCC,SCCHN,EOC) or Karnofsky PS (GBM)	X		X	X ^C	X ^C	X	X	X		Section 8.2.1
Vital Signs	X		X	X ^C	X ^C	X	X	X		See Section 8.2.2
Resting O2 saturation for SCCHN	X		X	X ^C	X ^C	X	X			See Section 8.2.2

Evaluation	Screening (up to 28 days before Day 1)		Treatment Phase ^a			End of Treatment (EOT)	Post Treatment Follow-up Phase			Notes	
									^a A cycle is 21 days		
	Cycle 1			Cycle 2 and Beyond				Safety follow-up Period		Survival follow-up	
	D- 28 to D- 15	D-14 to D-1	D1 (±1)	D8 (± 1)	D15 (±1)	D1 (± 2)	30 (±7) days after last IMPs admin	At 60 (±7) days after last IMPs admin	At 90 (±7) days after last IMPs admin	Every 90 days (±7) after last safety follow-up	
12-Lead ECG	X					As clinically indicated					See Section 8.2.3
Laboratory Assessments											
Pregnancy test (WOCBP only) ^b		X (within 7 days prior to first dose)				X	X	X	X (every 30±7 days until 5 months after last dose of study treatment)		^b Women of child bearing potential must have a negative serum pregnancy test result within 7 days prior to first IMP administra- tion.
Blood Chemistry		X	X	X ^c	X ^c	X	X	X	X		See Section 10.3 Table 12
Hematology		X	X	X ^c	X ^c	X	X	X	X		Section 10.3
Coagulation(GBM)		X	X	X ^c	X ^c	X	X				Section 10.3
Coagulation (for HCC, SCCHN, and EOC)		X				As clinically indicated					Section 10.3

Evaluation	Screening (up to 28 days before Day 1)		Treatment Phase ^a				End of Treatment (EOT)	Post Treatment Follow-up Phase			Notes
			Cycle 1			Cycle 2 and Beyond			Safety follow-up Period		Survival follow-up
	D- 28 to D- 15	D-14 to D-1	D1 (±1)	D8 (± 1)	D15 (±1)	D1 (± 2)	30 (±7) days after last IMPs admin	At 60 (±7) days after last IMPs admin	At 90 (±7) days after last IMPs admin	Every 90 days (±7) after last safety follow-up	
Blood Typing Interference Test		X				Cycle 2 Day 1 only					Section 10.3 Before each transfusion.
Serology HBV and HCV (for HCC only)	X										Section 10.3
Urinalysis (at baseline and if required) /urine dipstick		X	X	As clinically indicated			X	X	X		Section 10.3
Disease Assessment											
CT/MRI (for HCC, SCCHN, and EOC)	X					X (Weeks 9, 18, 27, and then every 12 weeks)	X (if necessary)		X (until PD is confirmed if no PD documented & confirmed)		Section 8.1
Brain MRI (for GBM only)		X (within 14 days prior to first dose)				X (Weeks 6, 12, 18, 24, and then every 9 weeks)	X (if necessary)		X (until PD is confirmed if no PD documented & confirmed)		Section 8.1
AFP (for HCC) / CA125 (for EOC)			X			X (Weeks 9, 18, 27, and then every 12 weeks)	X (if necessary)		X (until PD is confirmed if no PD documented & confirmed)		Section 8.1
Isatuximab Administration			X	X ^c	X ^c	X					^c evaluation not applicable for Cohort E
Atezolizumab Administration			X			X					

Evaluation	Screening (up to 28 days before Day 1)		Treatment Phase ^a				End of Treatment (EOT)	Post Treatment Follow-up Phase			Notes ^a A cycle is 21 days
			Cycle 1		Cycle 2 and Beyond				Safety follow-up Period		
	D- 28 to D- 15	D-14 to D-1	D1 (±1)	D8 (± 1)	D15 (±1)	D1 (± 2)	30 (±7) days after last IMPs admin	At 60 (±7) days after last IMPs admin	At 90 (±7) days after last IMPs admin	Every 90 days (±7) after last safety follow-up	
AE/SAE Assessment	X		Continuously throughout period					X (ongoing related AEs, ongoing SAEs at EOT and new related AE/SAEs)			
Prior/Concomitant Medication	X (within 14 days prior to first dose)		Continuously throughout period					X (related to AE/SAEs listed above)			
PK	See Pharmacokinetics and immunogenicity Flow Chart										
ADA	See Pharmacokinetics and immunogenicity Flow Chart										
Tumor Biopsy, Archival Tumor Tissue Collection, Biomarker Blood Draw	See Biomarker Flow Chart										
Subsequent Anticancer Therapy Status							X	X	X	X	
Survival Status										X	

PHARMACOKINETICS AND IMMUNOGENICITY FLOW CHART

Study Phase	Treatment Phase																			Post-treatment				
Cycle	Cycle 1									Cycle 2			Cycle 3		Cycle 4			Subsequent Cycles		EOT	FUP (90±7 days)			
Day within the Cycle	D1						D15			D1			D1		D1			D1						
Sample RNT (h) Ref: SOI Atezolizumab	SOI	EOI	EOI +30 min	-						SOI	-		SOI	-	SOI	-			SOI	-	-	-		
Sample time window Atezolizumab	[-24h, SOI]	±5 min	±10 min	-						[-24h, SOI]	-		[-24h, SOI]	-	[-24h, SOI]	-			[-24h, SOI]	-	-	-		
Sample RNT (h) Ref: SOI Isatuximab	SOI	-		EOI	EOI +4h	72h	168h (SOI of Day 8)		SOI	EOI	SOI	-	EOI	SOI	-	SOI	-	EOI	EOI +1h	SOI	-	-	-	
Sample time window Isatuximab	-	-		±10 min	±30 min	±5 h	[-24h, SOI]		[-24h, SOI]	±10 min	-	-	±10 min	-	-	-	-	±10 min	±10 min	-	-	-	-	
IMP administration (IV infusion)																								
Atezolizumab	X-----X									X			X			X				X				
Isatuximab			X-----X				X	X-----X				X-----X			X	X-----X				X				
Pharmacokinetics ^{a,i}																								
Atezolizumab ^j Sample ID	S00	-	S01	-						S00 ^c	-		S00 ^c	-	S00 ^c	-			S00 ^{c,j}	-	SF00	-		
Isatuximab ⁱ Sample ID	P00 ^b	-	-	P01 ^f	P02	P03	P04 ^{d,g}		P05 ^g	P06 ^f	P00 ^{b, g}	-	P01 ^f	P00 ^{b, g}	-	P00 ^{b, g}	-	P01 ^f	P02	P00 ^{b, g}	-	-	-	
Immunogenicity (ADA) ^{a, e, i}																								
Atezolizumab ^{c, j}	AS00	-								AS00	-		AS00	-	AS00	-			AS00 ^j	-	ASF00	-		
Isatuximab ^{g, h i}	AP00 ^b	-								AP00 ^b	-		AP00 ^b	-	AP00 ^b	-			AP00 ^b	-		APF00 ^h		

^a Refer to the laboratory manual for sample collection, processing and shipping.

^b Isatuximab PK or ADA predose samples should be collected before isatuximab administration and at the same time as atezolizumab predose samples when applicable.

^c Atezolizumab sample collected strictly before the start of atezolizumab infusion (during treatment phase).

^d P04 sample must be collected even if the participant withdraws the study without receiving the second administration of isatuximab.

^e After start of study treatment, if 1 drug is prematurely discontinued, ADA samples for this drug will be collected and tested on Day 1 of the 2 next cycles. The ADA sample collection for other drug should be done as planned.

- f* Sample collected just before actual end of infusion of isatuximab.
- g* Isatuximab sample collected strictly before the start of isatuximab infusion (during treatment phase).
- h* After FUP 90($\pm$ 7) days after last isatuximab administration no further ADA will be sampled even if the status is positive or inconclusive.
- i* Isatuximab PK and ADA samples to be collected each cycle until the last study treatment for PK and until FUP 90($\pm$ 7) days after last IMP administration for ADA or at the cut-off date, whichever comes first.
However, collection can be reduced or stopped earlier upon notification from the Sponsor based on the updated knowledge of isatuximab on PK/ADA.
- j* For subsequent cycles, atezolizumab PK and ADA samples (pre-dose) are to be collected at Cycle 8 Day 1 and Cycle 16 Day 1 only.

Abbreviations: SOL: Start of infusion, EOI: End of infusion, P= Plasma, S= Serum, AP= Antibody-Plasma, AS= Antibody-Serum, IMP= Investigational medicine product, D= Day, EOT= End of treatment, FUP= follow-up period, RNT= relative nominal time, IV= Intravenous, ID= Identification, ADA= Anti-drug antibody

EXPLORATORY BIOMARKER FLOW CHART

Sample Collection	Biomarker Analysis	Screening	Treatment Phase				Post-Treatment Phase		Notes
			Cycle 1	Cycle 2	Cycle 3	Cycle 5 and Beyond	EOT	Follow-up	
		D-28 to D-1	D1 (-1 day) a	D1	D1	D1	30 (±7) days after last IMP admin	60 (±7) days after last IMP admin	a. Biomarker blood draw may be performed within 1 working day prior to IMP administration.
Peripheral Blood	Immune genetic markers		X						
	Tumor mutational profile (plasma cell free DNA)		X	X	X		X		
	Immunophenotyping		X	X	X	X (odd cycles only)	X	X	Only for Phase 2 Stage 2 participants
	Plasma or serum cytokines	X	X	X	X	X (odd cycles only)	X	X	Only for Phase 2 Stage 2 participants
	Adaptive immunity		X	X (only 1 on-treatment sample, see Study Reference Manual for details)					Only Phase 2 Stage 2 participants, if sufficient clinical response is observable in Phase 1 and Phase 2 Stage 1 participants
	Plasma or serum Metabolites		X	X	X	X (odd cycles only)	X		Only Phase 2 Stage 2 participants, if sufficient clinical response is observable in Phase 1 and Phase 2 Stage 1 participants
Archival Pre-Treatment Tumor Tissue Collection (for HCC, SCCHN, EOC and GBM)		X							Mandatory for all GBM participants. For HCC, SCCHN, and EOC participants, provision of adequate archival tumor tissue is allowed to replace mandatory baseline biopsy
Baseline or on treatment tumor biopsy core in order of priority (Biopsy to be performed for HCC, SCCHN, and EOC if adequate archival tissue is not available)	Core 1	CD38 and PD-L1 expression, immune contexture (IHC analyses)	X		X (if clinically feasible)				Refer to 102 for details. At least 1 tumor core is required, but 3 cores are optimal. Refer to Laboratory Manual for details.
	Core 2	CD38 and PD-L1 expression, immune contexture (IHC analyses)	X		X (if clinically feasible)				
	Core 3	Transcriptomic analysis	X		X (if clinically feasible)				