

1.3. Schedule of Activities (SoA)

Note: For applicable participants who have their combination therapy held while SC weekly ramucirumab is continued, procedures noted in column DX are to be performed at the time of the additional ramucirumab doses.

Screening, On-Study, and Post-Treatment Schedule of Activities																	
	Screening		On-Treatment												Post-Treatment		
			Cycle = 21 days			Cycle = 21 days (or 28 days for Cohorts B & C per combination regimen)											
	(Day Relative to C1D1)	Cycle 1			Cycle 2			Cycle 2-n (If combo therapy is held and ramucirumab is given)		Cycle 3-n				Short-term follow-up ^a	Instructions		
		(±3 days)			(±3 days)			(±3 days)		(±3 days)						(±3 days)	
	≤28	≤7	D1	D8	D15	D1	D8	D15	D22	DX	D1	D8	D15	D22	V801	D22 for 28-day cycles only.	
Procedure																	
Informed consent	X															ICF must be signed before any protocol-specific procedures are performed	
Inclusion/Exclusion criteria	X																
Medical history	X															Including assessment of preexisting conditions, historical illnesses, and habits (such as tobacco and alcohol use).	
Cancer treatment history	X															Record prior anti-cancer therapy	
Concomitant medication	X		X													- At baseline, record prior and concurrent medications. - Record all premedication, supportive care, and concomitant medication continuously at every visit and throughout the study.	
Physical examination	X		X	X	X	X					X				X		

Screening, On-Study, and Post-Treatment Schedule of Activities																
	Screening		On-Treatment												Post-Treatment	
			Cycle = 21 days			Cycle = 21 days (or 28 days for Cohorts B & C per combination regimen)										
	(Day Relative to C1D1)		Cycle 1			Cycle 2			Cycle 2-n (If combo therapy is held and ramucirumab is given)	Cycle 3-n				Short-term follow-up ^a	Instructions	
			(±3 days)			(±3 days)			(±3 days)	(±3 days)				(±3 days)		
	≤28	≤7	D1	D8	D15	D1	D8	D15	D22	DX	D1	D8	D15	D22	V801	D22 for 28-day cycles only.
Procedure																
Vital signs	X		See instructions			X	X	X	X	X	X	X	X	X	X	Measure vital signs (height [at baseline], weight, temperature, blood pressure, pulse rate, and respiration rate). On days with concurrent ECGs, vital signs, and/or PK sampling, these measurements should occur at approximately the same time. Vital sign measurements will be obtained: • Day 1 at predose (prior to SC ramucirumab dose, but within 90 minutes prior to predose blood draw) • C1D2 (24 hours postdose) • C1D8 • C1D15 • Any time for other specified visits, but prior to SC ramucirumab dose at same visit.
AE collection	X		X												X	• Collect continuously at every visit and throughout the study. • CTCAE Version 5.0 • See Section 10.3 for AE grading for lab investigations.

Screening, On-Study, and Post-Treatment Schedule of Activities																
	Screening		On-Treatment												Post-Treatment	
			Cycle = 21 days			Cycle = 21 days (or 28 days for Cohorts B & C per combination regimen)										
	(Day Relative to C1D1)	Cycle 1			Cycle 2			Cycle 2-n (If combo therapy is held and ramucirumab is given)		Cycle 3-n				Short-term follow-up ^a	Instructions	
		(±3 days)			(±3 days)			(±3 days)		(±3 days)						
	≤28	≤7	D1	D8	D15	D1	D8	D15	D22	DX	D1	D8	D15	D22	V801	D22 for 28-day cycles only.
Procedure																
ECOG PS	X		X			X					X				X	During study treatment, perform ≤3 days prior to treatment.
ECG	X		See instructions												X	Single, local. On days with concurrent measurements, see vital sign row for more details. Times are referenced to the end of dosing. ECGs will be obtained: • C1D1 at predose. • Every 6 weeks thereafter, predose. • Any other time as clinically indicated.
Hematology		X	X	X	X	X	X	X		X	X				X	See Appendix 2 .
Coagulation		X				X										See Appendix 2 . Perform at baseline, C2D1, D1 of every other cycle afterwards (C4D1, C6D1, etc.), and as clinically indicated.
Clinical chemistry		X	X	X	X	X	X	X		X	X				X	See Appendix 2 .
Urinalysis		X	X	X	X	X	X	X	X	X	X	X	X	X	X	See Appendix 2 . In addition, perform as clinically indicated.
Pregnancy test		X	See instructions												X	• Applies only to women of childbearing potential.

Screening, On-Study, and Post-Treatment Schedule of Activities																
	Screening		On-Treatment												Post-Treatment	Instructions
			Cycle = 21 days			Cycle = 21 days (or 28 days for Cohorts B & C per combination regimen)										
	(Day Relative to C1D1)	Cycle 1			Cycle 2			Cycle 2-n (If combo therapy is held and ramucirumab is given)		Cycle 3-n				Short-term follow-up ^a		
		(±3 days)			(±3 days)			(±3 days)		(±3 days)					(±3 days)	
	≤28	≤7	D1	D8	D15	D1	D8	D15	D22	DX	D1	D8	D15	D22	V801	D22 for 28-day cycles only.
Procedure																
																Note: During study treatment, perform monthly or as required per local regulations and/or institutional guidelines. See Appendix 2.
Thyroid panel		X	See instructions												X	See Appendix 2 . Starting C1D1, then every 3 months thereafter.
Radiologic imaging and measurement of palpable or visible lesions	X		See instructions												X	Perform according to RECIST 1.1, by the same method used at baseline, every 8 weeks (±7 days) for the first 6 months after C1D1 and every 12 weeks (±7 days) thereafter until radiographic disease progression, death, or study completion, whichever occurs first. Perform as scheduled, even if study treatment is delayed or omitted.
Injection site assessments (solicited)			See instructions			X	X	X	X	X	X	X	X	X		Prior to each injection and at the designated timepoints, the investigator/investigator designee will ask the participant if they have any injection site concerns and record the participant's response. 1.Cycle 1 collection:

Screening, On-Study, and Post-Treatment Schedule of Activities																
	Screening		On-Treatment												Post-Treatment	
			Cycle = 21 days			Cycle = 21 days (or 28 days for Cohorts B & C per combination regimen)										
	(Day Relative to C1D1)	Cycle 1			Cycle 2			Cycle 2-n (If combo therapy is held and ramucirumab is given)	Cycle 3-n				Short-term follow-up ^a	Instructions		
		(±3 days)			(±3 days)			(±3 days)	(±3 days)							
	≤28	≤7	D1	D8	D15	D1	D8	D15	D22	DX	D1	D8	D15	D22	V801	D22 for 28-day cycles only.
Procedure																
																a.C1D1: 5-15 min after injection is complete and 60 min after injection is complete b.C1D2 (approximately 24 hr after injection) c.C1D4 (± 1 day) d.C1D8: 5-15 min after injection and 60 min after injection e.C1D15: 5-15 min after injection and 60 min after injection 2.Cycle 2-n collection: a.5-15 min after injection and 60 min after injection The findings from the assessment are recorded on the ISR questionnaire. All positive responses of pain require an additional assessment using the participant-completed Pain VAS. If there are ≥2 injections, the ISR questionnaire will be filled out separately for each site, after all injections have been administered. See Section 8.2.5 for details.

Screening, On-Study, and Post-Treatment Schedule of Activities																	
	Screening		On-Treatment												Post-Treatment		
			Cycle = 21 days			Cycle = 21 days (or 28 days for Cohorts B & C per combination regimen)											
	(Day Relative to C1D1)	Cycle 1			Cycle 2			Cycle 2-n (If combo therapy is held and ramucirumab is given)		Cycle 3-n				Short-term follow-up ^a	Instructions		
		(±3 days)			(±3 days)			(±3 days)		(±3 days)						(±3 days)	
	≤28	≤7	D1	D8	D15	D1	D8	D15	D22	DX	D1	D8	D15	D22	V801	D22 for 28-day cycles only.	
Procedure																	
Injection site assessments (spontaneous)			See instructions													1. ISR questionnaire completed outside of solicited assessment schedule. 2. May be triggered by participant anytime during the cycle based on diary. 3. If the participant indicates presence of pain during spontaneous ISR reporting, no VAS assessment is required.	
Participant diary			See instructions													• Participant will assess injection site(s) daily. If any ISR symptoms present, the participant should contact the investigator. • Diary should be returned to the investigator at each weekly visit.	
Sample collection																See Section 1.3.1 for PK and IG.	
PK				See Section 1.3.1													
IG				See Section 1.3.1													

Screening, On-Study, and Post-Treatment Schedule of Activities																
	Screening		On-Treatment												Post-Treatment	
			Cycle = 21 days			Cycle = 21 days (or 28 days for Cohorts B & C per combination regimen)										
	(Day Relative to C1D1)	Cycle 1			Cycle 2			Cycle 2-n (If combo therapy is held and ramucirumab is given)		Cycle 3-n				Short-term follow-up ^a	Instructions	
		(±3 days)			(±3 days)			(±3 days)		(±3 days)						(±3 days)
	≤28	≤7	D1	D8	D15	D1	D8	D15	D22	DX	D1	D8	D15	D22	V801	D22 for 28-day cycles only.
Procedure																
Administer ramucirumab			See instructions													See Section 6.1. Participants must be closely monitored for a 1-hour observation period following the administration of SC ramucirumab (see Section 6.1).
Administer combination medications, if applicable (Cohorts B and C only)			See instructions							See instructions					Begin Combination anti-cancer therapies along with weekly SC ramucirumab at Cycle-2. See details in Section 6.1. Erlotinib (1L EGFR+ NSCLC), docetaxel (2L NSCLC), paclitaxel (2L gastric), FOLFIRI (2L CRC).	

Abbreviations: 1L = first line; 2L = second line; AE = adverse event; C = cycle; CRC = colorectal cancer; CTCAE = common terminology criteria for adverse events; D = day; ECG = electrocardiogram; ECOG PS = Eastern Cooperative Oncology Group Performance Score; EGFR = epidermal growth factor receptor; FOLFIRI = fluorouracil and irinotecan; hr = hour; ICF = informed consent form; IG = immunogenicity; ISR = injection site reaction; min = minutes; NSCLC = non-small cell lung cancer; PK = pharmacokinetic; RECIST 1.1 = Response Evaluation Criteria in Solid Tumors version 1.1; SC = subcutaneous; V = visit; VAS = visual analogue scale.

- a Short-term follow-up begins when the participant and investigator agree that the participant will no longer continue study treatment and lasts a minimum of 30 days following the last dose of study drug.

Continued Access Schedule of Activities			
Visit	Study Treatment	Follow-Up ^a	Instructions
Procedure	501-5XX	901	
AE Collection	X	X	Grading via CTCAE, Version 5.0. As part of AE collection, monitor vital signs and perform standard laboratory tests (hematology, chemistry, urinalysis, and pregnancy testing) at the same frequency as during the study treatment period. See Section 10.3 for AE grading for lab investigations. All laboratory tests during the continued access period will be performed in the local laboratories only.
PK, IG, and exploratory hypersensitivity			In the event of hypersensitivity, blood samples will be collected for PK, IG, and exploratory hypersensitivity analyses at the following time points, as close as possible to: (i) the onset of the hypersensitivity, (ii) the resolution of the hypersensitivity, and (iii) 30 (±3) days following the hypersensitivity.
Administer ramucirumab	X		See Section 6.1. Participants must be closely monitored for a 1-hour observation period following the ramucirumab injection (see Section 6.1).

Abbreviations: AE = adverse event; CTCAE = common terminology criteria for adverse events (NCI 2018); IG = immunogenicity; NCI = National Cancer Institute; PK = pharmacokinetic.

- ^a Continued-access follow-up begins when the participant and the investigator agree that the participant will no longer continue treatment in the continued-access period and lasts a minimum of 30 days following the last dose of study drug. In all cases, no follow-up procedures will be performed for a participant who withdraws informed consent unless he or she has explicitly provided permission and consent.

1.3.1. Pharmacokinetic Sampling Schedule

General Instructions:

The pharmacokinetic sampling table applies to all participants irrespective of 21-day or 28-day cycle length.

Adequately record the actual collection date and time (24-hour clock; for example, 16:00, not 4 pm) on the requisition form of the sample.

In case of cycle delay and/or ramucirumab dosing interruption, if possible, collect an additional unscheduled PK samples using the appropriate central lab collection kit:

- at time as close as possible to the onset of the adverse event (AE) (if the cycle delay or ramucirumab dosing interruption is due to an AE).
- at the time when ramucirumab dose should have been administered (if there had not been a delay or dose interruption).
- prior to resuming the ramucirumab dosing after dosing interruption/dosing delay.

Note: If the above time points for unscheduled PK sampling happen to occur at the same time as a scheduled PK time point (see table below), then only the scheduled sample will be taken.

Sample #	Study Cycle	Day within Cycle	Collection Time Point Relative to Ramucirumab Weekly Dose	Ramucirumab PK Collection	Immunogenicity Collection
1	1	1	-1 hr (predose)	X	X
2	1	2	24 hrs $\pm$ 2 postdose	X	
3	1	4 $\pm$ 1	48-96 hrs postdose window	X	
4	1	8	-1 hr (predose)	X	
5	1	15	-1 hr (predose)	X	X
6	1	18 $\pm$ 1	48-96 hrs postdose window	X	
7	2	1	-1 hr (predose)	X	
8	2	8	-1 hr (predose)	X	X
9	2	11 $\pm$ 1	48-96 hrs postdose window	X	
10	3	1	-1 hr (predose)	X	
11	3	15	-1 hr (predose)	X	
12	3	18 $\pm$ 1	48-96 hrs postdose window	X	
13	4	1	-1 hr (predose)	X	X
14	5	1	-1 hr (predose)	X	
End of treatment			30 days post last dose	X	X

Abbreviations: hr = hour; PK = pharmacokinetic.