

1.3. Schedule of Activities (SoA)

1.3.1. Schedule of Activities for the Screening and Induction Treatment Periods

Table 1. Schedule of Activities for the Screening and Induction Treatment Periods of Study J1P-MC-KFAH										
V1 and V9 procedures may be conducted over more than 1 day as long as all activities are completed within the allowable visit tolerance and all scheduled activities are completed prior to dosing.										
For procedures at an ETV, see ETV in Table 4.										
For procedures at an unscheduled visit, see V997 in Table 4.										
Visit number	V1	V2	V3	V4	V5	V6	V7	V8	V9	Comment
Weeks from randomization	-5	—	1	2	4	6	8	10	12	
Study day	—	1	8	15	29	43	57	71	85	
Visit interval tolerance (days)	≤35 before V2	—	±3	±3	±3	±3	±3	±3	±3	
Fasting visit		X							X	
Informed consent	X									
Site staff explains, participant reports remission/normal stool frequency	X									The purpose of UC remission question should be explained before the participant answers the question. Errors in the response cannot be corrected after the participant saves and confirms the response.
Inclusion and exclusion criteria, review and confirm	X	X								
Demographics	X									
Preexisting conditions and medical history	X									Includes relevant surgical history.
Prespecified medical history	X									Indication and history of interest.
Prior treatments for indication	X									Includes all UC-specific therapies since date of diagnosis.
Concomitant medications	X	X	X	X	X	X	X	X	X	
Adverse events	X	X	X	X	X	X	X	X	X	For AESIs, additional data are collected (Section 8.3.6).
Review modified Mayo score (MMS)		X							X	
Tobacco use	X								X	
Alcohol and caffeine use	X									

Table 1. Schedule of Activities for the Screening and Induction Treatment Periods of Study J1P-MC-KFAH

V1 and V9 procedures may be conducted over more than 1 day as long as all activities are completed within the allowable visit tolerance and all scheduled activities are completed prior to dosing.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visit number	V1	V2	V3	V4	V5	V6	V7	V8	V9	Comment
Weeks from randomization	-5	—	1	2	4	6	8	10	12	
Study day	—	1	8	15	29	43	57	71	85	
Visit interval tolerance (days)	≤35 before V2	—	±3	±3	±3	±3	±3	±3	±3	
Fasting visit		X							X	
Physical Evaluation										
Height	X									Participant should remove shoes.
Weight	X								X	
Vital signs	X	X	X	X	X	X	X	X	X	Blood pressure, body temperature, and pulse rate. Vital signs should be measured after participant has been sitting at least 5 minutes.
Physical examination	X									The complete physical exam excludes pelvic, rectal, and breast exams, unless clinically indicated, and includes assessment for signs and symptoms of tuberculosis (TB), including peripheral lymph nodes. See Section 8.2.2.
Symptom-directed physical examination, including assessment for signs and symptoms of tuberculosis (TB), including peripheral lymph nodes		X							X	See Section 8.2.2.
CCI	X	X			X		X		X	
12-lead electrocardiogram (ECG)	X								X	Locally performed.

Table 1. Schedule of Activities for the Screening and Induction Treatment Periods of Study J1P-MC-KFAH

V1 and V9 procedures may be conducted over more than 1 day as long as all activities are completed within the allowable visit tolerance and all scheduled activities are completed prior to dosing.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visit number	V1	V2	V3	V4	V5	V6	V7	V8	V9	Comment
Weeks from randomization	-5	—	1	2	4	6	8	10	12	
Study day	—	1	8	15	29	43	57	71	85	
Visit interval tolerance (days)	≤35 before V2	—	±3	±3	±3	±3	±3	±3	±3	
Fasting visit		X							X	
Chest x-ray (posterior-anterior view)	X									Interpreted and reported by radiologist or pulmonologist. Not done at screening if done within 3 months before screening <u>and</u> if qualifying radiographs or equivalent imaging study and/or formal report are available for investigator's review. Lateral view may also be taken. Locally performed.
Patient Diary (Electronic)										
Patient diary dispensed	X									
Diary compliance check		X	X	X	X	X	X	X	X	Check compliance for Stool Frequency Rectal Bleeding Nocturnal Stools Bristol Stool Scale Urgency NRS Abdominal Pain NRS Fatigue NRS, and Patient Global Rating of Severity (PGR-S).
Patient-Reported Outcomes (Electronic)										
Inflammatory Bowel Disease Questionnaire (IBDQ)		X							X	

Table 1. Schedule of Activities for the Screening and Induction Treatment Periods of Study J1P-MC-KFAH

V1 and V9 procedures may be conducted over more than 1 day as long as all activities are completed within the allowable visit tolerance and all scheduled activities are completed prior to dosing.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visit number	V1	V2	V3	V4	V5	V6	V7	V8	V9	Comment
Weeks from randomization	-5	—	1	2	4	6	8	10	12	
Study day	—	1	8	15	29	43	57	71	85	
Visit interval tolerance (days)	≤35 before V2	—	±3	±3	±3	±3	±3	±3	±3	
Fasting visit		X							X	
Patient Global Impression of Change (PGI-C)									X	
QIDS-SR16		X							X	
Clinician-Administered Questionnaires (Paper)										
Physician's Global Assessment (PGA)	X								X	
C-SSRS Screening/Baseline	X									Adapted for the assessment of ideation and behavior categories only.
Self-Harm Supplement Form	X									
Self-Harm Follow-Up Form	X									Required if triggered by the Self-Harm Supplement Form per instructions.
Laboratory Tests and Sample Collections										
Hematology	X	X	X	X	X	X	X	X	X	
Clinical chemistry	X	X	X	X	X	X	X	X	X	
Lipid panel		X							X	Fasting samples are preferred, but if participant is non fasting at time of collection, this is not a protocol deviation.
Urinalysis		X			X		X		X	
Serum pregnancy	X									Perform only for women of childbearing potential and women with a history of tubal ligation.

Table 1. Schedule of Activities for the Screening and Induction Treatment Periods of Study J1P-MC-KFAH

V1 and V9 procedures may be conducted over more than 1 day as long as all activities are completed within the allowable visit tolerance and all scheduled activities are completed prior to dosing.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visit number	V1	V2	V3	V4	V5	V6	V7	V8	V9	Comment
Weeks from randomization	-5	—	1	2	4	6	8	10	12	
Study day	—	1	8	15	29	43	57	71	85	
Visit interval tolerance (days)	≤35 before V2	—	±3	±3	±3	±3	±3	±3	±3	
Fasting visit		X							X	
Urine pregnancy (local)		X			X		X		X	Perform only for women of childbearing potential and women with a history of tubal ligation. At dosing visits which have pregnancy testing, the pregnancy test result must be confirmed to be negative prior to dosing. Pregnancy testing may be performed more frequently than designated on the SoA, if local laws or regulations require more frequent testing.
Follicle-stimulating hormone (FSH)	X									Optional, perform only for women <u>not</u> of childbearing potential to confirm postmenopausal status.
CCI		X	X	X	X	X	X	X	X	
Tuberculosis (TB) test	X									Patients who had a tuberculin skin test (TST) will return from 48 to 72 hours after placement to have the test result read. Samples may be sent to central or local laboratory based on the type of test. A local laboratory must be qualified by local regulations.
Human immunodeficiency virus (HIV) screening tests	X									
Hepatitis C virus (HCV) screening tests	X									HCV RNA will be measured to confirm positive hepatitis C virus antibody.
Hepatitis B virus (HBV) screening tests	X									Includes testing for HBsAg and anti-HBc.

Table 1. Schedule of Activities for the Screening and Induction Treatment Periods of Study J1P-MC-KFAH

V1 and V9 procedures may be conducted over more than 1 day as long as all activities are completed within the allowable visit tolerance and all scheduled activities are completed prior to dosing.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visit number	V1	V2	V3	V4	V5	V6	V7	V8	V9	Comment
Weeks from randomization	-5	—	1	2	4	6	8	10	12	
Study day	—	1	8	15	29	43	57	71	85	
Visit interval tolerance (days)	≤35 before V2	—	±3	±3	±3	±3	±3	±3	±3	
Fasting visit		X							X	
HBV DNA	X								X	Only for participants who are positive for anti-HBc at screening.
Pharmacokinetic (PK) samples		X	X	X	X		X		X	Collect samples before dosing, if dosing is scheduled. Collect additional samples at specified times relative to onset of hypersensitivity events.
CCI		X		X	X		X		X	Collect samples before dosing, if dosing is scheduled. Collect additional samples at specified times relative to onset of hypersensitivity events.
Flow cytometry panel		X		X					X	
Genetics sample		X								Blood sample for DNA pharmacogenetics can be obtained any time at or after V2.
CCI		X			X		X		X	
Endoscopic Procedure										
Endoscopy	X								X	Screening endoscopy must occur ≤14 days prior to randomization. At screening, perform only for participants who are eligible for the study by all other study entry criteria.
Colon biopsy sample collection	X								X	Performed at time of endoscopy. Instructions for collection are provided in the lab manual. Flexible sigmoidoscopy is recommended, except for protocol-specified reasons when a colonoscopy should be performed.

Table 1. Schedule of Activities for the Screening and Induction Treatment Periods of Study J1P-MC-KFAH

V1 and V9 procedures may be conducted over more than 1 day as long as all activities are completed within the allowable visit tolerance and all scheduled activities are completed prior to dosing.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visit number	V1	V2	V3	V4	V5	V6	V7	V8	V9	Comment
Weeks from randomization	-5	—	1	2	4	6	8	10	12	
Study day	—	1	8	15	29	43	57	71	85	
Visit interval tolerance (days)	≤35 before V2	—	±3	±3	±3	±3	±3	±3	±3	
Fasting visit		X							X	
Stool Samples										
Stool culture	X									Additional local stool testing (for example, for ova and parasites) is allowed at the investigator's discretion.
<i>C. difficile</i> testing	X									Additional <i>C. difficile</i> testing is allowed throughout the study at the investigator's discretion.
CCI		X			X		X		X	
		X			X		X		X	
Randomization and Dosing										
Randomization		X								
Dosing		X		X	X	X	X	X	X	No dosing at V3.

Abbreviations: AESIs = adverse event of special interest; anti-HBc = antibody to hepatitis B core antigen; C-SSRS = Columbia–Suicide Severity Rating Scale;

DNA = deoxyribonucleic acid; ETV = early termination visit; NRS = numeric rating scale; HBsAg = hepatitis B surface antigen; QIDS-SR16 = 16-Item

Quick Inventory of Depressive Symptomatology–Self Report; RNA = ribonucleic acid; SoA = schedule of activities; UC= ulcerative colitis; V = visit.

1.3.2. Schedule of Activities for the Maintenance Treatment Period

Table 2. Schedule of Activities for the Maintenance Treatment Period of Study J1P-MC-KFAH

This schedule applies to study participants who are **responders** at Week 12.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 10 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof), if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring.

Visit number	V10	V11	V12	V13	V14	V15	V16	V17	V18	V19	Comment
Weeks from randomization	14	16	18	20	22	24	26	28	30	32	
Study day	99	113	127	141	155	169	183	197	211	225	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit						X					
Concomitant medications	X	X	X	X	X	X	X	X	X	X	
Adverse events	X	X	X	X	X	X	X	X	X	X	For AESIs, additional data are collected (Section 8.3.6).
Tobacco use						X					
Physical Evaluation											
Weight						X					
Vital signs	X	X	X	X	X	X	X	X	X	X	Blood pressure, body temperature, and pulse rate. Vital signs should be measured after participant has been sitting for at least 5 minutes.
Symptom-directed physical examination, including assessment for signs and symptoms of tuberculosis (TB), including peripheral lymph nodes						X					See Section 8.2.2.
CCI						X					
12-lead electrocardiogram (ECG)						X					Locally performed.
Patient Diary (Electronic)											

Table 2. Schedule of Activities for the Maintenance Treatment Period of Study J1P-MC-KFAH

This schedule applies to study participants who are **responders** at Week 12.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 10 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof), if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring.

Visit number	V10	V11	V12	V13	V14	V15	V16	V17	V18	V19	Comment
Weeks from randomization	14	16	18	20	22	24	26	28	30	32	
Study day	99	113	127	141	155	169	183	197	211	225	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit						X					
Diary compliance check	X	X	X	X	X	X	X	X	X	X	Check compliance for Stool Frequency Rectal Bleeding Nocturnal Stools Bristol Stool Scale Urgency NRS Abdominal Pain NRS Fatigue NRS, and Patient Global Rating of Severity (PGR-S).
Patient-Reported Outcomes (Electronic)											
Inflammatory Bowel Disease Questionnaire (IBDQ)						X					
Patient Global Impression of Change (PGI-C)						X					
QIDS-SR16						X					
Laboratory Tests and Sample Collections											
Hematology		X		X		X		X		X	
Clinical chemistry		X		X		X		X		X	

Table 2. Schedule of Activities for the Maintenance Treatment Period of Study J1P-MC-KFAH

This schedule applies to study participants who are **responders** at Week 12.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 10 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof), if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring.

Visit number	V10	V11	V12	V13	V14	V15	V16	V17	V18	V19	Comment
Weeks from randomization	14	16	18	20	22	24	26	28	30	32	
Study day	99	113	127	141	155	169	183	197	211	225	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit						X					
Lipid panel						X					Fasting samples are preferred, but if participant is nonfasting at time of collection, this is not a protocol deviation.
Urinalysis		X		X		X					
Urine pregnancy (local)		X		X		X		X		X	Perform only for women of childbearing potential and women with a history of tubal ligation. At dosing visits which have pregnancy testing, the pregnancy test result must be confirmed to be negative prior to dosing. Pregnancy testing may be performed more frequently than designated on the SoA, if local laws or regulations require more frequent testing.
CCI						X					
HBV DNA						X					Only for participants who are positive for anti-HBc at screening.
Pharmacokinetic (PK) samples						X					Collect samples before dosing, if dosing is scheduled. Collect additional samples at specified times relative to onset of hypersensitivity events.

Table 2. Schedule of Activities for the Maintenance Treatment Period of Study J1P-MC-KFAH

This schedule applies to study participants who are **responders** at Week 12.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 10 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof), if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring.

Visit number	V10	V11	V12	V13	V14	V15	V16	V17	V18	V19	Comment
Weeks from randomization	14	16	18	20	22	24	26	28	30	32	
Study day	99	113	127	141	155	169	183	197	211	225	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit						X					
CCI						X					Collect samples before dosing, if dosing is scheduled. Collect additional samples at specified times relative to onset of hypersensitivity events.
Flow cytometry panel						X					
Genetics sample											Blood sample for DNA pharmacogenetics can be obtained any time at or after V2.
CCI						X					
Endoscopic Procedure											Not applicable for responders during the time period of V10 through V19
Stool Samples											
CCI						X					
						X					
Randomization and Dosing											
Dosing	X	X	X	X	X	X	X	X	X	X	

Table 2 (continued). Schedule of activities for the Maintenance Treatment Period of Study J1P-MC-KFAH

This schedule applies to study participants who are **responders** at Week 12.

For procedures at an early termination visit, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 10 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring.

Visit number	V20	V21	V22	V23	V24	V25	V26	V27	V28	V29	Comment
Weeks from randomization	34	36	38	40	42	44	46	48	50	52	
Study day	239	253	267	281	295	309	323	337	351	365	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit										X	
Concomitant medications	X	X	X	X	X	X	X	X	X	X	
Adverse events	X	X	X	X	X	X	X	X	X	X	For AESIs, additional data are collected (Section 8.3.6).
Review modified Mayo score (MMS)										X	
Tobacco use										X	
Physical Evaluation											
Weight										X	
Vital signs	X	X	X	X	X	X	X	X	X	X	Blood pressure, body temperature, and pulse rate. Vital signs should be measured after participant has been sitting for at least 5 minutes.
Symptom-directed physical examination, including assessment for signs and symptoms of tuberculosis (TB), including peripheral lymph nodes		X								X	See Section 8.2.2.
CCI		X								X	
12-lead electrocardiogram (ECG)										X	Locally performed.

Table 2 (continued). Schedule of activities for the Maintenance Treatment Period of Study J1P-MC-KFAH

This schedule applies to study participants who are **responders** at Week 12.

For procedures at an early termination visit, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 10 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring.

Visit number	V20	V21	V22	V23	V24	V25	V26	V27	V28	V29	Comment
Weeks from randomization	34	36	38	40	42	44	46	48	50	52	
Study day	239	253	267	281	295	309	323	337	351	365	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit										X	
Patient Diary (Electronic)											
Diary compliance check	X	X	X	X	X	X	X	X	X	X	Check compliance for Stool Frequency Rectal Bleeding Nocturnal Stools Bristol Stool Scale Urgency NRS Abdominal Pain NRS Fatigue NRS, and Patient Global Rating of Severity (PGR-S).
Diary return										X	
Patient-Reported Outcomes (Electronic)											
Inflammatory Bowel Disease Questionnaire (IBDQ)										X	
Patient Global Impression of Change (PGI-C)										X	
QIDS-SR16										X	
Clinician-Administered Questionnaires (Paper)											
Physician's Global Assessment (PGA)										X	

Table 2 (continued). Schedule of activities for the Maintenance Treatment Period of Study J1P-MC-KFAH

This schedule applies to study participants who are **responders** at Week 12.

For procedures at an early termination visit, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 10 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring.

Visit number	V20	V21	V22	V23	V24	V25	V26	V27	V28	V29	Comment
Weeks from randomization	34	36	38	40	42	44	46	48	50	52	
Study day	239	253	267	281	295	309	323	337	351	365	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit										X	
Laboratory Tests and Sample Collections											
Hematology		X		X		X		X		X	
Clinical chemistry		X		X		X		X		X	
Lipid panel										X	Fasting samples are preferred, but if participant is nonfasting at time of collection, this is not a protocol deviation.
Urinalysis										X	
Urine pregnancy (local)		X		X		X		X		X	Perform only for women of childbearing potential and women with a history of tubal ligation. At dosing visits which have pregnancy testing, the pregnancy test result must be confirmed to be negative prior to dosing. Pregnancy testing may be performed more frequently than designated on the SoA, if local laws or regulations require more frequent testing.
CCI										X	
HBV DNA		X						X			Only for participants who are positive for anti-HBc at screening.

Table 2 (continued). Schedule of activities for the Maintenance Treatment Period of Study J1P-MC-KFAH

This schedule applies to study participants who are **responders** at Week 12.

For procedures at an early termination visit, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 10 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring.

Visit number	V20	V21	V22	V23	V24	V25	V26	V27	V28	V29	Comment
Weeks from randomization	34	36	38	40	42	44	46	48	50	52	
Study day	239	253	267	281	295	309	323	337	351	365	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit										X	
Pharmacokinetic (PK) samples		X								X	Collect samples before dosing, if dosing is scheduled. Collect additional samples at specified times relative to onset of hypersensitivity events.
CCI		X								X	Collect samples before dosing, if dosing is scheduled. Collect additional samples at specified times relative to onset of hypersensitivity events.
Flow cytometry panel										X	
Genetics sample											Blood sample for DNA pharmacogenetics can be obtained any time at or after V2.
CCI		X								X	
Endoscopic Procedure											
Endoscopy										X	
Colon biopsy sample collection										X	Performed at time of endoscopy. Instructions for collection are provided in the lab manual. Flexible sigmoidoscopy is recommended, except for protocol-specified reasons when a colonoscopy should be performed.
Stool Samples											
CCI		X								X	
		X								X	

Table 2 (continued). Schedule of activities for the Maintenance Treatment Period of Study J1P-MC-KFAH

This schedule applies to study participants who are **responders** at Week 12.

For procedures at an early termination visit, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 10 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring.

Visit number	V20	V21	V22	V23	V24	V25	V26	V27	V28	V29	Comment
Weeks from randomization	34	36	38	40	42	44	46	48	50	52	
Study day	239	253	267	281	295	309	323	337	351	365	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit										X	
Randomization and Dosing											
Dosing	X	X	X	X	X	X	X	X	X		No dosing at V29.

Abbreviations: AESIs = adverse event of special interest; anti-HBc = antibody to hepatitis B core antigen; DNA = deoxyribonucleic acid; ETV = early termination visit; HBV = hepatitis B virus; IT = information technology; NRS = numeric rating scale; QIDS-SR16 = 16-Item Quick Inventory of Depressive Symptomatology–Self Report; SoA = schedule of activities; V = visit.

1.3.3. Schedule of Activities for the Extension Induction and Extension Maintenance Periods

Table 3. Schedule of Activities for the Extension Induction and Extension Maintenance Treatment Periods of Study J1P-MC-KFAH

This schedule applies to study participants who are **nonresponders** at Week 12.

V16 procedures may be conducted over more than 1 day as long as all activities are completed within the allowable visit tolerance and all scheduled activities are completed prior to dosing

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 17 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring.

Visit number	V10	V11	V12	V13	V14	V15	V16	V17	V18	V19	Comment
Weeks from randomization	14	16	18	20	22	24	26	28	30	32	
Study day	99	113	127	141	155	169	183	197	211	225	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit							X				
Concomitant medications	X	X	X	X	X	X	X	X	X	X	
Adverse events	X	X	X	X	X	X	X	X	X	X	For AESIs, additional data are collected (Section 8.3.6).
Review modified Mayo score (MMS)							X				
Tobacco use							X				
Physical Evaluation											
Weight	X						X				
Vital signs	X	X	X	X	X	X	X	X	X	X	Blood pressure, body temperature, and pulse rate. Vital signs should be measured after participant has been sitting for at least 5 minutes.
Symptom-directed physical examination, including assessment for signs and symptoms of tuberculosis (TB), including peripheral lymph nodes	X						X				See Section 8.2.2.
CCI	X		X		X		X				
12-lead electrocardiogram (ECG)	X						X				Locally performed.
Patient Diary (Electronic)											

Table 3. Schedule of Activities for the Extension Induction and Extension Maintenance Treatment Periods of Study J1P-MC-KFAH

This schedule applies to study participants who are **nonresponders** at Week 12.

V16 procedures may be conducted over more than 1 day as long as all activities are completed within the allowable visit tolerance and all scheduled activities are completed prior to dosing

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 17 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring.

Visit number	V10	V11	V12	V13	V14	V15	V16	V17	V18	V19	Comment
Weeks from randomization	14	16	18	20	22	24	26	28	30	32	
Study day	99	113	127	141	155	169	183	197	211	225	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit							X				
Diary compliance check	X	X	X	X	X	X	X	X	X	X	Check compliance for Stool Frequency Rectal Bleeding Nocturnal Stools Bristol Stool Scale Urgency NRS Abdominal Pain NRS Fatigue NRS, and Patient Global Rating of Severity (PGR-S).
Patient-Reported Outcomes (Electronic)											
Inflammatory Bowel Disease Questionnaire (IBDQ)							X				
Patient Global Impression of Change (PGI-C)							X				
QIDS-SR16							X				
Clinician-Administered Questionnaires (Paper)											
Physician's Global Assessment (PGA)							X				

Table 3. Schedule of Activities for the Extension Induction and Extension Maintenance Treatment Periods of Study J1P-MC-KFAH

This schedule applies to study participants who are **nonresponders** at Week 12.

V16 procedures may be conducted over more than 1 day as long as all activities are completed within the allowable visit tolerance and all scheduled activities are completed prior to dosing

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 17 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring.

Visit number	V10	V11	V12	V13	V14	V15	V16	V17	V18	V19	Comment
Weeks from randomization	14	16	18	20	22	24	26	28	30	32	
Study day	99	113	127	141	155	169	183	197	211	225	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit							X				
Laboratory Tests and Sample Collections											
Hematology	X	X	X	X	X	X	X	X		X	
Clinical chemistry	X	X	X	X	X	X	X	X		X	
Lipid panel							X				Fasting samples are preferred, but if participant is nonfasting at time of collection, this is not a protocol deviation.
Urinalysis	X	X	X	X	X		X				
Urine pregnancy (local)		X		X		X		X		X	Perform only for women of childbearing potential and women with a history of tubal ligation. At dosing visits which have pregnancy testing, the pregnancy test result must be confirmed to be negative prior to dosing. Pregnancy testing may be performed more frequently than designated on the SoA if local laws or regulations require more frequent testing.
CCI	X		X		X		X				

Table 3. Schedule of Activities for the Extension Induction and Extension Maintenance Treatment Periods of Study J1P-MC-KFAH

This schedule applies to study participants who are **nonresponders** at Week 12.

V16 procedures may be conducted over more than 1 day as long as all activities are completed within the allowable visit tolerance and all scheduled activities are completed prior to dosing

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 17 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring.

Visit number	V10	V11	V12	V13	V14	V15	V16	V17	V18	V19	Comment
Weeks from randomization	14	16	18	20	22	24	26	28	30	32	
Study day	99	113	127	141	155	169	183	197	211	225	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit							X				
HBV DNA							X				Only for participants who are positive for anti-HBc at screening.
Pharmacokinetic (PK) samples							X				Collect samples before dosing, if dosing is scheduled. Collect additional samples at specified times relative to onset of hypersensitivity events.
CCI							X				Collect samples before dosing, if dosing is scheduled. Collect additional samples at specified times relative to onset of hypersensitivity events.
Flow cytometry panel							X				
Genetics sample											Blood sample for DNA pharmacogenetics can be obtained any time at or after V2.
CCI			X				X				
Endoscopic Procedure											
Endoscopy							X				

Table 3. Schedule of Activities for the Extension Induction and Extension Maintenance Treatment Periods of Study J1P-MC-KFAH

This schedule applies to study participants who are **nonresponders** at Week 12.

V16 procedures may be conducted over more than 1 day as long as all activities are completed within the allowable visit tolerance and all scheduled activities are completed prior to dosing

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 17 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring.

Visit number	V10	V11	V12	V13	V14	V15	V16	V17	V18	V19	Comment
Weeks from randomization	14	16	18	20	22	24	26	28	30	32	
Study day	99	113	127	141	155	169	183	197	211	225	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit							X				
Colon biopsy sample collection							X				Performed at time of endoscopy. Instructions for collection are provided in the lab manual. Flexible sigmoidoscopy is recommended, except for protocol-specified reasons when a colonoscopy should be performed.
Stool Samples											
CCI			X				X				
			X				X				
Randomization and Dosing											
Dosing	X	X	X	X	X	X	X	X	X	X	

Table 3 (continued). Schedule of activities for the Extension Induction and Extension Maintenance Treatment Periods of Study J1P-MC-KFAH

This schedule applies to study participants who are **nonresponders** at Week 12.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 17 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring

Visit number	V20	V21	V22	V23	V24	V25	V26	V27	V28	V29	Comment
Weeks from randomization	34	36	38	40	42	44	46	48	50	52	
Study day	239	253	267	281	295	309	323	337	351	365	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit										X	
Concomitant medications	X	X	X	X	X	X	X	X	X	X	
Adverse events	X	X	X	X	X	X	X	X	X	X	For AESIs, additional data are collected (Section 8.3.6).
Review modified Mayo score (MMS)										X	
Tobacco use										X	
Physical Evaluation											
Weight			X							X	
Vital signs	X	X	X	X	X	X	X	X	X	X	Blood pressure, body temperature, and pulse rate. Vital signs should be measured after participant has been sitting for at least 5 minutes.
Symptom-directed physical examination, including assessment for signs and symptoms of tuberculosis (TB), including peripheral lymph nodes			X							X	See Section 8.2.2.
CCI			X							X	
12-lead electrocardiogram (ECG)			X							X	Locally performed.

Table 3 (continued). Schedule of activities for the Extension Induction and Extension Maintenance Treatment Periods of Study J1P-MC-KFAH

This schedule applies to study participants who are **nonresponders** at Week 12.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 17 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring

Visit number	V20	V21	V22	V23	V24	V25	V26	V27	V28	V29	Comment
Weeks from randomization	34	36	38	40	42	44	46	48	50	52	
Study day	239	253	267	281	295	309	323	337	351	365	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit										X	
Patient Diary (Electronic)											
Diary compliance check	X	X	X	X	X	X	X	X	X	X	Check compliance for Stool Frequency Rectal Bleeding Nocturnal Stools Bristol Stool Scale Urgency NRS Abdominal Pain NRS Fatigue NRS, and Patient Global Rating of Severity (PGR-S).
Diary return										X	
Patient-Reported Outcomes (Electronic)											
Inflammatory Bowel Disease Questionnaire (IBDQ)										X	
Patient Global Impression of Change (PGI-C)										X	
QIDS-SR16										X	
Clinician-Administered Questionnaires (Paper)											
Physician's Global Assessment (PGA)										X	

Table 3 (continued). Schedule of activities for the Extension Induction and Extension Maintenance Treatment Periods of Study J1P-MC-KFAH

This schedule applies to study participants who are **nonresponders** at Week 12.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 17 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring

Visit number	V20	V21	V22	V23	V24	V25	V26	V27	V28	V29	Comment
Weeks from randomization	34	36	38	40	42	44	46	48	50	52	
Study day	239	253	267	281	295	309	323	337	351	365	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit										X	
Laboratory Tests and Sample Collections											
Hematology		X	X	X		X		X		X	
Clinical chemistry		X	X	X		X		X		X	
Lipid panel										X	Fasting samples are preferred, but if participant is nonfasting at time of collection, this is not a protocol deviation.
Urinalysis										X	
Urine pregnancy (local)		X		X		X		X		X	Perform only for women of childbearing potential and women with a history of tubal ligation. At dosing visits which have pregnancy testing, the pregnancy test result must be confirmed to be negative prior to dosing. Pregnancy testing may be performed more frequently than designated on the SoA if local laws or regulations require more frequent testing.
CCI			X							X	
HBV DNA			X					X			Only for participants who are positive for anti-HBc at screening.

Table 3 (continued). Schedule of activities for the Extension Induction and Extension Maintenance Treatment Periods of Study J1P-MC-KFAH

This schedule applies to study participants who are **nonresponders** at Week 12.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 17 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring

Visit number	V20	V21	V22	V23	V24	V25	V26	V27	V28	V29	Comment
Weeks from randomization	34	36	38	40	42	44	46	48	50	52	
Study day	239	253	267	281	295	309	323	337	351	365	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit										X	
Pharmacokinetic (PK) samples			X							X	Collect samples before dosing, if dosing is scheduled. Collect additional samples at specified times relative to onset of hypersensitivity events.
CCI			X							X	Collect samples before dosing, if dosing is scheduled. Collect additional samples at specified times relative to onset of hypersensitivity events.
Flow cytometry panel										X	
Genetics sample											Blood sample for DNA pharmacogenetics can be obtained any time at or after V2.
CCI										X	
Endoscopic Procedure											
Endoscopy										X	
Colon biopsy sample collection										X	Performed at time of endoscopy. Instructions for collection are provided in the lab manual. Flexible sigmoidoscopy is recommended, except for protocol-specified reasons when a colonoscopy should be performed.
Stool Samples											
CCI			X							X	
			X							X	

Table 3 (continued). Schedule of activities for the Extension Induction and Extension Maintenance Treatment Periods of Study J1P-MC-KFAH

This schedule applies to study participants who are **nonresponders** at Week 12.

For procedures at an ETV, see ETV in Table 4.

For procedures at an unscheduled visit, see V997 in Table 4.

Visits 17 through 28 are eligible to be conducted remotely (that is, by telephone, IT-assisted virtual visit, mobile healthcare, or a combination thereof) if written approval is provided by the sponsor and according to the preferences of the participant and the study site. However, no more than 34 days can elapse without an on-site study visit occurring

Visit number	V20	V21	V22	V23	V24	V25	V26	V27	V28	V29	Comment
Weeks from randomization	34	36	38	40	42	44	46	48	50	52	
Study day	239	253	267	281	295	309	323	337	351	365	
Visit interval tolerance (days)	±3	±3	±3	±3	±3	±3	±3	±3	±3	±3	
Fasting visit										X	
Randomization and Dosing											
Dosing	X	X	X	X	X	X	X	X	X		No dosing at V29.

Abbreviations: AESIs = adverse event of special interest; anti-HBc = antibody to hepatitis B core antigen; DNA = deoxyribonucleic acid; ETV = early termination visit; HBV = hepatitis B virus; IT = information technology; NRS = numeric rating scale; QIDS-SR16 = 16-Item Quick Inventory of Depressive Symptomatology–Self Report; SoA = schedule of activities; V = visit.

1.3.4. Schedule of activities for ETV, unscheduled visits, and Post-Treatment Follow-Up Period

Table 4. Schedule of Activities for Early Termination, Unscheduled Visits, and Post-Treatment Follow-Up Period					
V802 is only for randomized participants who were positive for anti-HBc at screening. All other participants will have their final visit at V801 or ETV.					
V997: Additional study procedures can be performed at the discretion of the investigator.					
Visit number	ETV	V997	V801	V802	Comment
Weeks from randomization	—	—	58 or ETV + 8 weeks after last dose	66 or ETV + 16 weeks after last dose	The intervals between ETV and the post-treatment follow-up visit(s) should be adequate to ensure that V801 occurs at least 8 weeks since last dose and V802 (if applicable) occurs at least 16 weeks since last dose.
Study day	—	—	—	—	
Visit interval tolerance (days)	—	—	±7	±7	
Fasting visit					No fasting in this period
Concomitant medications	X	X	X	X	
Adverse events	X	X	X	X	For AESIs, additional data are collected (Section 8.3.6).
Review modified Mayo score (MMS)	X				
Tobacco use	X				
Physical Evaluation					
Weight	X		X		
Vital signs	X	X	X	X	Blood pressure, body temperature, and pulse rate. Vital signs should be measured after participant has been sitting for at least 5 minutes. V997: Vital signs collection is optional.
Symptom-directed physical examination, including assessment for signs and symptoms of tuberculosis (TB), including peripheral lymph nodes	X		X	X	See Section 8.2.2.
CCI	X		X	X	
12-lead electrocardiogram (ECG)	X				Locally performed.
Patient Diary (Electronic)					
Diary return	X				
Patient-Reported Outcomes (Electronic)					
Inflammatory Bowel Disease Questionnaire (IBDQ)	X				
Patient Global Impression of Change (PGI-C)	X				

Table 4. Schedule of Activities for Early Termination, Unscheduled Visits, and Post-Treatment Follow-Up Period					
V802 is only for randomized participants who were positive for anti-HBc at screening. All other participants will have their final visit at V801 or ETV.					
V997: Additional study procedures can be performed at the discretion of the investigator.					
Visit number	ETV	V997	V801	V802	Comment
Weeks from randomization	—	—	58 or ETV + 8 weeks after last dose	66 or ETV + 16 weeks after last dose	The intervals between ETV and the post-treatment follow-up visit(s) should be adequate to ensure that V801 occurs at least 8 weeks since last dose and V802 (if applicable) occurs at least 16 weeks since last dose.
Study day	—	—	—	—	
Visit interval tolerance (days)	—	—	±7	±7	
Fasting visit					No fasting in this period
QIDS-SR16	X				
Clinician-Administered Questionnaires (Paper)					
Physician's Global Assessment (PGA)	X				
Laboratory Tests and Sample Collections					
Hematology	X		X	X	
Clinical chemistry	X		X	X	
Lipid panel	X				Fasting is not required for the lipid panel at ETV.
Urinalysis	X		X	X	
Urine pregnancy (local)	X		X	X	Perform only for women of childbearing potential and women with a history of tubal ligation. At dosing visits which have pregnancy testing, the pregnancy test result must be confirmed to be negative prior to dosing. Pregnancy testing may be performed more frequently than designated on the SoA if local laws or regulations require more frequent testing.
CCI	X				
HBV DNA	X			X	Only for participants who are positive for anti-HBc at screening.
Pharmacokinetic (PK) samples	X		X		Collect samples before dosing, if dosing is scheduled. Collect additional samples at specified times relative to onset of hypersensitivity events.

Table 4. Schedule of Activities for Early Termination, Unscheduled Visits, and Post-Treatment Follow-Up Period					
V802 is only for randomized participants who were positive for anti-HBc at screening. All other participants will have their final visit at V801 or ETV.					
V997: Additional study procedures can be performed at the discretion of the investigator.					
Visit number	ETV	V997	V801	V802	Comment
Weeks from randomization	—	—	58 or ETV + 8 weeks after last dose	66 or ETV + 16 weeks after last dose	The intervals between ETV and the post-treatment follow-up visit(s) should be adequate to ensure that V801 occurs at least 8 weeks since last dose and V802 (if applicable) occurs at least 16 weeks since last dose.
Study day	—	—	—	—	
Visit interval tolerance (days)	—	—	±7	±7	
Fasting visit					No fasting in this period
CCI	X		X		Collect samples before dosing, if dosing is scheduled. Collect additional samples at specified times relative to onset of hypersensitivity events.
Flow cytometry panel	X				
Genetics sample					Blood sample for DNA pharmacogenetics can be obtained any time at or after V2.
CCI	X				
Endoscopic Procedure					
Endoscopy	X				Recommended at ETV based on judgment of the investigator and after discussion with the sponsor's medical monitor. If not performed at ETV, this will not be considered a protocol deviation.
Colon biopsy sample collection	X				Recommended at ETV based on judgment of the investigator and after discussion with the sponsor's medical monitor. Instructions for collection are provided in the lab manual. Flexible sigmoidoscopy is recommended, except for protocol-specified reasons when a colonoscopy should be performed. If not performed at ETV, this will not be considered a protocol deviation.
Stool Samples					
CCI	X				
	X				

Table 4. Schedule of Activities for Early Termination, Unscheduled Visits, and Post-Treatment Follow-Up Period					
V802 is only for randomized participants who were positive for anti-HBc at screening. All other participants will have their final visit at V801 or ETV.					
V997: Additional study procedures can be performed at the discretion of the investigator.					
Visit number	ETV	V997	V801	V802	Comment
Weeks from randomization	—	—	58 or ETV + 8 weeks after last dose	66 or ETV + 16 weeks after last dose	The intervals between ETV and the post-treatment follow-up visit(s) should be adequate to ensure that V801 occurs at least 8 weeks since last dose and V802 (if applicable) occurs at least 16 weeks since last dose.
Study day	—	—	—	—	
Visit interval tolerance (days)	—	—	±7	±7	
Fasting visit					No fasting in this period
Randomization and Dosing					
Dosing					No dosing at ETV, V997, or post-treatment follow-up visits.

Abbreviations: AESIs = adverse event of special interest; anti-HBc = antibody to hepatitis B core antigen; DNA = deoxyribonucleic acid; ETV = early termination visit; HBV = hepatitis B virus; QIDS-SR16 = 16-Item Quick Inventory of Depressive Symptomatology–Self Report; SoA = schedule of activities; V = visit.