

2. Schedule of Activities

Study Schedule Protocol I8F-MC-GPGS

	Screening	Periods 1 and 2 Study Days – at least 35 days washout between Day 1 doses												ED ^c	Comments
Procedure	D-28 to D-2	D-1 ^b	D1 ^b	D2	D3	D4	D5	D6	D7	D8	D15	D21	D36 (±1) ^b		
Informed Consent	X														
Subject Admission to CRU		X													
Tirzepatide Dosing			0 hour												Study drug will be administered after an overnight fast of at least 8 hours.
Subject Discharge from CRU						X									Subjects' CRU stay may be extended at the investigator's discretion for safety monitoring.
Outpatient Visit							X	X	X	X	X	X	X	X	
Medical History and Demographics	X														
Physical Examination /Medical Assessment	X		Predose			X							X	X	Physical examination at screening. Thereafter, targeted reviews and medical assessments as appropriate.
Weight	X		Predose										X	X	
Height	X														
Temperature	X		Predose												
Safety 12-lead ECG ^a	X		Predose	X		X				X			X	X	Single safety ECG will be collected. ECGs must be recorded before collecting any blood samples at each timepoint. Subjects must be supine for approximately 5 to 10 minutes before ECG collection, and remain supine but awake during ECG collection.
Supine Vital Signs (hours)	X		Predose, 12	24	48	72				X	X	X	X	X	Additional time points may be added, if warranted and agreed upon between Lilly and the investigator.
Clinical Laboratory Tests	X		Predose			X				X	X	X	X	X	See Appendix 2 for details. Day 1 predose sample is for baseline only.
Pregnancy Test	X	X											X	X	Female subjects only. Serum pregnancy test will be performed at screening and urine pregnancy tests at subsequent visits.
FSH test, if applicable	X														Female subjects considered to be postmenopausal only. Women with confirmed nonchildbearing potential status can be exempted from further pregnancy tests during the study after screening.
AEs/Concomitant Medications	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Pharmacogenetic Sample ^a			Predose												Single sample in Period 1 only.

	Screening	Periods 1 and 2 Study Days – at least 35 days washout between Day 1 doses												ED ^c	Comments
Procedure	D-28 to D-2	D-1 ^b	D1 ^b	D2	D3	D4	D5	D6	D7	D8	D15	D21	D36 (±1) ^b		
Immunogenicity ^a			Predose								X		X	X	Where applicable, collection times should match with PK sampling time points. In the event of immediate or non-immediate drug hypersensitivity reactions, unscheduled samples will be collected as detailed in Section 9.4.6. Subjects with TE ADA at follow-up/ED will undergo additional follow-up as detailed in Section 9.7.
Blood glucose monitoring (hours) ^a			Predose, 12	24, 36	48	72									Performed using a bedside glucose monitor. Additional unscheduled measurements may be taken at the discretion of the investigator where clinically indicated.
PK Sampling (hours) ^a			Predose, 8, 12	24, 36	48	72	96	120	144	168	336	480	X	X	Up to a 10% deviation from the nominal collection time is permissible as long as actual sampling time is recorded.

Abbreviations: ADA = antidrug antibodies; AE = adverse event; CRU = clinical research unit; D or d = day; ECG = electrocardiogram; ED = early discontinuation; PK = pharmacokinetic(s); TE ADA = treatment-emergent antidrug antibodies.

Note: All sampling times are given relative to dosing (Time 0 hour) with tirzepatide (predose or hours postdose). Unless otherwise indicated, predose procedure may be performed any time prior to dosing.

If multiple procedures take place at the same time point, the following order of the procedures should be used: ECGs, vital signs, PK sample (record of actual PK sampling time is the priority), clinical laboratory sample, immunogenicity, blood glucose and pharmacogenetic sample.

^a Specified times are approximate, and actual times will be recorded. Actual sampling time should not exceed 1 hour prior to dosing for the predose sample.

^b Period 2, Day -1 procedures (pregnancy tests) and some Period 2, Predose procedures (physical examinations, medical assessments, and clinical laboratory assessments) may be omitted if Period 1, D36 (±1) procedure dates occur within 4 days of Period 2 dosing. Period 2, Day 36 (±1) will also be considered as the final follow-up visit.

^c Within 7 to 14 days upon confirmation of early discontinuation.