


1.3. Schedule of Activities (SoA)

Study Procedure	Screening (D-42 to -2)	Treatment Period (Study Days)																			FU/ ED ^b
		-1	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18 ^a	24 (± 2 days)
Informed consent	X																				
Admission to CRU		X																			
Discharge from CRU																				X	
Outpatient visit	X																				X
Medical history and demographics	X																				
Participant eligibility	X	X																			
Rosuvastatin administration			X					X							X						
Pirtobrutinib administration								X	X	X	X	X	X	X	X	X	X	X	X		
Height	X																				
Weight	X																				
Pregnancy test ^c	X	X																		X	
Urine drug screen (including cotinine)	X	X																			
Ethanol test		X																			
Supine vital signs (PR and BP) ^d	X		P, 2 h	24 h				P, 2 h	24 h						P, 2 h	24 h				X	X
Oral body temperature	X																				

Study Procedure	Screening (D-42 to -2)	Treatment Period (Study Days)																			FU/ ED ^b
		-1	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18 ^a	24 (± 2 days)
12-lead ECG	X		P		X			P		X					P		X			X	X
Clinical laboratory tests	X		P		X			P		X					P		X			X	X
Physical examination ^c		X																		X	X
Genetic blood sample for screening	X																				
CCI																					
Rosuvastatin PK samples ^g			P, 0.5, 1, 1.5, 2, 2.5, 3, 4, 5, 6, 8, 12 h	24 h	48 h	72 h	96 h	Pf, 0.5, 1, 1.5, 2, 2.5, 3, 4, 5, 6, 8, 12 h	24 h	48 h	72 h	96 h	120 h		P, 0.5, 1, 1.5, 2, 2.5, 3, 4, 5, 6, 8, 12 h	24 h	48 h	72 h	96 h	120 h	
CCI																					
Adverse event review	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Concomitant medication review	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X


Abbreviations: BP = blood pressure; CRU = clinical research unit; D = day; ECG = electrocardiogram; ED = early discontinuation; FU = follow-up; h = hours postdose; P = predose; PK = pharmacokinetic(s); PR = pulse rate.

- a Participants will be discharged from the CRU on Day 18 once all study procedures are completed.
- b Participants will attend an outpatient follow-up visit 7 ($\pm$ 2) days after the final dose of study intervention. If participants are not able to attend the CRU for this visit, the CRU should contact the participant via phone call to conduct adverse event and concomitant medication review. If applicable, ED assessments should be performed on the day of discontinuation (or as close to it as possible).
- c Female participants only. Serum pregnancy test at screening and urine pregnancy test at all other times.



CCI

- e Complete physical examination at check-in, discharge from CRU, and follow-up. Additional symptom-driven physical examinations may be performed at the discretion of the investigator.



CCI