## 1.3. Schedule of Activities (SoA)

This SoA shows visits and procedures unique to the intervention-specific appendix (ISA) H0P-MC-OA01 (OA01) for LY3016859. Please refer to master protocol H0P-MC-CPMP (CPMP) and the osteoarthritis (OA) disease-state addendum (DSA) H0P-MC-CPMP(1) SoAs for additional information.

| H0P-MC-OA01 ISA | Randomization | Double-Blind Treatment | | | | Follow-up | | | Early Discontinuation | Notes |
|---|---|---|---|---|---|---|---|---|---|---|
| **Visit Number** | **V3** | **V4** | **V5** | **V6** | **V7** | **V801** | **V802** | **V803** | **ED** | |
| **Study Week** | 0 | 2 | 4 | 6 | 8 | 12 | 16 | 26 | | |
| **Visit Window (days)** | | ±3 | ±3 | ±3 | ±3 | ±5 | ±7 | ±7 | | Minimal interval between 2 doses cannot be shorter than 11 days. |
| Physical examination | X | X | X | X | X | X | X | X | X | Targeted exam for skin, eyes, mouth, lungs and GI tract should be performed at each visit, in addition to any symptom driven exam. Full physical exam should be performed at Visit 803 or ED. |
| Weight | | | | | | | | X | | |
| Vital signs | X | X | X | X | X | X | X | X | | Includes respiratory rate, blood pressure, temperature and pulse rate. All vital signs are collected before study intervention infusion on V3-6. |
| ECG | | | X | | X | | | X | | Single safety ECGs |
| Study intervention IV infusion | X | X | X | X | | | | | | 1 hour IV administration<br>The timing between randomization and the first IV infusion may be up to, but not longer than 2 business days to reduce participant burden. |
| Adverse events | | | | | | X | X | X | | |
| Concomitant medication | | | | | | X | X | X | | |
| Rescue medication usage reporting | | | | | | X | | | | If the participant discontinues after V801, this assessment is not needed for ED per the CPMP SoA. |
| NRS | | | | | | X | | | | If the participant discontinues after V801, this assessment is not needed for ED per the CPMP SoA. |
| PGI | | | | | | X | | | | If the participant discontinues after V801, this assessment is not needed for ED per the CPMP SoA. |

| H0P-MC-OA01 ISA | Randomization | Double-Blind Treatment | | | | Follow-up | | | Early Discontinuation | Notes |
|---|---|---|---|---|---|---|---|---|---|---|
| **Visit Number** | **V3** | **V4** | **V5** | **V6** | **V7** | **V801** | **V802** | **V803** | **ED** | |
| **Study Week** | 0 | 2 | 4 | 6 | 8 | 12 | 16 | 26 | | |
| **Visit Window (days)** | | ±3 | ±3 | ±3 | ±3 | ±5 | ±7 | ±7 | | Minimal interval between 2 doses cannot be shorter than 11 days. |
| WOMAC | | | | | | X | | | | If the participant discontinues after V801, this assessment is not needed for ED per the CPMP SoA. |
| C-SSRS | | | | | | X | | X | | Use C-SSRS Since Last Visit version. |
| Self-Harm Supplement Form | | | | | | X | | X | | |
| Self-Harm Follow-up Form | | | | | | X | | X | | Required if triggered by the Self-Harm Supplement Form per instructions. |
| Hematology | | | | | | X | X | X | | |
| Chemistry | | X | | X | | X | X | X | | |
| Urine Chemistries | X | X | X | X | X | X | | X | X | |
| Urinalysis | | | | | | | | X | | |
| Urine pregnancy | X | X | X | X | X | | | X | X | For all premenopausal females or postmenopausal females <50 years of age<br>V3: collect sample before dosing |
| Urine drug screen | X | | X | | X | | | | | |
| Hemoglobin A1c | X | | | | X | | | X | X | |

| H0P-MC-OA01 ISA | Randomization | Double-Blind Treatment | | | | Follow-up | | | Early Discontinuation | Notes |
|---|---|---|---|---|---|---|---|---|---|---|
| **Visit Number** | **V3** | **V4** | **V5** | **V6** | **V7** | **V801** | **V802** | **V803** | **ED** | |
| **Study Week** | 0 | 2 | 4 | 6 | 8 | 12 | 16 | 26 | | |
| **Visit Window (days)** | | ±3 | ±3 | ±3 | ±3 | ±5 | ±7 | ±7 | | Minimal interval between 2 doses cannot be shorter than 11 days. |
| PK sample | X | X | X | X | X | X | X | X | X | V3-V6: collect samples before IV infusion starts and within 15 minutes after IV infusion is complete.<br>V7 – V803: single sample collected at any time.<br>Obtain PK samples from opposite arm used for IV infusion.<br>Record date and time of collection. |
| Immunogenicity (ADA) sample | X | X | X | | X | X | | X | X | V3-6: collect sample before IV infusion starts |
| Epiregulin sample | X | X | X | | X | X | X | X | X | V3-6: collect sample before IV infusion starts.<br>V7-V803: single sample collected at any time to match PK. |
| CCI | | | | | | | | | | |

Abbreviations: ADA = anti-drug antibody; C-SSRS = Columbia Suicide Severity Rating Scale; ECG = electrocardiogram; GI = gastrointestinal; IV = intravenous; NRS = Numeric Rating Scale for pain; PGI = Patient Global Impression of change; PK = pharmacokinetic; V = visit; WOMAC = Western Ontario and McMaster University Arthritis Index.