
1.2. Schema



* Medication washout and preliminary data entry period begins.
** Randomization to either LY3016859 or placebo.
‡ Each participant receives a 1 hour infusion at Visits 3, 4, 5, and 6.
Abbreviations: IV = intravenous; PDEP = preliminary data entry period; V = visit.

1.3. Schedule of Activities (SoA)

This SoA shows visits and procedures unique to the intervention-specific appendix (ISA) H0P-MC-NP01 (NP01) for LY3016859. Please refer to master protocol H0P-MC-CPMP (CPMP) and the diabetic peripheral neuropathic pain (DPNP) disease-state addendum (DSA) H0P-MC-CPMP(3) SoAs for additional information.

H0P-MC-NP01 ISA	Randomization	Double-Blind Treatment				Follow-up			Early Discontinuation	Notes
Visit Number	**V3**	**V4**	**V5**	**V6**	**V7**	**V801**	**V802**	**V803**	**ED**	
Study Week	0	2	4	6	8	12	16	26		
Visit Window (days)		±3	±3	±3	±3	±5	±7	±7		Minimal interval between 2 doses cannot be shorter than 11 days.
Physical examination	X	X	X	X	X	X	X	X	X	Targeted exam for skin, eyes, mouth, lungs and GI tract should be performed at each visit, in addition to any symptom driven exam. Full physical exam should be performed at Visit 803 or ED.
Weight								X		
Vital signs	X	X	X	X	X	X	X	X		Includes respiratory rate, blood pressure, temperature and pulse rate. All vital signs are collected before study intervention infusion on V3-6.
ECG			X		X			X		Single safety ECGs
Study Intervention IV infusion	X	X	X	X						1 hour IV administration The timing between randomization and the first IV infusion may be up to, but not longer than 2 business days to reduce participant burden.
Adverse Events						X	X	X		
Concomitant medication						X	X	X		
Rescue medication usage reporting						X				If the participant discontinues after V801, this assessment is not needed for ED per the CPMP SoA.
NRS						X				If the participant discontinues after V801, this assessment is not needed for ED per the CPMP SoA.
PGI						X				If the participant discontinues after V801, this assessment is not needed for ED per the CPMP SoA.

H0P-MC-NP01 ISA	Randomization	Double-Blind Treatment				Follow-up			Early Discontinuation	Notes
Visit Number	**V3**	**V4**	**V5**	**V6**	**V7**	**V801**	**V802**	**V803**	**ED**	
Study Week	0	2	4	6	8	12	16	26		
Visit Window (days)		±3	±3	±3	±3	±5	±7	±7		Minimal interval between 2 doses cannot be shorter than 11 days.
BPI						X				If the participant discontinues after V801, this assessment is not needed for ED per the CPMP SoA.
C-SSRS						X		X		Use C-SSRS Since Last Visit version.
Self-Harm Supplement Form						X		X		
Self-Harm Follow-up Form						X		X		Required if triggered by the Self-Harm Supplement Form per instructions.
Hematology						X	X	X		
Chemistry		X		X		X	X	X		
Urine Chemistries	X	X	X	X	X	X		X	X	
Urinalysis								X		
Urine pregnancy	X	X	X	X	X			X	X	For all premenopausal females or postmenopausal females <50 years of age V3: collect sample before randomization
Urine drug screen	X		X		X					
Hemoglobin A1c	X				X			X	X	

H0P-MC-NP01 ISA	Randomization	Double-Blind Treatment				Follow-up			Early Discontinuation	Notes
Visit Number	**V3**	**V4**	**V5**	**V6**	**V7**	**V801**	**V802**	**V803**	**ED**	
Study Week	0	2	4	6	8	12	16	26		
Visit Window (days)		±3	±3	±3	±3	±5	±7	±7		Minimal interval between 2 doses cannot be shorter than 11 days.
PK sample	X	X	X	X	X	X	X	X	X	V3-6: collect samples before IV infusion starts and within 15 minutes after IV infusion is complete. V7 – V803: single sample collected at any time. Obtain PK samples from opposite arm used for IV infusion. Record date and time of collection.
Immunogenicity (ADA) sample	X	X	X		X	X		X	X	V3-6: collect sample before IV infusion starts
Epiregulin sample	X	X	X		X	X	X	X	X	V3-6: collect samples before IV infusion starts. V7-V803: single sample collected at any time to match PK.
[redacted]										

Abbreviations: ADA = anti-drug antibody; BPI = Brief pain inventory-modified short form; C-SSRS = Columbia Suicidal Severity Rating Scale; ECG = electrocardiogram; GI = gastrointestinal; IV = intravenous; NRS = Numeric Rating Scale for pain; PGI = Patient Global Impression of Change; PK = pharmacokinetic; V = visit.