1.3. Schedule of Activities (SoA)

This SoA shows visits and procedures unique to the intervention-specific appendix (ISA) H0P-MC-BP01 (BP01) for LY3016859. Please refer to Master Protocol H0P-MC-CPMP (CPMP) and the CLBP disease-state addendum (DSA) H0P-MC-CPMP(2) SoAs for additional information.

H0P-MC-BP01 ISA										
	Randomization	**Double-Blind Treatment**				**Follow-up**			**Early Discontinuation**	**Notes**
Visit Number	**V3**	**V4**	**V5**	**V6**	**V7**	**V801**	**V802**	**V803**	**ED**	
Study Week	0	2	4	6	8	12	16	26		
Visit Window (days)		±3	±3	±3	±3	±5	±7	±7		Minimal interval between 2 doses cannot be shorter than 11 days.
Physical examination	X	X	X	X	X	X	X	X	X	Targeted exam for skin, eyes, mouth, lungs, and GI tract should be performed at each visit, in addition to any symptom-driven exam. Full physical exam should be performed at Visit 803 or ED.
Weight								X		
Vital signs	X	X	X	X	X	X	X	X		Includes respiratory rate, blood pressure, temperature, and pulse rate. All vital signs are collected before study intervention infusion on V3-V6.
ECG			X		X			X		Single safety ECGs
Study intervention IV infusion	X	X	X	X						1-hour IV administration The timing between randomization and the first IV infusion may be up to, but not longer than, 2 business days to reduce participant burden.
Adverse events						X	X	X		
Concomitant medication						X	X	X		
Rescue medication usage reporting						X				If the participant discontinues after V801, this assessment is not needed for ED per the CPMP SoA.
NRS						X				If the participant discontinues after V801, this assessment is not needed for ED per the CPMP SoA.
PGI						X				If the participant discontinues after V801, this assessment is not needed for ED per the CPMP SoA.

H0P-MC-BP01 ISA										
	Randomization	**Double-Blind Treatment**				**Follow-up**			**Early Discontinuation**	**Notes**
Visit Number	**V3**	**V4**	**V5**	**V6**	**V7**	**V801**	**V802**	**V803**	**ED**	
Study Week	0	2	4	6	8	12	16	26		
Visit Window (days)		±3	±3	±3	±3	±5	±7	±7		Minimal interval between 2 doses cannot be shorter than 11 days.
RMDQ						X				If the participant discontinues after V801, this assessment is not needed for ED per the CPMP SoA.
C-SSRS						X		X		Use C-SSRS Since Last Visit version.
Self-Harm Supplement Form						X		X		
Self-Harm Follow-up Form						X		X		Required if triggered by the Self-Harm Supplement Form per instructions.
Hematology						X	X	X		
Chemistry		X		X		X	X	X		
Urine Chemistries	X	X	X	X	X	X		X	X	
Urinalysis								X		
Urine pregnancy	X	X	X	X	X			X	X	For all premenopausal females or postmenopausal females <50 years of age V3: collect sample before dosing.
Urine drug screen	X		X		X					
Hemoglobin A1c	X				X			X	X	

H0P-MC-BP01 ISA										
	Randomization	**Double-Blind Treatment**				**Follow-up**			**Early Discontinuation**	**Notes**
Visit Number	**V3**	**V4**	**V5**	**V6**	**V7**	**V801**	**V802**	**V803**	**ED**	
Study Week	0	2	4	6	8	12	16	26		
Visit Window (days)		±3	±3	±3	±3	±5	±7	±7		Minimal interval between 2 doses cannot be shorter than 11 days.
PK sample	X	X	X	X	X	X	X	X	X	V3-V6: collect samples before IV infusion starts and within 15 minutes after IV infusion is complete. V7-V803: single sample collected at any time. Obtain PK samples from opposite arm used for IV infusion. Record date and time of collection.
Immunogenicity (ADA) sample	X	X	X		X	X		X	X	V3-V6: collect sample before IV infusion starts.
Epiregulin sample	X	X	X		X	X	X	X	X	V3-V6: collect sample before IV infusion starts. V7-V803: single sample collected at any time to match PK.
[redacted]										
[redacted]										

Abbreviations: ADA = anti-drug antibody; C-SSRS = Columbia-Suicide Severity Rating Scale; ECG = electrocardiogram; ED = early discontinuation; GI = gastrointestinal; ISA = intervention-specific appendix; IV = intravenous; NRS = Numeric Rating Scale for pain; PGI = Patient Global Impression of Change; PK = pharmacokinetic; RMDQ = Roland-Morris Disability Questionnaire; SoA = schedule of activities; V = visit.