1.3. Schedule of Activities (SoA)

This SoA shows visits and procedures unique to the intervention-specific appendix (ISA) H0P-MC-OA03 (OA03) for LY3556050. Please refer to master protocol CPMP and the OA DSA CPMP(1) SoAs for additional information.

H0P-MC-OA03 ISA	**Randomization**	**Double-Blind Treatment**				**Early Discontinuation**	**Notes**
Visit Number	**V3**	**V4**	**V5**	**V6**	**V7**	**ED**	
Study Week	0	2	4	6	8		
Visit Window (days)		±3	±3	±3	±3		
Telephone visit	X[a,b]						Visit window +2 days There will be 2 telephone visits between Visit 3 and Visit 4: [a] Visit occurs 4 days after Visit 3 [redacted] [b] Visit occurs 8 days after Visit 3 [redacted]
Procedures							
Evaluate all I/E criteria	X						
Physical examination	X	X	X	X	X	X	V3-V7: Do symptom-directed physical examination
ECG	X	X	X	X	X	X	V3, V5 and V7: Single ECG before in-clinic dosing and at 3 hours after dosing V4, V6 and ED: Single ECG ECG includes collection of PR interval, QRS duration and interval, and heart rate.
Vital signs	X	X	X	X	X	X	**Blood Pressure** V3, V5, and V7: Collect blood pressure and pulse in supine and standing position, before in-clinic dosing and at 3 hours after dosing V4, V6 and ED: Collect blood pressure once Collect orthostatic blood pressure after 5 minutes supine and then 2-3 minutes after standing and collect pulse after supine and standing blood pressure. **Temperature and Respiratory rate** Collect once at each visit

H0P-MC-OA03 ISA	Randomization	Double-Blind Treatment				Early Discontinuation	Notes
Visit Number	**V3**	**V4**	**V5**	**V6**	**V7**	**ED**	
Study Week	0	2	4	6	8		
Visit Window (days)		±3	±3	±3	±3		
Telephone visit	X[a,b]						
Study intervention	X	X	X	X	X		Intervention is taken orally, twice daily (BID), approximately every 12 hours. V3, V5, and V7: Participants take the morning dose in the clinic. Refer to Section 6.6 for dose titration scheme.
Intervention compliance		X	X	X	X	X	
Laboratory Tests and Sample Collection							
Hematology	X	X	X	X	X	X	
Chemistry	X	X	X	X	X	X	
Urine drug screen							Performed at investigator discretion
Serum pregnancy		X	X	X	X	X	For WOCBP only
Urine pregnancy	X						For WOCBP only Collect sample within 24 hours prior to first dose.
Lipid panel	X	X	X	X	X	X	
Thyroid panel	X	X	X	X	X	X	
Cystatin C	X	X	X	X	X	X	

H0P-MC-OA03 ISA	Randomization	Double-Blind Treatment				Early Discontinuation	Notes
Visit Number	**V3**	**V4**	**V5**	**V6**	**V7**	**ED**	
Study Week	0	2	4	6	8		
Visit Window (days)		±3	±3	±3	±3		
Telephone visit	X[a,b]						Visit window +2 days There will be 2 telephone visits between Visit 3 and Visit 4: [a] Visit occurs 4 days after Visit 3 C [redacted] [b] Visit occurs 8 days after Visit 3 C [redacted]
HbA1c					X	X	
PK sample	X	X	X	X	X	X	V3: Collect 3 hours ±15 minutes after dosing. V5 and V7: Collect samples before dosing and 3 hours ±15 minutes after dosing. V4, V6, and ED: Collect single sample at any time. Record date and time of collection.
Somatostatin-related biomarker sample	X	X	X	X	X	X	
CCI [redacted]							
Participant Device							
Participant returns device					X	X	CCI [redacted]

Abbreviations: ECG = electrocardiogram; ED = early discontinuation; HbA1c = glycated hemoglobin; I/E = inclusion and exclusion; ISA = intervention-specific appendix; PK = pharmacokinetic; V = visit; WOCBP = women of childbearing potential.