1.3. Schedule of Activities (SoA)

This SoA shows visits and procedures unique to the intervention-specific appendix (ISA) H0P-MC-NP03 (NP03) for LY3556050. Please refer to master protocol and the DPNP DSA SoAs for additional information.

H0P-MC-NP03 ISA	Randomization	Double-Blind Treatment				Early Discontinuation	Follow-up	Notes
Visit Number	**V3**	**V4**	**V5**	**V6**	**V7**	**ED**	V801	
Study Week	0	2	4	6	8			
Visit Window (days)		±3	±3	±3	±3			
Telephone Visit	X[a,b]						X[c]	Visit window +2 days CCI [redacted] [c] Visit 801 occurs 4-10 days after last dose of study intervention.
Procedures								
Evaluate all I/E criteria	X							
Physical examination	X	X	X	X	X	X		V3-V7: Do symptom-directed physical examination
ECG	X	X	X	X	X	X		V3, V5, and V7: Single safety ECG before in-clinic dosing and at 3 hours after dosing V4, V6 and ED: Single ECG ECG includes collection of PR interval, QRS duration and interval, and heart rate.
Vital signs	X	X	X	X	X	X		**Blood Pressure and Pulse Rate** V3, V5, and V7: Collect blood pressure and pulse in supine and standing position before in-clinic dosing and at 3 hours after dosing V4, V6, and ED: Collect blood pressure once Collect orthostatic blood pressure after 5 minutes supine and then 2-3 minutes after standing and collect pulse after supine and standing blood pressure. **Temperature and Respiratory rate** Collect once at each visit

H0P-MC-NP03 ISA	Randomization	Double-Blind Treatment				Early Discontinuation	Follow-up	Notes
Visit Number	**V3**	**V4**	**V5**	**V6**	**V7**	**ED**	V801	
Study Week	0	2	4	6	8			
Visit Window (days)		±3	±3	±3	±3			
Telephone Visit	X[a,b]						X[c]	Visit window +2 days CCI [redacted] [c] Visit 801 occurs 4-10 days after last dose of study intervention.
Procedures								
AEs							X	
Study intervention	X	X	X	X	X			Intervention is taken orally, twice daily (BID), approximately every 12 hours. V3, V5, and V7: Participants take the morning dose in the clinic. Refer to Section 6.6 for dose titration scheme.
Intervention compliance		X	X	X	X	X		
						Laboratory Tests and Sample Collection		
Hematology	X	X	X	X	X	X		V3: Collect laboratory tests before dosing.
Chemistry	X	X	X	X	X	X		V3: Collect laboratory tests before dosing.
Urine drug screen								Performed at investigator discretion.
Serum pregnancy		X	X	X	X	X		For WOCBP only
Urine pregnancy	X							For WOCBP only Collect sample within 24 hours prior to first dose.
Lipid panel	X	X	X	X	X	X		

H0P-MC-NP03 ISA	Randomization	Double-Blind Treatment				Early Discontinuation	Follow-up	Notes
Visit Number	**V3**	**V4**	**V5**	**V6**	**V7**	**ED**	V801	
Study Week	0	2	4	6	8			
Visit Window (days)		±3	±3	±3	±3			
Telephone Visit	X[a,b]						X[c]	Visit window +2 days CCI [redacted] [c] Visit 801 occurs 4-10 days after last dose of study intervention.
Procedures								
Thyroid panel	X	X	X	X	X	X		
Cystatin C	X	X	X	X	X	X		
HbA1c					X	X		
PK sample	X	X	X	X	X	X		V3: Collect at 3 hours ±15 minutes after dosing. V5 and V7: Collect samples before dosing and at 3 hours ±15 minutes after dosing. V4, V6, and ED: Collect single sample at any time. Record date and time of collection.
Somatostatin-regulated hormone sample	X	X	X	X	X	X		
CCI [redacted]								

H0P-MC-NP03 ISA	Randomization	Double-Blind Treatment				Early Discontinuation	Follow-up	Notes
Visit Number	**V3**	**V4**	**V5**	**V6**	**V7**	**ED**	V801	
Study Week	0	2	4	6	8			
Visit Window (days)		±3	±3	±3	±3			
Telephone Visit	X[a,b]						X[c]	Visit window +2 days CCI [redacted] [c] Visit 801 occurs 4-10 days after last dose of study intervention.
Procedures								
Participant Device								
Participant returns device					X	X		CCI [redacted]

Abbreviations: AE = adverse event; ECG = electrocardiogram; ED = early discontinuation; HbA1c = glycated hemoglobin; I/E = inclusion and exclusion; ISA = intervention-specific appendix; PK = pharmacokinetic; V = visit; WOCBP = women of childbearing potential.