1.3. Schedule of Activities (SoA)

This SoA shows visits and procedures unique to the intervention-specific appendix (ISA) H0P-MC-BP02 (BP02) for LY3526318. Please refer to the CPMP Master Protocol and the CLBP DSA H0P-MC-CPMP(2) SoAs for additional information.

	Randomization to ISA	**Double-Blind Treatment**				**Early Discontinuation**	**Notes**
	V3	**V4**	**V5**	**V6**	**V7**	**ED**	
Study Week	0	2	4	6	8		
Visit Window (days)		±3					
Physical examination	X		X		X	X	Symptom-directed physical exam (see Section 8.2.1) at V5, V7, and ED (if applicable).
Vital signs	X	X	X	X	X	X	At V3, vitals should be taken predose and at approximately 2 and 4 hours post dosing. At V4, V5, V6, and V7, vitals should be taken predose. Pulse rate, respiratory rate, blood pressure, and temperature taken with participant in a sitting position.
ECG			X		X	X	Single ECGs at V5, V7, and ED (if applicable).
Study intervention	X	X	X	X	X		See Section 6.1.
Intervention compliance		X	X	X	X	X	
Clinical Laboratory Tests and Sample Collection							
Hematology		X	X	X	X		
Chemistry		X	X	X	X		
Lipid panel		X	X	X	X		
Urinalysis					X		
Urine pregnancy	X	X	X	X	X	X	For all WOCBP at V3: collect sample before dosing.
Urine drug screen	X		X		X	X	See Section 5.3.2.
HbA1c	X				X	X	

	Randomization to ISA	Double-Blind Treatment				Early Discontinuation	Notes
	V3	V4	V5	V6	V7	ED	
Study Week	0	2	4	6	8		
Visit Window (days)		±3					
PK sample	X	X	X	X	X	X	At V3, samples should be taken at 2 and 4 hours (±15 minutes) post dosing. At V4, V5, V6, and V7, a sample for PK should be taken before the dose is administered; participants should be asked to refrain from taking their dose on V4-V7 until they have their clinic visit and after the blood sample is taken. The date and time of LY3526318 administration prior to the PK sampling will be recorded, i.e., the dose administered in the clinic at V3 and the last at-home dose prior to each of V4, V5, V6, and V7.
CCI							
Participant Device							
Participant returns device					X	X	If a dBM device is used, the participant will return the device to the site at V7 or ED (if applicable).

Abbreviations: CCI ; ECG = electrocardiogram; ED = early discontinuation; HbA1c = hemoglobin A1c; ISA = intervention-specific appendix; PK = pharmacokinetic; V = Visit; WOCBP = women of childbearing potential.