1.3. Schedule of Activities (SoA)

The activities applicable to all disease states and interventions studied under the master protocol are described in the SoA below.

	Screening[a]		Double-Blind Treatment Period						
H0P-MC-CPMP Master Protocol	**Screening Visit for Master**	**Pre-Randomization Screening Visit**	**Randomization to ISA**					**Early Discontinuation**	**Notes**
Visit Number	**V1[b]**	**V2[c]**	**V3**	**V4**	**V5**	**V6**	**V7**	**ED**	Visits after V7 may be specified in the ISA.
Study Week	Up to -6 months	-10 to -7 days for most participants	0	2	4	6	8		
Procedure/ Assessments									
Informed consent for master protocol	X								
Informed consent for ISA	X								ISA-specific informed consent may occur after other V1 procedures and prior to beginning the medication washout if the ISA is not available at the time of initial screening.
Evaluate master protocol I/E criteria	X	X	X						
Pain medication washout		X							Duration of washout is at least 5 half-lives of the longest half-life medication taken for pain. Recommended minimum washout duration is 2 days. If participants are not taking medication for chronic pain during screening, a washout is not required.
PDEP		X							7-day minimum NRS data and rescue use medication collected daily at home post washout and prior to randomization.
Medical history	X								
Family history	X								

	Screening[a]		Double-Blind Treatment Period						
H0P-MC-CPMP Master Protocol	**Screening Visit for Master**	**Pre-Randomization Screening Visit**	**Randomization to ISA**					**Early Discontinuation**	**Notes**
Visit Number	**V1[b]**	**V2[c]**	**V3**	**V4**	**V5**	**V6**	**V7**	**ED**	Visits after V7 may be specified in the ISA.
Study Week	Up to -6 months	-10 to -7 days for most participants	0	2	4	6	8		
Demographics	X								Including age, gender, race, and ethnicity
Physical examination	X	X[d]							
Weight	X	X[d]	X	X	X	X	X	X	
Height	X								
Vital signs	X	X[d]							Pulse rate, respiratory rate, blood pressure, and temperature taken with participant in a sitting position.
ECG	X	X[d]	X					X	Triplicate ECGs. Post randomization ECGs are included in each ISA.
AEs		X	X	X	X	X	X	X	Any events that occur after signing the ICF are considered AEs as defined in Section 10.3.
Preexisting conditions	X								Preexisting conditions not identified at V1 and started prior to signing the ICF should be recorded.
Concomitant medications	X	X	X	X	X	X	X	X	
Scales, Questionnaires, and Outcome Measures									
VAS for pain	X	X	X	X	X	X	X	X	V2 procedure should be repeated if >60 days from V1
NRS for pain		X	X	X	X	X	X	X	Collected daily with a take-home device. Compliance reviewed at each clinic visit.

	Screening[a]		Double-Blind Treatment Period						
H0P-MC-CPMP Master Protocol	**Screening Visit for Master**	**Pre-Randomization Screening Visit**	**Randomization to ISA**					**Early Discontinuation**	**Notes**
Visit Number	**V1[b]**	**V2[c]**	**V3**	**V4**	**V5**	**V6**	**V7**	**ED**	Visits after V7 may be specified in the ISA.
Study Week	Up to -6 months	-10 to -7 days for most participants	0	2	4	6	8		
Rescue medication usage reporting		X	X	X	X	X	X	X	Total quantity from any source, collected daily with a take-home device. Compliance reviewed at each clinic visit. See Section 6.5. for usage guidance.
Pain catastrophizing scale	X								
PGI				X	X	X	X	X	
EQ-5D-5L			X	X	X	X	X	X	
MOS Sleep Scale			X	X	X	X	X	X	
C-SSRS	X	X[d]	X	X	X	X	X	X	At V1, use the C-SSRS Baseline version. All subsequent assessments will use the C-SSRS Since Last Visit version.
Self-Harm Supplement form	X	X[d]	X	X	X	X	X	X	
Self-Harm Follow-Up form	X	X[d]	X	X	X	X	X	X	Complete 1 form for each event identified on the Self-Harm Supplement form.
Clinical Laboratory Tests									
Hematology	X	X[d]	X					X	Post randomization collections will be defined in each respective ISA.

	Screening[a]		Double-Blind Treatment Period						
H0P-MC-CPMP Master Protocol	**Screening Visit for Master**	**Pre-Randomization Screening Visit**	**Randomization to ISA**					**Early Discontinuation**	**Notes**
Visit Number	**V1[b]**	**V2[c]**	**V3**	**V4**	**V5**	**V6**	**V7**	**ED**	Visits after V7 may be specified in the ISA.
Study Week	Up to -6 months	-10 to -7 days for most participants	0	2	4	6	8		
Chemistry	X	X[d]	X					X	Post randomization collections will be defined in each respective ISA.
Lipid panel	X	X[d]	X					X	Post randomization collections will be defined in each respective ISA.
Urinalysis	X	X[d]	X					X	Post randomization collections will be defined in each respective ISA.
B12	X								
Methylmalonic acid	X								
Homocysteine	X								
Serum pregnancy	X	X							For women of childbearing potential. V2 test is optional if within 30 days of V1 procedure
FSH	X	X[d]							
Estradiol	X	X[d]							
TSH	X	X[d]							Post randomization collections will be defined in each respective ISA.
Urine drug screen	X								Collected at screening visit for all participants and by investigator discretion after V1.
Alcohol consumption	X	X[d]	X	X	X	X	X	X	

	Screening[a]		Double-Blind Treatment Period						
H0P-MC-CPMP Master Protocol	**Screening Visit for Master**	**Pre-Randomization Screening Visit**	**Randomization to ISA**					**Early Discontinuation**	**Notes**
Visit Number	**V1[b]**	**V2[c]**	**V3**	**V4**	**V5**	**V6**	**V7**	**ED**	Visits after V7 may be specified in the ISA.
Study Week	Up to -6 months	-10 to -7 days for most participants	0	2	4	6	8		
Recreational drug use	X								
HbA1c	X	X[d]							Post randomization collections will be defined in each respective ISA.
HIV	X	X[d]							
Exploratory biomarker samples			X						Post randomization collections will be defined in each respective ISA.
Pharmacogenetic sample			X						
Participant Questionnaire									
Self-pain positioning questionnaire		X							
Participant Device									
CCI									

Abbreviations: AE = adverse event; C-SSRS = Columbia-Suicide Severity Rating Scale; ECG = electrocardiogram; ED = early discontinuation; EQ-5D-5L = EuroQol-5D 5 level questionnaire; FSH = follicle-stimulating hormone; HbA1c = hemoglobin A1c; HIV = human immunodeficiency virus; ICF = informed consent form; I/E = inclusion and exclusion; ISA = intervention-specific appendix; MOS = Medical Outcomes Study; NRS = numeric rating scale; PDEP = preliminary data entry period; PGI = Patient Global Impression of Change; TSH = thyroid stimulating hormone; V = visit; VAS = visual analog scale.

[a] Screening assessments may be conducted at other time points prior to randomization if they reduce participant burden.

[b] The site determines the half-life of each pain medication the participant is currently taking to schedule Visit 2. Visit 2 can be scheduled no earlier than 7 days prior to randomization at Visit 3 due to the required 7-day PDEP.

[c] The 5-half-life washout period for pain medications must come before the required 7-day PDEP, resulting in a minimum of 10 days for most participants.

[d] Visit 2 testing is optional for individuals who have completed these assessments within 60 days of Visit 1. If the tests are repeated, the most recent results will be used for I/E purposes.