

1.3. Schedule of Activities (SoA)

	Screening	Treatment Period														Follow-Up phone call	Comments
		Days															
Procedure	-28 to -2 days prior to Day 1	-1	1	2	3	4	5	6	7	8	9	10	11	ED		within 7 to 10 days after last dose	
Informed consent	X																
Inclusion and exclusion criteria	X																
Demography	X																
Participant admission to CRU		X															
Participant discharge from CRU													X				A study participant may be required to stay later than Day 11 for management or monitoring of an AE or a clinically significant finding, at the discretion of the investigator.
Outpatient visit	X																

	Screening	Treatment Period													Follow-Up phone call	Comments
		Days														
Procedure	-28 to -2 days prior to Day 1	-1	1	2	3	4	5	6	7	8	9	10	11	ED	within 7 to 10 days after last dose	
Medical history	X															
Medical assessment	X		P							P			X	X		Full medical assessment to be performed at screening, and symptom driven examination to be done at other/any time point at physician's discretion.
Height and weight	X												X	X		Height and weight to be measured at screening; weight only to be measured at Day 11 or ED as appropriate.

	Screening	Treatment Period													Follow-Up phone call	Comments
		Days														
Procedure	-28 to -2 days prior to Day 1	-1	1	2	3	4	5	6	7	8	9	10	11	ED	within 7 to 10 days after last dose	
Serum or urine pregnancy test (women of childbearing potential only)	X	X											X	X		Serum pregnancy tests will be performed at screening and urine pregnancy test at all other time points
Follicle- stimulating hormone (women of non- childbearing potential only)	X															
Human immunodeficiency virus, hepatitis B and C screen	X															
12-lead ECG	X		P, 2h		X					P, 2h		X	X	X		Single ECGs are required.
Vital signs (supine), pulse rate, and blood pressure	X		P, 1, 2h	24h	X	X				P, 1,2h	24h	X	X	X		Time points may be added, if warranted and agreed upon between Lilly and the investigator.
Clinical laboratory tests	X		P	24h		X				P	24h		X	X		See Appendix 10.2, Clinical Laboratory Tests, for details.

	Screening	Treatment Period														Follow-Up phone call	Comments
		Days															
Procedure	-28 to -2 days prior to Day 1	-1	1	2	3	4	5	6	7	8	9	10	11	ED	within 7 to 10 days after last dose		
Genetic sample			X														
CCI																	
Selpercatinib administration			X							X							Participant consumes high-fat meal within 30 minutes prior to dose based on randomization.
Adverse event /Serious adverse event review	X	X	←=====→												X		
Concomitant medication review	X	X	←=====→												X		

Abbreviations: CRU = clinical research unit; ECG = electrocardiogram; ED = early discontinuation; h = hour;

CCI