
o3d12: binned on predicted σ (17537 test atoms)
Mondrian σ_F (λ_g, ARD fit)
CR3 × λ_F (in sample on test)
B1: posterior × q̂
B2: POPS (5 % cut)
B3: CR3, λ = 1
calibrated
observed |ΔF| (eV/Å), bin RMS
predicted per-atom σ (eV/Å), bin RMS


o4d20: binned on predicted σ (17537 test atoms)
Mondrian σ_F (λ_g, ARD fit)
CR3 × λ_F (in sample on test)
B1: posterior × q̂
B2: POPS (5 % cut)
B3: CR3, λ = 1
calibrated
observed |ΔF| (eV/Å), bin RMS
predicted per-atom σ (eV/Å), bin RMS


o3d12: coverage per group (conformal: Mondrian, one quantile; Gaussian: others)
Mondrian σ_F (λ_g, ARD fit)
CR3 × λ_F (in sample on test)
B1: posterior × q̂
B2: POPS (5 % cut)
B3: CR3, λ = 1
one hold-out quantile (ARD fit)
nominal 90%
test coverage of the 90% region
Mondrian group g = 2b + f


o4d20: coverage per group (conformal: Mondrian, one quantile; Gaussian: others)
test coverage of the 90% region
Mondrian group g = 2b + f